

PRIOR AUTHORIZATION POLICY

POLICY: Multiple Sclerosis and Ulcerative Colitis (Oral – Sphingosine 1-Phosphate

Receptor Modulator) - Zeposia Prior Authorization Policy

Zeposia[®] (ozanimod capsules – Celgene)

REVIEW DATE: 07/23/2025

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT WHERE APPROPRIATE AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. WHERE COVERAGE FOR CARE OR SERVICES DOES NOT DEPEND ON SPECIFIC CIRCUMSTANCES, REIMBURSEMENT WILL ONLY BE PROVIDED IF A REQUESTED SERVICE(S) IS SUBMITTED IN ACCORDANCE WITH THE RELEVANT CRITERIA OUTLINED IN THE APPLICABLE COVERAGE POLICY, INCLUDING COVERED DIAGNOSIS AND/OR PROCEDURE CODE(S). REIMBURSEMENT IS NOT ALLOWED FOR SERVICES WHEN BILLED FOR CONDITIONS OR DIAGNOSES THAT ARE NOT COVERED UNDER THIS COVERAGE POLICY (SEE "CODING INFORMATION" BELOW). WHEN BILLING, PROVIDERS MUST USE THE MOST APPROPRIATE CODES AS OF THE EFFECTIVE DATE OF THE SUBMISSION. CLAIMS SUBMITTED FOR SERVICES THAT ARE NOT ACCOMPANIED BY COVERED CODE(S) UNDER THE APPLICABLE COVERAGE POLICY WILL BE DENIED AS NOT COVERED. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Zeposia, a sphingosine 1-phosphate receptor modulator, is indicated for the following uses:¹

- Relapsing forms of multiple sclerosis (MS), in adults, including clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
- **Ulcerative colitis** (UC), in adults with moderately to severely active disease.

Guidelines/Clinical Efficacy

Published guidelines address recommended treatments for the following conditions:

Multiple sclerosis (MS): Zeposia is not currently addressed in MS guidelines.
 In September 2019, a consensus paper was updated by the MS Coalition that discusses the use of disease-modifying therapies in MS.² Many options from

- various pharmacologic classes, involving different mechanisms of action and modes of administration, have shown benefits in patients with MS.
- **Ulcerative colitis (UC):** Zeposia is not currently addressed in UC guidelines. The American Gastroenterological Association (2020) and the American College of Gastroenterology (2019) have clinical practice guidelines on the management of moderate to severe UC and make recommendations for induction and maintenance of remission in adults.^{3,4} Both organizations endorse the use of biologic agents and give specific patient circumstances in the selection for induction and maintenance therapies. The 10-week induction pivotal trial for Zeposia included adults with moderately to severely active UC who had an inadequate response or were intolerant to any of the following agents: oral aminosalicylates, corticosteroids, immunomodulators (e.g., 6-mercaptopurine and azathioprine), or a biologic (e.g., tumor necrosis factor inhibitor, Entyvio[®] [vedolizumab intravenous infusion and subcutaneous injection]).¹

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Zeposia. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Zeposia as well as the monitoring required for adverse events and long-term efficacy, approval requires Zeposia to be prescribed by or in consultation with a physician who specializes in the condition being treated.

• Zeposia® (ozanimod capsules – Celgene) is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

- **1. Multiple Sclerosis.** Approve for the duration noted below if the patient meets ONE of the following (A <u>or</u> B):
 - A) <u>Initial Therapy</u>. Approve for 1 year if the patient meets BOTH of the following (i and ii):
 - i. Patient has a relapsing form of multiple sclerosis; AND Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
 - **ii.** The medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; OR
 - B) Patient is Currently Receiving Zeposia for ≥ 1 Year. Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
 - i. Patient has a relapsing form of multiple sclerosis; AND

<u>Note</u>: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.

- **ii.** Patient meets ONE of the following (a <u>or</u> b):
 - a) Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR

 Note: Examples include stabilization or reduced worsening in disease activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2 lesions]; stabilization or reduced worsening on the Expanded Disability State Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the fatigue symptom and impact questionnaire-relapsing multiple sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or 12-Item MS Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume
 - b) Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND
- **iii.** The medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.
- **2. Ulcerative Colitis.** Approve for the duration noted if the patient meets ONE of the following (A <u>or</u> B):
 - A) <u>Initial Therapy</u>. Approve for 6 months if the patient meets ALL of the following (i <u>and</u> ii):
 - i. Patient is ≥ 18 years of age; AND
 - ii. The medication is prescribed by or in consultation with a gastroenterologist;OR
 - B) <u>Patient is Currently Receiving Zeposia</u>. Approve for 1 year if the patient meets BOTH of the following (i <u>and</u> ii):
 - i. Patient has been established on therapy for at least 6 months; AND <u>Note</u>: A patient who has received < 6 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).
 - ii. Patient meets at least ONE of the following (a or b):
 - a) When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating the requested drug); OR
 - <u>Note</u>: Examples of assessment for inflammatory response include fecal markers (e.g., fecal calprotectin), serum markers (e.g., C-reactive protein), endoscopic assessment, and/or reduced dose of corticosteroids.
 - **b)** Compared with baseline (prior to initiating the requested drug), patient experienced an improvement in at least one symptom, such as

decreased pain, fatigue, stool frequency, and/or decreased rectal bleeding.

CONDITIONS NOT COVERED

- Zeposia® (ozanimod capsules Celgene) is(are) considered not medically necessary for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):
- Concurrent Use with Other Disease-Modifying Agents Used for Multiple Sclerosis. These agents are not indicated for use in combination (see <u>Appendix</u> <u>B</u> for examples). Additional data are required to determine if use of disease-modifying multiple sclerosis agents in combination is safe and provides added efficacy.
- 2. Non-Relapsing Forms of Multiple Sclerosis. The efficacy of Zeposia has not been established in patients with multiple sclerosis with non-relapsing forms of the disease.¹

<u>Note</u>: An example of a non-relapsing form of multiple sclerosis is primary progressive multiple sclerosis.

- **3.** Concurrent Use with a Biologic or with a Targeted Synthetic Oral Small Molecule Drug. This medication should not be administered in combination with another biologic or with a targeted synthetic oral small molecule drug used for an inflammatory condition (see Appendix A for examples). Combination therapy is generally not recommended due to a potentially higher rate of adverse events and lack of controlled clinical data supporting additive efficacy.
- **4. Concurrent Use with Other Potent Immunosuppressants.** In pivotal trials, patients who received Zeposia were not to receive concomitant treatment with non-corticosteroid immunosuppressive or immune-modulating therapies used for the treatment of ulcerative colitis. Combination therapy is generally not recommended due to a potential for a higher rate of adverse effects with combinations and lack of controlled clinical data supporting additive efficacy. Note: Examples of non-corticosteroid immunosuppressive or immune-modulating therapies include 6-mercaptopurine, azathioprine, cyclosporine, and methotrexate.

REFERENCES

- 1. Zeposia® capsules [prescribing information]. Princeton, NJ: Celgene/Bristol Myers Squibb; August 2024
- 2. A Consensus Paper by the Multiple Sclerosis Coalition. The use of disease-modifying therapies in multiple sclerosis. September 2019.
- 3. Feuerstein JD, Isaac s KL, Schneider Y, et al. AGA clinical practice guidelines on the management of moderate to severe ulcerative colitis. *Gastroenterology*. 2020;158:1450-1461.

4. Rubin DT, Ananthakrishnan AN, Siegel CA, et al. American College of Gastroenterology clinical guideline: ulcerative colitis in adults. *Am J Gastroenterol*. 2019;114:384-413.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	No criteria changes.	11/08/2023
Selected Revision	Conditions Not Covered: Concurrent use with a Biologic or with a Targeted Synthetic Oral Small Molecule Drug was changed to as listed (previously oral small molecule drug was listed as Disease-Modifying Antirheumatic Drug). Added Concurrent use with a Potent Immunosuppressant is not allowed.	09/11/2024
Annual Revision	Ulcerative Colitis: For Initial Therapy, it was added in a Note that trial of a mesalamine product does not count as meeting the requirement to try one systemic therapy. Multiple Sclerosis: Ocrevus Zunovo was added to Appendix B as an example of a disease-modifying medication used for multiple sclerosis.	10/09/2024
Early Annual Revision	The name of the policy was changed to add "Oral – Sphingosine 1-Phosphate Receptor Modulator". In addition, the following changes were made: Multiple Sclerosis: Extavia was removed from Appendix B. Ulcerative Colitis: For Initial Therapy, the requirement that the patient has tried ONE systemic agent for ulcerative colitis was removed, along with the related Note. Appendix A was updated to note that biosimilars are available to Stelara, as well as an update to the list of indications.	07/23/2025

APPENDIX A

	Mechanism of Action	Examples of Indications*		
Biologics				
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC		
Cimzia ® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA		
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA, RA		
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC		
Zymfentra [®] (infliximab-dyyb SC injection)	Inhibition of TNF	CD, UC		
Simponi [®] , Simponi Aria [®] (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC		
		IV formulation: AS, PJIA, PsA, RA		
Tocilizumab Products (Actemra® IV, biosimilar; Actemra SC, biosimilar)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA		
		IV formulation: PJIA, RA, SJIA		
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA		
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA IV formulation: JIA, PsA, RA		

Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA^, RA
Omvoh ® (mirikizumab IV infusion, SC injection)	Inhibition of IL-23	CD, UC
Ustekinumab Products (Stelara® SC biosimilars; ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC IV formulation: CD, UC
Siliq® (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection; secukinumab IV infusion)	Inhibition of IL-17A	SC formulation: AS, ERA, nr-axSpA, PsO, PsA IV formulation: AS, nr-axSpA, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Bimzelx ® (bimekizumab-bkzx SC injection)	Inhibition of IL- 17A/17F	PsO
Ilumya® (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi ® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PSA, PsO, UC IV formulation: CD, UC
Tremfya [®] (guselkumab SC injection, guselkumab IV infusion)	Inhibition of IL-23	SC formulation: CD, PsA, PsO, UC IV formulation: CD, UC
Entyvio® (vedolizumab IV infusion, vedolizumab SC injection)	Integrin receptor antagonist	CD, UC

APPENDIX A (CONTINUED)

	Mechanism of Action	Examples of Indications*			
Oral Therapies/Targeted Synthetic Oral Small Molecule Drugs					
Otezla® (apremilast tablets)	Inhibition of PDE4	PsO, PsA			
Cibinqo™ (abrocitinib tablets)	Inhibition of JAK	AD			
	pathways				
Olumiant® (baricitinib tablets)	Inhibition of JAK pathways	RA, AA			
Litfulo ® (ritlecitinib capsules)	Inhibition of JAK pathways	AA			
Leqselvi ® (deuruxolitinib tablets)	Inhibition of JAK pathways	AA			
Rinvoq ® (upadacitinib extended-release tablets)	Inhibition of JAK pathways	AD, AS, nr-axSpA, RA, PsA, CD, UC			
Rinvoq® LQ (upadacitinib oral solution)	Inhibition of JAK pathways	PsA, PJIA			
Sotyktu® (deucravacitinib tablets)	Inhibition of TYK2	PsO			
Xeljanz® (tofacitinib tablets/oral solution)	Inhibition of JAK pathways	RA, PJIA, PsA, UC			
Xeljanz® XR (tofacitinib extended- release tablets)	Inhibition of JAK pathways	RA, PsA, UC			
Zeposia® (ozanimod tablets)	Sphingosine 1 phosphate receptor modulator	UC			
Velsipity ® (etrasimod tablets)	Sphingosine 1 phosphate receptor modulator	UC			

^{*} Not an all-inclusive list of indications. Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn's disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; AA – Alopecia areata; TYK2 – Tyrosine kinase 2.

APPENDIX B

Medication	Mode of Administration
Aubagio® (teriflunomide tablets, generic)	Oral
Avonex® (interferon beta-1a intramuscular injection)	Injection (self-administered)
Bafiertam® (monomethyl fumarate delayed-release capsules)	Oral
Betaseron® (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Briumvi® (ublituximab-xiiy intravenous infusion)	Intravenous infusion
Copaxone® (glatiramer acetate subcutaneous injection,	Injection (self-administered)
generic)	
Gilenya® (fingolimod capsules, generic)	Oral
Glatopa® (glatiramer acetate subcutaneous injection)	Injection (self-administered)
Kesimpta® (ofatumumab subcutaneous injection)	Injection (self-administered)
Lemtrada® (alemtuzumab intravenous infusion)	Intravenous infusion
Mavenclad® (cladribine tablets)	Oral
Mayzent® (siponimod tablets)	Oral
Ocrevus® (ocrelizumab intravenous infusion)	Intravenous infusion

Ocrevus Zunovo [™] (ocrelizumab and hyaluronidase-ocsq subcutaneous injection)	Subcutaneous injection (not self- administered)
Plegridy® (peginterferon beta-1a subcutaneous or intramuscular injection)	Injection (self-administered)
Ponvory® (ponesimod tablets)	Oral
Rebif® (interferon beta-1a subcutaneous injection)	Injection (self-administered)
Tascenso ODT® (fingolimod orally disintegrating tablets)	Oral
Tecfidera® (dimethyl fumarate delayed-release capsules, generic)	Oral
Tyruko® (natalizumab-sztn intravenous infusion)	Intravenous infusion
Tysabri® (natalizumab intravenous infusion)	Intravenous infusion
Vumerity® (diroximel fumarate delayed-release capsules)	Oral
Zeposia® (ozanimod capsules)	Oral

[&]quot;Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.