



PRIOR AUTHORIZATION POLICY

POLICY: Weight Loss – Glucagon-Like Peptide-1 Agonists Prior Authorization Policy

- Saxenda® (liraglutide subcutaneous injection – Novo Nordisk, generic)
- Wegovy® (semaglutide subcutaneous injection – Novo Nordisk)
- Zepbound® (tirzepatide subcutaneous injection – Eli Lilly)

REVIEW DATE: 07/09/2025; selected revision 08/27/2025 and 09/10/2025

INSTRUCTIONS FOR USE

THE FOLLOWING COVERAGE POLICY APPLIES TO HEALTH BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. CERTAIN CIGNA COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES. IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT WHERE APPROPRIATE AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. WHERE COVERAGE FOR CARE OR SERVICES DOES NOT DEPEND ON SPECIFIC CIRCUMSTANCES, REIMBURSEMENT WILL ONLY BE PROVIDED IF A REQUESTED SERVICE(S) IS SUBMITTED IN ACCORDANCE WITH THE RELEVANT CRITERIA OUTLINED IN THE APPLICABLE COVERAGE POLICY, INCLUDING COVERED DIAGNOSIS AND/OR PROCEDURE CODE(S). REIMBURSEMENT IS NOT ALLOWED FOR SERVICES WHEN BILLED FOR CONDITIONS OR DIAGNOSES THAT ARE NOT COVERED UNDER THIS COVERAGE POLICY (SEE "CODING INFORMATION" BELOW). WHEN BILLING, PROVIDERS MUST USE THE MOST APPROPRIATE CODES AS OF THE EFFECTIVE DATE OF THE SUBMISSION. CLAIMS SUBMITTED FOR SERVICES THAT ARE NOT ACCOMPANIED BY COVERED CODE(S) UNDER THE APPLICABLE COVERAGE POLICY WILL BE DENIED AS NOT COVERED. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

CIGNA NATIONAL FORMULARY COVERAGE:

OVERVIEW

Liraglutide (Saxenda, generic), Wegovy, and Zepbound are glucagon-like peptide-1 (GLP-1) receptor agonists; Zepbound is also a glucose-dependent insulinotropic polypeptide (GIP) receptor agonist.^{1,2,9}

Liraglutide, Wegovy, and Zepbound are indicated in combination with a reduced-calorie diet and increased physical activity:^{1,2,9}

- To **reduce excess body weight and maintain weight reduction long term** in:
 - **Liraglutide, Wegovy and Zepbound:** Adults with overweight in the presence of at least one weight-related comorbid condition.^{1,2,9,11}

- **Liraglutide, Wegovy and Zepbound:** Adults with obesity.^{1,9}
- **Liraglutide:** Pediatric patients ≥ 12 years of age and ≥ 60 kg with obesity.²
- **Wegovy:** Pediatric patients ≥ 12 years of age with obesity.^{1,12}

Wegovy is indicated in combination with a reduced-calorie diet and increased physical activity:¹

- To **reduce the risk of major adverse cardiovascular (CV) events (MACE)** [CV death, non-fatal myocardial infarction {MI}, or non-fatal stroke] in adults with established CV disease and either **obesity or overweight**.^{1,10}
- For the treatment of **non-cirrhotic metabolic dysfunction-associated steatohepatitis (MASH)**, formerly known as non-alcoholic steatohepatitis (NASH), with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis) in adults.^{1,25,26}

Zepbound is indicated in combination with a reduced-calorie diet and increased physical activity:⁹

- To treat **moderate to severe obstructive sleep apnea (OSA)** in adults with **obesity**.

Dosing

In the prescribing information for Wegovy, a recommended dose escalation schedule of 16 weeks is outlined (the 2.4 mg dose would be reached at the start of Week 17).¹ If a patient does not tolerate a dose during dose escalation, consider delaying dose escalation for 4 weeks. For CV risk reduction and weight reduction, the maintenance dose of Wegovy is 2.4 mg (recommended) or 1.7 mg injected subcutaneously (SC) once weekly (QW); consider treatment response and tolerability when selecting the maintenance dose. For MASH, the recommended maintenance dose is 2.4 mg QW. If the patient does not tolerate 2.4 mg QW, the dose can be decreased to 1.7 mg QW. Consider re-escalation to 2.4 mg QW.

In the prescribing information for liraglutide, a recommended dose escalation schedule of 4 weeks is outlined.² If a patient does not tolerate an increased dose during dose escalation, consider delaying dose escalation for approximately one additional week. For adults, the recommended maintenance dose of liraglutide is 3 mg SC once daily (QD); discontinue liraglutide if the patient cannot tolerate the 3 mg dose. Additionally, for adults, the prescribing information states to evaluate the change in body weight 16 weeks after initiating liraglutide and discontinue liraglutide if the patient has not lost $\geq 4\%$ of baseline body weight, since it is unlikely the patient will achieve and sustain clinically meaningful weight loss with continued treatment. For pediatric patients, the recommended maintenance dose is 3 mg SC QD; patients who do not tolerate 3 mg QD may have the maintenance dose reduced to 2.4 mg QD. Discontinue liraglutide if the patient cannot tolerate the 2.4 mg dose. If a pediatric patient does not tolerate an increased dose during dose escalation, the dose may be lowered to the prior dose level; dose escalation may take up to 8 weeks. Evaluate the change in body mass index (BMI) after 12 weeks on the maintenance dose in pediatric patients; if the patient has not had a reduction in BMI of $\geq 1\%$ from baseline, discontinue liraglutide as it is unlikely the

patient will achieve and sustain clinically meaningful weight loss with continued treatment.

In the prescribing information for Zepbound, the recommended starting dose is 2.5 mg SC QW.⁹ The 2.5 mg dose is for treatment initiation and is not intended for chronic weight management. After 4 weeks, the dose can be increased to 5 mg QW. The dose can then be increased in 2.5 mg increments, after at least 4 weeks on the current dose. The recommended maintenance doses for weight reduction and long-term maintenance are 5 mg, 10 mg, or 15 mg QW. The recommended maintenance dose in OSA is 10 mg or 15 mg QW. The treatment response and tolerability should be considered when selecting the maintenance dose. If a patient does not tolerate a maintenance dose, consider a lower maintenance dose. The maximum dose is 15 mg QW. The 5 mg, 10 mg, and 15 mg maintenance doses are reached after Week 4, Week 12, and Week 20, respectively.

None of the GLP-1 or GLP-1/GIP agonists are recommended for coadministration with other GLP-1 or GLP-1/GIP agonists.^{1,2,9}

Clinical Efficacy

Secondary Prevention of MACE

SELECT was a randomized, double-blind, placebo-controlled, event-driven study that assessed Wegovy vs. placebo, when added to standard of care, for the secondary prevention of CV events in adults ≥ 45 years of age with BMI ≥ 27 kg/m² and established CV disease without diabetes (n = 17, 604).¹⁰ Established CV disease was defined as one of the following: prior MI, prior stroke (ischemic or hemorrhagic), and/or symptomatic peripheral arterial disease (as evidenced by intermittent claudication with ankle-brachial index < 0.85 , peripheral arterial revascularization procedure, or amputation due to atherosclerotic disease). The trial excluded patients who had received a GLP-1 or GLP-1/GIP agonist within 90 days of enrollment. Patients who developed diabetes during the study remained in the study and received treatment (excluding use of another GLP-1 or GLP-1/GIP agonist). Wegovy was titrated to reach the 2.4 mg QW maintenance dose over 16 weeks. However, if dose escalation led to unacceptable effects, the dose escalation intervals could be extended, treatment could be paused, or maintenance doses < 2.4 mg QW could be used. Very few patients ($< 0.1\%$) were treated with weight-lowering pharmacotherapy at baseline (further detail is not available; however, concomitant GLP-1 agonist use was not allowed).¹¹ The mean hemoglobin A_{1c} (HbA_{1c}) was just over 5.7%; 67% of patients had a HbA_{1c} $\geq 5.7\%$ (pre-diabetes). The most common prior CV event was MI (68% of patients), followed by stroke (18%), and 4.5% of patients had symptomatic peripheral arterial disease; 8% of patients had two or more of these conditions. At baseline, 91.8% of patients were receiving CV risk-lowering pharmacotherapy, 90% of patients were receiving lipid-lowering agents (87.3% of patients were taking statins, 13.0% of patients were taking ezetimibe, 2.7% of patients were taking fibrates, and 2.0% of patients were taking proprotein convertase subtilisin/kexin type 9 inhibitors), 86.2% of patients were receiving platelet aggregation inhibitors, and 12.6% of patients were receiving antithrombotic medications.^{10,11} In addition, 70.2% of patients were taking beta-blockers, 45.0% of patients were taking angiotensin converting enzyme inhibitors,

and 29.5% of patients were taking angiotensin receptor blockers.¹¹ The primary efficacy endpoint was a composite of death from CV causes, non-fatal MI, or non-fatal stroke.¹⁰

Results. Patients were followed for a mean of 39.8 months.¹⁰ At Week 104, approximately 77% of patients receiving Wegovy were taking the target 2.4 mg QW dose (details on the exact proportions of patients on other Wegovy doses are not available; efficacy results are only provided for the 2.4 mg dose). The trial achieved its primary endpoint, demonstrating a statistically significant and superior reduction in MACE for Wegovy vs. placebo. A primary endpoint event occurred in 6.5% vs. 8.0% of patients in the Wegovy vs. placebo groups, respectively (hazard ratio [HR] 0.80; 95% confidence interval [CI]: 0.72, 0.90; $P < 0.001$). Death from CV events, the first confirmatory secondary endpoint, occurred in 2.5% vs. 3.0% of Wegovy- vs. placebo-treated patients, respectively (HR 0.85; 95% CI: 0.71, 1.01; $P =$ not significant for superiority). Because the difference between groups for death from CV events did not meet the required P -value for superiority, testing was not performed for the remaining confirmatory and secondary endpoints.

MASH

The ESSENCE trial (Part 1 $n = 800$), a two-part, ongoing. Phase III, multicenter, double-blind, parallel-group trial randomized adults with MASH and stage F2 to F3 fibrosis to Wegovy or placebo, both in addition to standard of care (optimization of treatment for type 2 diabetes, dyslipidemia, and CV risk management).^{25,26} Results from Part 1 have been published. Eligible patients were ≥ 18 years of age with histological presence of steatohepatitis with stage F2 to F3 fibrosis according to NASH Clinical Research Network classification and a non-alcoholic fatty liver disease (NAFLD) activity score (NAS) of ≥ 4 , with a score of ≥ 1 in each component (steatosis, lobular inflammation, and hepatocyte ballooning), based on central pathologist evaluation of a baseline liver biopsy. Patients with an average alcohol consumption of ≥ 20 grams/day for women or ≥ 30 grams/day for men or alcohol dependence were excluded.^{25,26} Patients taking stable doses of pioglitazone, vitamin E, glucose lowering agent(s), lipid-lowering medication(s), and weight loss medication(s) were allowed to continue these therapies. Rezdiffra[™] (resmetirom tablets) was not approved at the time the trial commenced; therefore, no patients were taking Rezdiffra in Part 1 of this trial. Concomitant use of any other GLP-1 or GLP-1/GIP agonist was not allowed. In patients randomized to Wegovy, the dose was escalated over 16 weeks to reach Wegovy 2.4 mg SC QW. Patients could continue to receive a lower dose of Wegovy if they had unacceptable side effects while receiving the designated target dose and would otherwise discontinue the study.²⁶ The two primary histologic endpoints were: 1) Resolution of steatohepatitis and no worsening of liver fibrosis; and 2) Improvement in liver fibrosis and no worsening of steatohepatitis. In Part 2 of the trial, the primary endpoint will be cirrhosis-free survival at Week 240 (ongoing). Overall, 56% of patients had type 2 diabetes. The mean BMI was 34.6 kg/m² and most patients had a BMI ≥ 35 kg/m² (41.3%); 31.5% of patients had a BMI ≥ 30 kg/m² to < 35 kg/m², 20.5% of patients had a BMI of ≥ 25 kg/m² to < 30 kg/m², and 6.6% of patients had a BMI < 25 kg/m². Most patients fulfilled four (27.8%) or five (43.3%) of five metabolic dysfunction-associated metabolic liver disease (MASLD) cardiometabolic criteria (i.e., BMI ≥ 25 kg/m² [≥ 23 kg/m² Asia] or waist

circumference > 94 cm [male] or > 80 cm [female] or ethnicity adjusted equivalent; fasting serum glucose $\geq$ 100 mg/dL or 2-hour post-prandial glucose $\geq$ 140 mg/dL or type 2 diabetes or treatment for type 2 diabetes; blood pressure $\geq$ 130/85 mmHg or specific antihypertensive drug treatment; plasma high-density lipoprotein cholesterol [HDL-C] $\leq$ 40 mg/dL [male] and $\leq$ 50 mg/dL [female] or lipid-lowering treatment). Most patients had stage F3 fibrosis (68.8%); 31.3% of patients had stage F2 fibrosis. The mean NAS was 5.05 (5.11 in patients with stage F3 fibrosis and 4.92 in patients with stage F2 fibrosis).

Results. At the time of the primary endpoint assessment, 83.5% of patients were taking Wegovy 2.4 mg QW.²⁶ At the interim analysis (the first 800 patients enrolled in the trial), the between-group differences for both primary endpoints were significant for Wegovy vs. placebo. Wegovy demonstrated a significant improvement in liver fibrosis with no worsening of steatohepatitis, as well as resolution of steatohepatitis with no worsening of liver fibrosis. At Week 72, 62.9% vs. 34.1% of patients treated with Wegovy vs. placebo, respectively, achieved resolution of steatohepatitis with no worsening of liver fibrosis (estimated difference 28.7%; 95% CI: 21.1%, 36.2%; $P < 0.001$). In addition, at Week 72, 36.8% vs. 22.4% of patients treated with Wegovy vs. placebo, respectively, had a reduction in liver fibrosis with no worsening of steatohepatitis (estimated difference 14.4%; 95% CI: 7.5%, 21.3%; $P < 0.001$). Confirmatory secondary endpoints also generally favored Wegovy (e.g., resolution of steatohepatitis with improvement in liver fibrosis, weight change). At Week 72, the proportion of patients with both resolution of steatohepatitis and improvement in fibrosis was 32.7% vs. 16.1% of patients receiving Wegovy vs. placebo, respectively (difference 16.5%; 95% CI: 10.2% to 22.8%, respectively; $P < 0.001$). The mean change in body weight was -10.5% vs. -2.0% for Wegovy vs. placebo, respectively (difference -8.5%; 95% CI: -9.6%, -7.4%; $P < 0.001$). Improvements in liver enzymes, non-invasive liver tests, and other metabolic parameters also favored Wegovy. Among patients with F2 fibrosis (Wegovy vs. placebo), 45.0% vs. 22.7% of patients, respectively, had improvement of fibrosis, 37.1% vs. 42.4% of patients, respectively, had no change in fibrosis, and 17.9% vs. 34.8% of patients, respectively, had worsening of fibrosis. Part 2 of the trial is ongoing and expected to read out in 2029.

OSA

The SURMOUNT-OSA ($n = 469$) trials were two 52-week, Phase III, multicenter, double-blind, randomized trials that evaluated the efficacy and safety of maximally tolerated Zepbound (10 mg or 15 mg QW) in adults with obesity (without diabetes) and moderate to severe OSA.¹⁴ Two patient populations were included. In Trial 1, patients were unable or unwilling to use positive airway pressure (PAP) therapy, and in Trial 2, patients had been using PAP therapy for ≥ 3 months at the time of screening and planned to continue PAP therapy during the trial. All patients had a diagnosis of moderate to severe OSA with an apnea-hypopnea index (AHI) ≥ 15 events/hour as diagnosed with polysomnography, home sleep apnea test, or other methods that met local guidelines prior to Visit 1. Patients had a BMI of ≥ 30 kg/m² (≥ 27 kg/m² in Japan) despite the history of at least one self-reported unsuccessful dietary effort to lose weight. Key exclusion criteria were the presence of type 1 or type 2 diabetes ($HbA_{1c} \geq 6.5\%$ at Visit 1), change in weight of > 5 kg

in the past 3 months, planned surgery for sleep apnea or obesity, diagnosis of central or mixed sleep apnea with the percentage of mixed or central apneas/hypopneas $\geq 50\%$, or diagnosis of Cheyne Stokes respiration, diagnosis of obesity hypoventilation syndrome or daytime hypocapnia, active device treatment of OSA other than PAP therapy (e.g., dental appliance), and major craniofacial abnormalities that may affect breathing. In addition, use of medications (prescribed or over-the-counter) or alternative remedies to promote weight loss in the past 3 months was not allowed, this included other GLP-1 or GLP-1/GIP agonists. Of note, although patients with diabetes at baseline were excluded, if a patient developed diabetes while in the study, the patient was referred to their usual care provider. The decision to further evaluate, to initiate antihyperglycemic therapy, and the choice of antihyperglycemic medication was at the discretion of the provider. Following a 4-week screening period, patients were assigned to Trial 1 or Trial 2 and randomly assigned to receive Zepbound or placebo SC QW. All patients received regular lifestyle counseling sessions focused on the maintenance of healthy nutrition, adherence to a 500 calorie/day deficit, and ≥ 150 minutes per week of physical activity. The dose of Zepbound was escalated over a period of up to 20 weeks starting at 2.5 mg SC QW and increased by 2.5 mg every 4 weeks during the dose-escalation period until the patient reached the maximum tolerated dose of 10 mg or 15 mg SC QW at Week 20. Dose modification was permitted for the management of intolerable gastrointestinal (GI) symptoms. Patients who did not tolerate ≥ 10 mg even after one de-escalation and re-escalation attempt were discontinued from the study intervention but remained in the study for continued follow-up. During the first 24 weeks of the treatment period (20-week dose escalation plus 4 weeks), patients unable to tolerate 2.5 mg or 5 mg were discontinued from the study intervention but remained in the study. For patients unable to tolerate any dose escalation between 7.5 mg and 15 mg (inclusive), a dose de-escalation step with subsequent re-escalation by 2.5 mg every 4 weeks up to the maximum tolerated dose was allowed in a blinded fashion, to reach either the 10 mg or 15 mg dose. Only one cycle of dose de-escalation and re-escalation was permitted during the first 24 weeks of the treatment period. The 10 mg maintenance dose was used in patients who tolerated 10 mg, but not 12.5 mg or 15 mg even following one de-escalation and re-escalation attempt. In addition, patients who tolerated 12.5 mg, but not 15 mg even after one de-escalation and re-escalation attempt, continued 10 mg as their maximum tolerated dose. Patients who tolerated 15 mg continued 15 mg as their maximum tolerated dose.

The primary endpoint was the superiority of Zepbound vs. placebo for the change in the AHI from baseline.¹⁴ Several key secondary endpoints were assessed, including the proportion of patients with an AHI reduction of $\geq 50\%$, the proportion of patients with an AHI of < 5 events/hour or with an AHI of 5 to 14 events/hour and a score of ≤ 10 on the Epworth Sleepiness Scale (ESS; scores range from 0 to 24 with higher scores indicating greater daytime sleepiness), percent change in body weight, change in high-sensitivity C-reactive protein (hsCRP), change in sleep apnea specific hypoxic burden, changes in patient-reported outcome measures, and the change in systolic blood pressure. In Trial 1, the mean BMI was 39.1 kg/m² and the mean AHI was 51.5 events/hour. Most patients had severe OSA (63%). In

Trial 2, the mean BMI was 38.7 kg/m² and the mean AHI was 49.5 events/hour. Most patients had severe OSA (68%).

Results. In both trials, Zepbound was superior to placebo for the primary endpoint. In Trial 1, the change in AHI at Week 52 with Zepbound was superior to placebo (-25.3 events/hour [95% CI: -29.3, -21.2] vs. -5.3 events/hour [95% CI: -9.4, -1.1], respectively; estimated treatment difference of -20.0 events/hour; 95% CI: -25.8, -14.2; $P < 0.001$).¹⁴ In Trial 2, the change in AHI at Week 52 with Zepbound was superior to placebo (-29.3 events/hour [95% CI: -33.2, -25.4] vs. -5.5 events/hour [95% CI: -9.9, -1.2], respectively; estimated treatment difference of -23.8 events/hour; 95% CI: -26.9, -17.9; $P < 0.001$). Additionally, patients in both trials who received Zepbound had significant reductions in sleep apnea-specific hypoxic burden. The proportion of patients with a reduction in the AHI of $\geq 50\%$ at Week 52 and the proportion of patients with an AHI of < 5 events/hour or an AHI of 5 to 14 events/hour and an ESS of ≤ 10 at Week 52 also favored Zepbound. Patients receiving Zepbound in both trials had significant reductions in body weight, systolic blood pressure, and hsCRP concentrations as well.

Guidelines

Weight Management

Adult

Guidelines from the American Gastroenterological Association on pharmacological interventions for adults with obesity (2022) state that in adults with obesity or overweight with weight-related complications, who have had an inadequate response to lifestyle interventions, it is recommended to add pharmacological agents to lifestyle interventions over continuing lifestyle interventions alone (strong recommendation, moderate quality evidence).⁶ Wegovy and liraglutide are listed among the therapeutic options. It is also noted that given the magnitude of net benefit, Wegovy may be prioritized over other approved anti-obesity medications for the long-term treatment of obesity for most patients.

Guidelines from the Endocrine Society regarding pharmacological management of obesity (2015) recommend pharmacotherapy as an adjunct to behavioral modification to reduce food intake and increase physical activity for patients with BMI ≥ 30 kg/m² or ≥ 27 kg/m² in the presence of at least one comorbidity, such as hypertension, dyslipidemia, type 2 diabetes, or OSA.³ If a patient's response to a weight loss medication is deemed effective (weight loss $\geq 5\%$ of body weight at 3 months) and safe, it is recommended that the medication be continued. In clinical studies of liraglutide and semaglutide, eligible patients were required to have a prior unsuccessful dietary weight loss attempt. The American Diabetes Association also cites weight loss $\geq 5\%$ of body weight at 3 months as "effective"; when early response is insufficient (typically $< 5\%$ weight loss after 3 months), other therapies should be evaluated.⁸

Per American Association of Clinical Endocrinologists/American College of Endocrinology obesity guidelines (2016), pharmacotherapy for overweight and obesity should be used only as an adjunct to lifestyle therapy and not alone.⁴ The addition of pharmacotherapy produces greater weight loss and weight-loss

maintenance compared with lifestyle therapy alone. The concurrent initiation of lifestyle therapy and pharmacotherapy should be considered in patients with weight-related complications that can be ameliorated by weight loss. Pharmacotherapy should be offered to patients with obesity, when potential benefits outweigh the risks, for the chronic treatment of the disease. Short-term treatment (3 to 6 months) using weight-loss medications has not been demonstrated to produce longer-term health benefits and cannot be generally recommended based on scientific evidence.

Pediatric

Guidelines from the American Academy of Pediatrics on evaluation and treatment of children and adolescents with obesity (2023) note that pediatricians and other primary healthcare providers should offer adolescents ≥ 12 years of age with obesity (BMI $\geq 95^{\text{th}}$ percentile) weight loss pharmacotherapy, according to medication indications, risks, and benefits, as an adjunct to health behavior and lifestyle treatment.⁷

A 2017 Endocrine Society clinical practice guideline on pediatric obesity recommends that pharmacotherapy in combination with lifestyle modification be considered in obese children or adolescents only after failure of a formal program of intensive lifestyle (dietary, physical activity and behavioral) modification to limit weight gain or to ameliorate comorbidities.⁵ The Endocrine Society recommends pharmacotherapy in overweight children and adolescents < 16 years of age only in the context of a clinical trial. Pharmacotherapy should be provided only by clinicians who are experienced in the use of anti-obesity agents and aware of the potential for adverse events. These guidelines recommend limited use of pharmacotherapy because pediatric obesity should be managed preferably as a serious lifestyle condition with important lifelong consequences. The Endocrine Society defines overweight as BMI in at least the 85th percentile but less than the 95th percentile, and obesity as BMI in at least the 95th percentile for age and sex, unless the height velocity is attenuated or inappropriate for the family background or stage of puberty.⁵

MASH

Available recommendations for MASH predate the availability of the published ESSENCE trial data and the approval of Wegovy for MASH. Note: The titles and verbiage within individual publications may not reflect the current MASH or MASLD nomenclature due to the timing of the respective publication. The nomenclature used in this section is reflective of that used within each publication.

The American Association for the Study of Liver Diseases (AASLD) Practice Guidance on the Clinical Management of NAFLD (2023) was updated in October 2024 to address the approval of Rezdiffra for MASLD.²⁷ Some recommendations regarding other therapies (e.g., GLP-1 agonists) are also made. Regardless of treatment, the management of MASLD should include comprehensive lifestyle modification (e.g., nutrition, exercise, and behavior modification) and optimal control of comorbid metabolic conditions. Further, given the comorbidity profile of individuals with MASLD, CV risk management is an important aspect of medical

management. MASH can only be definitively diagnosed by histologic exam (biopsy), in practice, patient selection is based on evidence of steatosis and fibrosis as determined by non-invasive liver disease assessments (NILDAs) in patients with cardiometabolic risk factors without other causes of steatosis, notably, alcohol consumption of > 20 g/day for women and > 30 g/day for men. There are no FDA-approved NILDAs to diagnose MASH with stage F2 to F3 fibrosis or to monitor the response to pharmacotherapy. In general, imaging-based NILDAs such as liver stiffness measurement by vibration-controlled transient elastography (VCTE) or magnetic resonance elastography have better accuracy in assessing fibrosis vs. blood-based tests. In general, magnetic resonance spectroscopy and magnetic resonance imaging proton density fat fraction are considered the most accurate quantitative measures of hepatic steatosis, followed by VCTE-controlled attenuation parameter score and gray scale ultrasound. However, for the purpose of selecting patients for treatment with non-quantitative imaging evidence of hepatic steatosis (e.g., ultrasonographic evidence) in individuals with at least one cardiometabolic risk factor and F2 or F3 fibrosis may be sufficient. Liver biopsy is not typically recommended for fibrosis staging in current clinical practice; however, histologic exam remains the gold standard to quantify fibrosis if performed previously (historical biopsy obtained reasonably recently, e.g., within 3 years). Since NILDAs are more readily available than liver biopsy, it is recommended that more current data (e.g., within 6 to 12 months) be utilized to determine patients who are appropriate candidates for treatment. Regardless of treatment, the management of MASLD should include comprehensive lifestyle modification (e.g., nutrition, exercise, and behavior modification) and optimal control of comorbid metabolic conditions.

The American Diabetes Association (ADA) Standards of Care (2025) provide recommendations for the management of patients with type 2 diabetes and MASLD or MASH.⁸ In adults with type 2 diabetes, MASLD, and overweight or obesity, a GLP-1 agonist (i.e., liraglutide, semaglutide) or GLP-1/GIP agonist (i.e., tirzepatide) with potential benefits in MASH, in addition to healthy interventions for weight loss, is recommended for glycemic management. In adults with type 2 diabetes and MASH or those at high-risk for liver fibrosis (based on non-invasive tests), pioglitazone, GLP-1 agonists, or a GLP-1/GIP agonist is preferred for glycemic management due to potential beneficial effects on MASH.

The ADA Consensus Report on MASLD in People with Diabetes (2025) largely provides recommendations similar to those outlined in the ADA standards of care for patients with type 2 diabetes and MASLD or MASH.²⁸ In patients with at-risk MASH (stage F2 to F3 fibrosis), Rezdiffra may be considered after treatment optimization for obesity and/or diabetes therapies for at least 6 to 12 months. The addition of Rezdiffra should be initiated by a hepatologist or gastroenterologist with expertise in MASH. For the treatment of diabetes, preference should be given to treatments with evidence of safety and effectiveness for steatohepatitis from high-quality trials (e.g., GLP-1 agonists and/or pioglitazone, or GLP-1/GIP agonist), with the dual purpose of treating hyperglycemia and steatohepatitis. Once cirrhosis is established there are no effective treatments. Similar to the AASLD update, the report acknowledges the data are limited for combination use of Rezdiffra with

medications often used for comorbidities in MASLD (e.g., pioglitazone, GLP-1 agonist, or GLP-1/GIP agonist). People with obesity and type 2 diabetes should make it a priority to optimize lifestyle and medical management with a GLP-1 agonist, pioglitazone, or their combination, or a GLP-1/GIP agonist with potential benefits for steatohepatitis.

The American Academy of Clinical Endocrinology Clinical Practice Guideline (co-sponsored by AASLD) [2022] for the Diagnosis and Management of NAFLD in Primary Care and Endocrinology Settings discusses medications that have been proven to be effective for the treatment of liver disease and cardiometabolic conditions associated with NAFLD or NASH.²⁹ The use of obesity pharmacotherapy in adjunct to lifestyle modification for individuals with obesity and NAFLD or NASH with a goal of $\geq 5\%$, and preferably $\geq 10\%$, weight loss is recommended, when this is not effectively achieved by lifestyle modification alone. For chronic weight management in individuals with a BMI ≥ 27 kg/m² and NAFLD or NASH obesity, pharmacotherapy should be considered (preference is given to Wegovy [2.4 mg/week, best evidence] or liraglutide [3 mg/day]). These therapies must be considered as adjuncts to lifestyle modification for individuals with obesity and NAFLD or NASH to promote cardiometabolic health and treat or prevent type 2 diabetes, CV disease, and other end-stage manifestations of obesity.

The American Heart Association Scientific Statement on NAFLD and CV risk (2022) provides several key take-away messages for healthcare providers.³⁰ NASH is recognized as a contributor to, and marker for, increased atherosclerotic CV disease risk. Many risk factors for NAFLD are also risk factors for atherosclerotic CV disease; NAFLD can be considered a risk enhancer when atherosclerotic CV disease risk is assessed. Similar to the other publications, lifestyle intervention is the key therapeutic intervention for patients with NAFLD. Dietary modification, increased physical activity, weight loss, and alcohol avoidance are strongly encouraged. Weight loss pharmacotherapy such as phentermine, Qsymia[®] (phentermine/topiramate extended-release capsules), Contrave[®] (naltrexone/bupropion extended-release tablets), high-dose liraglutide, high-dose semaglutide, and orlistat may be appropriate and effective to achieve sustained weight loss in some patients with BMI > 30 or ≥ 27 kg/m² with comorbidities; however, the role of these agents in the management of NAFLD and NASH is currently undefined. Bariatric surgery should be considered in patients with NAFLD or NASH with BMI > 35 kg/m² as it is the most effective intervention for achieving sustained weight loss in such individuals with a median body weight loss of 21% to 30%.

Sleep Apnea

The American Academy of Sleep Medicine (2017) recommends that diagnostic testing for OSA be performed in combination with a comprehensive sleep evaluation.¹⁵ Polysomnography is the standard diagnostic test for the diagnosis of OSA in adults in whom there is concern for OSA based on the sleep evaluation. Polysomnography is accepted as the gold standard test for the diagnosis of OSA. In some cases, and within the appropriate context, the use of home sleep apnea test as the initial sleep study may be acceptable, however, polysomnography should be

used when home sleep apnea test results do not provide satisfactory posttest probability of confirming or ruling out OSA.

Available treatment guidelines for OSA do not specifically mention the GLP-1 agonists. The American Thoracic Society clinical practice guideline on the role of weight management in the treatment with adults with OSA (2018) recommends that patients with OSA who are overweight or obese ($\text{BMI} \geq 25 \text{ kg/m}^2$) participate in comprehensive lifestyle intervention that includes a reduced calorie diet, exercise/increased physical activity, and behavioral counseling.¹⁶ For patients with OSA and a $\text{BMI} \geq 27 \text{ kg/m}^2$ who have not had an improvement in weight despite a comprehensive weight-loss lifestyle program, and have no contraindications (no active CV disease), evaluation for anti-obesity medication is suggested. The guideline also cites agreement with the American Association of Clinical Endocrinologists and the American College of Endocrinology guidelines (2016)⁴, which state the weight-loss goal in patients with overweight or obesity with OSA should be $\geq 7\%$ to 11% of total body weight.¹⁶ In patients with a $\text{BMI} \geq 35 \text{ kg/m}^2$, referral for bariatric surgery evaluation is suggested.

The American College of Physicians clinical practice guideline for the management of OSA (2013) recommends that all overweight and obese patients diagnosed with OSA be encouraged to lose weight.¹⁷ Continuous PAP is recommended as initial therapy for patients with OSA. Mandibular advancement devices are recommended for patients with OSA who prefer such devices or for those with adverse events associated with continuous PAP treatment.

Clinical success in OSA has been described by several publications. The American Academy of Sleep Medicine (2019) cites a clinically significant threshold of ≥ 15 events/hour (on AHI)¹⁸ and a clinical practice guideline for the treatment of OSA and snoring with oral appliance therapy (2015) from the American Academy of Sleep Medicine and American Academy of Dental Sleep Medicine¹⁹ notes that treatment success is usually defined as a reduction in the AHI to a specific level (e.g., post-treatment AHI < 5 events/hour, or a $> 50\%$ reduction in AHI). Of note, a meta-analysis on the impact of weight reduction on AHI reported that weight reduction in patients with obesity and OSA was associated with improvements in the severity of OSA. A BMI reduction of 20% was associated with an AHI reduction of 57% ; further weight reduction beyond 20% in BMI was associated with a smaller effect on AHI.²⁰

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of liraglutide (Saxenda, generic), Wegovy, and Zepbound. Of note, this policy targets liraglutide (Saxenda, generic), Wegovy, and Zepbound; other glucagon-like peptide-1 agonists that do not carry an FDA-approved indication for weight loss are not targeted in this policy. In the clinical criteria, as appropriate, an asterisk (*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a male, regardless of the individual's gender identity or gender expression; females are defined as individuals

with the biological traits of a female, regardless of the individual's gender identity or gender expression. Because of the specialized skills required for evaluation and diagnosis of patients treated with Wegovy for metabolic dysfunction-associated steatohepatitis (MASH)/non-alcoholic steatohepatitis (NASH) as well as the monitoring required for adverse events and long-term efficacy, approval requires Wegovy for MASH/NASH to be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days.

Documentation: Documentation is required for use of Wegovy for MASH/NASH as noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information.

I. Saxenda® (liraglutide subcutaneous injection – Novo Nordisk, generic) is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

1. Weight Loss in an Adult with Overweight or Obesity. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 4 months if the patient meets ALL of the following (i, ii, iii, and iv):

- i.** Patient is ≥ 18 years of age; AND
- ii.** Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND
- iii.** Patient meets ONE of the following (a or b):
 - a)** At baseline, patient had a BMI ≥ 30 kg/m²; OR
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

- (1)** At baseline, patient had a BMI ≥ 27 kg/m²; AND
- (2)** At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with liraglutide (Saxenda, generic). Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 4 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is ≥ 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI ≥ 30 kg/m²; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI ≥ 27 kg/m²; AND

(2) At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost $\geq 4\%$ of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

2. Weight Loss in a Pediatric Patient with Obesity. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 4 months if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is ≥ 12 years of age and < 18 years of age; AND

ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

iii. At baseline, patient had a BMI $\geq 95^{\text{th}}$ percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-

dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with liraglutide (Saxenda, generic). Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 4 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is ≥ 12 years of age and < 18 years of age; AND

ii. At baseline, patient had a BMI $\geq 95^{\text{th}}$ percentile for age and sex; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has had a reduction in BMI of $\geq 1\%$ from baseline; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

II. Wegovy[®] (semaglutide subcutaneous injection – Novo Nordisk) is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

1. Weight Loss in an Adult with Overweight or Obesity. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 7 months if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is ≥ 18 years of age; AND

ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

iii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI ≥ 30 kg/m²; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI ≥ 27 kg/m²; AND

(2) At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular

disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with Wegovy. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 7 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is ≥ 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI ≥ 30 kg/m²; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI ≥ 27 kg/m²; AND

(2) At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost $\geq 5\%$ of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

2. Weight Loss in a Pediatric Patient with Obesity. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 7 months if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is ≥ 12 years of age and < 18 years of age; AND

- ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND
- iii. At baseline, patient had a BMI $\geq$ 95th percentile for age and sex; AND
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with Wegovy. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 7 months of initial therapy, refer to Initial Therapy criteria above.

- i. Patient is $\geq$ 12 years of age and $<$ 18 years of age; AND
- ii. At baseline, patient had a BMI $\geq$ 95th percentile for age and sex; AND
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iii. Patient has had a reduction in BMI of $\geq$ 1% from baseline; AND
Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound).
- iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

3. Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease with Overweight or Obesity. Approve for 1 year if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve if the patient meets ALL of the following (i, ii, iii, iv, and v):

- i. Patient is $\geq$ 18 years of age; AND
- ii. Patient has a current BMI $\geq$ 27 kg/m²; AND
- iii. Patient meets ONE of the following (a, b, or c):
 - a) Patient has had a prior myocardial infarction; OR
 - b) Patient has had a prior stroke; OR
Note: This does not include a transient ischemic attack (TIA).
 - c) Patient has a history of symptomatic peripheral arterial disease as evidenced by ONE of the following [(1), (2), or (3)]:
 - (1) Intermittent claudication with ankle-brachial index $<$ 0.85; OR
 - (2) Peripheral arterial revascularization procedure; OR
 - (3) Amputation due to atherosclerotic disease; AND
- iv. According to the prescriber, the medication will be used in combination with optimized pharmacotherapy for established cardiovascular disease; AND
- v. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with Wegovy. Approve if the patient meets ALL of the following (i, ii, iii, iv, and v):

Note: For a patient who has not completed 1 year of initial therapy, refer to Initial Therapy criteria above.

i. Patient is ≥ 18 years of age; AND

ii. At baseline, patient had a BMI ≥ 27 kg/m²; AND

Note: This refers to baseline prior to Wegovy.

iii. Patient meets ONE of the following (a, b, or c):

a) Patient has had a prior myocardial infarction; OR

b) Patient has had a prior stroke; OR

c) Patient has a history of symptomatic peripheral arterial disease as evidenced by ONE of the following [(1), (2), or (3)]:

(1) Intermittent claudication with ankle-brachial index < 0.85 ; OR

(2) Peripheral arterial revascularization procedure; OR

(3) Amputation due to atherosclerotic disease; AND

iv. According to the prescriber, the medication will be used in combination with optimized pharmacotherapy for established cardiovascular disease; AND

v. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

4. Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH). Approve for 1 year if the patient meets the ONE of the following (A or B):

A) Initial Therapy: Approve if the patient meets ALL of the following (i, ii, iii, iv, v, vi, vii, and viii):

i. Patient is ≥ 18 years of age; AND

ii. Patient does not have cirrhosis; AND

iii. The diagnosis of MASH/NASH is confirmed by ONE of the following (a or b):

a) Patient has had a liver biopsy AND meets BOTH of the following [(1) and (2)]:

(1) Liver biopsy has been performed within the 3 years preceding treatment with Wegovy **[documentation required]**; AND

(2) Liver biopsy shows non-alcoholic fatty liver disease activity score of ≥ 4 with a score of ≥ 1 in ALL of the following ([i], [ii], and [iii]) **[documentation required]**:

(i) Steatosis; AND

(ii) Ballooning; AND

(iii) Lobular inflammation; OR

b) Patient has had ONE of the following imaging exams performed within the 6 months preceding treatment with Wegovy [(1), (2), or (3)] **[documentation required]**:

(1) Elastography; OR

Note: Examples of elastography include, but are not limited to vibration-controlled transient elastography (e.g., FibroScan), transient elastography, magnetic resonance elastography,

acoustic radiation force impulse imaging, shear wave elastography.

(2) Computed tomography; OR

(3) Magnetic resonance imaging; AND

iv. Patient meets ONE of the following prior to treatment with Wegovy (a or b) **[documentation required]**:

a) Patient has stage F2 fibrosis; OR

b) Patient has stage F3 fibrosis; AND

v. According to the prescriber, the patient has ONE or more of the following metabolic risk factors that are managed according to standard of care (a, b, c, d, e):

a) Central obesity;

b) Hypertriglyceridemia;

c) Reduced high-density lipoprotein cholesterol;

d) Hypertension;

e) Elevated fasting plasma glucose indicative of diabetes or pre-diabetes; AND

vi. According to the prescriber, patient meets ONE of the following (a or b):

a) Female* patient: Alcohol consumption is < 20 grams/day; OR

Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.

b) Male* patient: Alcohol consumption < 30 grams/day; AND

Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.

vii. The medication will be used in combination with appropriate diet and exercise therapy; AND

viii. The medication is prescribed by or in consultation with an endocrinologist, gastroenterologist, or hepatologist; OR

B) Patient is Currently Receiving Wegovy: Approve if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

Note: A patient who has received < 1 year of therapy or who is restarting therapy should be considered under criterion A (Initial Therapy).

i. Patient has completed ≥ 1 year of therapy with Wegovy AND according to the prescriber, the patient has not had worsening of fibrosis or MASH/NASH; AND

Note: This applies to a patient starting their second (or more) year of therapy with Wegovy (i.e., the patient has already completed 1 year or more of therapy with Wegovy).

ii. According to the prescriber, patient has not progressed to stage F4 (cirrhosis); AND

iii. According to the prescriber, metabolic risk factors are managed according to standard of care; AND

iv. According to the prescriber, patient meets ONE of the following (a or b):

a) Female* patient: Alcohol consumption is < 20 grams/day; OR

Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.

b) Male* patient: Alcohol consumption < 30 grams/day; AND

Note: One standard drink (or one alcoholic drink equivalent) contains roughly 14 grams of pure alcohol, which is found in 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of distilled spirits.

v. The medication will be used in combination with appropriate diet and exercise therapy; AND

vi. The medication is prescribed by or in consultation with an endocrinologist, gastroenterologist, or hepatologist.

*Refer to the Policy Statement

III. Zepbound® (tirzepatide subcutaneous injection - Eli Lilly) is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

FDA-Approved Indications

1. Weight Loss in an Adult with Overweight or Obesity. Approve for the duration noted if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve for 8 months if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is ≥ 18 years of age; AND

ii. Patient has engaged in a trial of behavioral modification and dietary restriction for at least 3 months; AND

iii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI ≥ 30 kg/m²; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI ≥ 27 kg/m²; AND

(2) At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

B) Patient is Continuing Therapy with Zepbound. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 8 months of initial therapy, refer to Initial Therapy criteria above.

i. Patient is ≥ 18 years of age; AND

ii. Patient meets ONE of the following (a or b):

a) At baseline, patient had a BMI ≥ 30 kg/m²; OR

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

b) Patient meets BOTH of the following [(1) and (2)]:

(1) At baseline, patient had a BMI ≥ 27 kg/m²; AND

(2) At baseline, patient had, or patient currently has, at least ONE of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, cardiovascular disease, knee osteoarthritis, asthma, chronic obstructive pulmonary disease, metabolic dysfunction-associated steatotic liver disease/non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iii. Patient has lost $\geq 5\%$ of baseline body weight; AND

Note: This refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., liraglutide [Saxenda, generic], Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).

iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

2. Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity.

Approve for 1 year if the patient meets ONE of the following (A or B):

A) Initial Therapy. Approve if the patient meets ALL of the following (i, ii, iii, iv, and v):

i. Patient is ≥ 18 years of age; AND

ii. Patient has a current BMI ≥ 30 kg/m²; AND

iii. Patient has had a sleep study that shows BOTH of the following (a and b):

a) Patient has been diagnosed with moderate to severe obstructive sleep apnea; AND

b) Patient has an apnea-hypopnea index ≥ 15 events per hour; AND

Note: A diagnosis of moderate obstructive sleep apnea is an apnea-hypopnea index of ≥ 15 events per hour and a diagnosis of severe sleep apnea is an apnea-hypopnea index ≥ 30 events per hour. The apnea-

hypopnea index is the number of apneas and hypopneas during 1 hour of sleep.

- iv. The patient does NOT meet either of the following (a or b):

Note: A patient who has one or more of the following conditions/diagnoses below is not approved.

- a) Central sleep apnea with percent of central apneas/hypopneas $\geq 50\%$;
OR

- b) Cheyne Stokes respiration; AND

- v. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet; OR

- B) Patient is Continuing Therapy with Zepbound.** Approve if the patient meets ALL of the following (i, ii, iii, and iv):

Note: For a patient who has not completed 1 year of initial therapy, refer to Initial Therapy criteria above.

- i. Patient is ≥ 18 years of age; AND

- ii. At baseline, patient had a BMI ≥ 30 kg/m²; AND

Note: This refers to baseline before Zepbound.

- iii. Patient has completed ≥ 1 year of therapy with Zepbound AND the patient meets BOTH of the following (a and b):

- a) Patient has lost $\geq 10\%$ of baseline body weight; AND

Note: This refers to baseline prior to Zepbound.

- b) According to the prescriber, patient has stability in obstructive sleep apnea signs or symptoms; AND

Note: Examples of signs or symptoms of obstructive sleep apnea include but are not limited to snoring, excessive daytime sleepiness, fatigue.

- iv. The medication will be used concomitantly with behavioral modification and a reduced-calorie diet.

CONDITIONS NOT COVERED

- **Saxenda® (liraglutide subcutaneous injection – Novo Nordisk, generic)**
- **Wegovy® (semaglutide subcutaneous injection – Novo Nordisk)**
- **Zepbound® (tirzepatide subcutaneous injection - Eli Lilly)**

is(are) considered not medically necessary for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

1. Concomitant Use with Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/ Glucose-Dependent Insulinotropic Polypeptide (GIP) Agonists.

The GLP-1 agonists and the GLP-1/GIP agonist should not be combined with each other or with any other GLP-1 agonists or GLP-1/GIP agonist.^{1,2,9,12,20-26}

There are other GLP-1 agonist and GLP-1/GIP agonist products not included in this policy that are FDA-approved for type 2 diabetes, and not for chronic weight management. Note: Examples of other GLP-1 agonists include but are not limited to, exenatide subcutaneous (SC) injection, Ozempic (semaglutide SC

injection), Rybelsus (semaglutide tablets), Trulicity (dulaglutide SC injection), and liraglutide SC injection (Victoza, generic). An example of a GLP-1/GIP agonist is Mounjaro (tirzepatide SC injection).

REFERENCES

1. Wegovy® subcutaneous injection [prescribing information]. Plainsboro, NJ: Novo Nordisk; August 2025.
2. Saxenda® subcutaneous injection [prescribing information]. Plainsboro, NJ: Novo Nordisk; May 2025.
3. Apovian CM, Aronne LJ, Bessesen DH, et al; Endocrine Society. Pharmacological management of obesity: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2015;100(2):342-62.
4. Garvey WT, Mechanick JI, Brett EM, Garber AJ, Hurley DL, Jastreboff AM, Nadolsky K, Pessah-Pollack R, Plodkowski R; Reviewers of the AACE/ACE Obesity Clinical Practice Guidelines. American Association of Clinical Endocrinologists and American College of Cardiology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract*. 2016;22 Suppl 3:1-203.
5. Styne DM, Arslanian SA, Connor EL, Farooqi IS, Murad MH, Silverstein JH, Yanovski JA. Pediatric Obesity-Assessment, Treatment, and Prevention: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2017;102(3):709-757.
6. Grunwald E, Shah R, Hernaez R, et al; AGA Clinical Guidelines Committee. AGA Clinical Practice Guideline on Pharmacological Interventions for Adults with Obesity. *Gastroenterology*. 2022;163(5):1198-1225.
7. Hampl SE, Hassink SG, Skinner AC, et al. Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents with Obesity. *Pediatrics*. 2023;151(2):e2022060640.
8. American Diabetes Association. Standards of care in diabetes – 2025. *Diabetes Care*. 2025;48(Suppl 1):S1-S352.
9. Zepbound® subcutaneous injection [prescribing information]. Indianapolis, IN: Eli Lilly; April 2025.
10. Lincoff AM, Brown-Frandsen K, Colhoun HM, et al; for the SELECT Trial Investigators. Semaglutide and cardiovascular outcomes in obesity without diabetes. *N Engl J Med*. 2023;389(24):2221-2232.
11. Lingvay I, Brown-Frandsen K, Colhoun HM et al. Semaglutide for cardiovascular event reduction in people with overweight or obesity: SELECT study baseline characteristics. *Obesity*. 2023;31(1):111-122.
12. Wilding JPH, Batterham RL, Calanna S, et al; STEP 1 Study Group. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med*. 2021;384(11):989.
13. Weghuber D, Barrett T, Barrientos-Pérez M, et al; STEP TEENS Investigators. Once-weekly semaglutide in adolescents with obesity. *N Engl J Med*. 2022;387(24):2245-2257.
14. Malhorta A, Grunstein RR, Fietze I, et al; for the SURMOUNT-OSA Investigators. Tirzepatide for the treatment of obstructive sleep apnea and obesity. *N Engl J Med*. 2024;391(13):1193-1205.
15. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical practice guideline for diagnostic testing for adult obstructive sleep apnea: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2017;13(3):479-504.
16. Hudgel DW, Patel SR, Ahasic AM, et al; on behalf of the American Thoracic Society Assembly on Sleep and Respiratory Neurology. The role of weight management in the treatment of adult obstructive sleep apnea. *Am J Respir Crit Care Med*. 2018;198(6):e70-e87.
17. Qaseem A, Hotly JEC, Owens DK, et al; for the Clinical Guidelines Committee of the American College of Physicians. Management of obstructive sleep apnea in adults: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2013;159:471-483.
18. Ramar K, Dort LC, Katz SG et al. Clinical practice guideline for the treatment of obstructive sleep apnea and snoring with oral appliance therapy: an update for 2015. An American Academy of Sleep Medicine and American Academy of Dental Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2015;11(7):773-827.

19. Patil SP, Ayappa IA, Caples SM, et al. Treatment of adult obstructive sleep apnea with positive airway pressure: An American Academy of Sleep Medicine Systematic Review, Meta-analysis, and GRADE Assessment. *J Clin Sleep Med*. 2019;15(2):301-334.
20. Malhorta A, Heilman CR, Banerjee, et al. Weight reduction and the impact on apnea-hypopnea index: a systematic meta-analysis. *Sleep Medicine*. 2023;121:26-31.
21. Jasterboff AM, Aronne LJ, Ahmad NN, et al; for the SURMOUNT-1 Investigators. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med*. 2022;387(3):205-216.
22. Garvey TW, Frias JP, Jasterboff, et al; for the SURMOUNT-2 Investigators. Tirzepatide once weekly for the treatment of obesity in people with type 2 diabetes (SURMOUNT-2): a double-blind, randomized, multicenter, placebo-controlled, phase 3 trial. *Lancet*. 2023;402(10402):613-6262.
23. Wadden TA, Chao AM, Machineni, et al. Tirzepatide after intensive lifestyle intervention in adults with overweight or obesity: The SURMOUNT-3 phase 3 trial. *Nature Med*. 2023;29(11):2909-2918.
24. Wadden TA, Hollander P, Klein S, et al; on behalf of the NN8022-1923 Investigators. Weight maintenance and additional weight loss with liraglutide after low-calorie-diet-induced weight loss: The SCALE Maintenance randomized study. *Int J Obes*. 2013;37:1443-1451.
25. Newsome PN, Sanyal AJ, Engerbretsen KA, et al. Semaglutide 2.4 mg in participants with metabolic dysfunction-associated steatohepatitis: baseline characteristics and design of the phase 3 ESSENCE trial. *Aliment Pharmacol Ther*. 2024;60(11-12):1525-1533.
26. Sanyal AJ, Newsome PN, Kliers I, et al. Phase 3 trial of semaglutide in metabolic dysfunction-associated steatohepatitis. *N Engl J Med*. 2025;392(21):2089-2099.
27. Chen VL, Morgan TR, Rotman Y, et al. Resmetirom therapy for metabolic dysfunction-associated steatotic liver disease: October 2024 updates to AASLD Practice Guidance. *Hepatology*. 2025;81(1):312-320.
28. Cusi K, Abdelmalek MF, Apovian CM, et al. Metabolic dysfunction-associated steatotic liver disease (MASLD) in people with diabetes: The need for screening and early intervention. A consensus report of the American Diabetes Association. *Diabetes Care*. 2025;48(7):1057-1082.
29. Cusi K, Isaacs S, Barb D, et al. American Association of Clinical Endocrinology clinical practice guideline for the diagnosis and management of nonalcoholic fatty liver disease in primary care and endocrinology settings. Co-sponsored by the American Association for the Study of Liver Diseases (AASLD). *Endocrine Pract*. 2022;28:528-562.
30. Duell PB, Welty FK, Miller M, et al; on behalf of the American Heart Association Council on Arteriosclerosis, Thrombosis and Vascular biology; Council on Hypertension; Council on the Kidney in Cardiovascular Disease; Council on Lifestyle and Cardiometabolic Health; and Council on Peripheral Vascular Disease. Nonalcoholic fatty liver disease and cardiovascular risk: a scientific statement from the American Heart Association. *Arterioscler Thromb Vasc Biol*. 2022;42:e168-e185.

HISTORY

Type of Revision	Summary of Changes	Review Date
Annual Revision	No criteria changes.	07/12/2023
Selected revision	<u>Wegovy</u> Weight Loss, Adult: Continuation criteria were updated to reflect the new approved maintenance dose of Wegovy (1.7 mg once weekly) in adults. The continuation criterion that approves continuation of Wegovy for 1 year, was modified to approve if the patient is able to tolerate a Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly. The continuation criterion that approves continuation of Wegovy for up to 5 months, was modified to approve if according to the prescriber, the patient is continuing to titrate the Wegovy dose to a target of 1.7 mg weekly or 2.4 mg once weekly. Other conditions of coverage still apply for continued approval of Wegovy.	07/26/2023

Selected revision	<u>Wegovy</u> Weight Loss, Adult: In the initial therapy criteria, the requirement for a current BMI ≥ 30 kg/m² or ≥ 27 kg/m² and at least one of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, or cardiovascular disease was modified to require that at baseline (prior to the initiation of Wegovy), the patient had a BMI ≥ 30 kg/m² or ≥ 27 kg/m² and at least one of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, or cardiovascular disease. Weight Loss, Pediatric: In the initial therapy criteria, the requirement for a current BMI $\geq 95^{\text{th}}$ percentile for age and sex was modified to require that at baseline (prior to the initiation of Wegovy), patient had a BMI $\geq 95^{\text{th}}$ percentile for age and sex. <u>Saxenda</u> Weight Loss, Adult: In the initial therapy criteria, the requirement for a current BMI ≥ 30 kg/m² or ≥ 27 kg/m² and at least one of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, or cardiovascular disease was modified to require that at baseline (prior to the initiation of Saxenda), the patient had a BMI ≥ 30 kg/m² or ≥ 27 kg/m² and at least one of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, or cardiovascular disease. Weight Loss, Pediatric: In the initial therapy criteria, the requirement for a current BMI $\geq 95^{\text{th}}$ percentile for age and sex was modified to require that at baseline (prior to the initiation of Saxenda), patient had a BMI $\geq 95^{\text{th}}$ percentile for age and sex.	09/13/2023
Selected revision	Zepbound was added to the policy. New criteria were created for this product. Initial approval is for 8 months, continuation approval is for 1 year (up to 4 months if still titrating). <u>Saxenda</u> Weight Loss, Adult: Initial Therapy: Baseline body mass index (BMI) criteria were modified to remove the requirement that the BMI is prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound). <u>Patient is Continuing Therapy with Saxenda:</u> Baseline BMI criteria were modified to remove the requirement that the BMI is prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound). The criterion that a patient has lost $\geq 4\%$ of baseline weight was modified to remove the requirement that baseline body weight was prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound). Weight Loss, Pediatric: Initial Therapy: The baseline BMI criterion was modified to remove the requirement that the BMI is prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulinotropic polypeptide (GIP) receptor agonist (e.g., Zepbound). <u>Patient is Continuing Therapy with Saxenda:</u> The baseline BMI criterion was modified to remove the requirement that the BMI is prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g.,	11/15/2023

	Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). The criterion that a patient has lost $\geq 1\%$ of baseline weight was modified to remove the requirement that baseline body weight was prior to initiation of Saxenda. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). <u>Wegovy</u> Weight Loss, Adult: Initial Therapy: Baseline BMI criteria were modified to remove the requirement that the BMI is prior to initiation of Wegovy. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). <u>Patient is Continuing Therapy with Wegovy:</u> Baseline BMI criteria were modified to remove the requirement that the BMI is prior to initiation of Wegovy. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). The criterion that a patient has lost $\geq 5\%$ of baseline weight was modified to remove the requirement that baseline body weight was prior to initiation of Wegovy. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). Weight Loss, Pediatric: Initial Therapy: The baseline BMI criterion was modified to remove the requirement that the BMI is prior to initiation of Wegovy. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound).	
Selected Revision	Weight Loss, Adult: Patient is Continuing Therapy with Wegovy: The baseline BMI criterion was modified to remove the requirement that the BMI is prior to initiation of Wegovy. A note was added that baseline refers to baseline prior to any glucagon-like peptide-1 (GLP-1) agonist (e.g., Saxenda, Wegovy) or GLP-1/glucose-dependent insulintropic polypeptide (GIP) receptor agonist (e.g., Zepbound). The criterion that a patient has lost $\geq 1\%$ of baseline weight was modified to remove the requirement that baseline body weight was prior to initiation of Wegovy. <u>Conditions Not Covered</u> Concomitant Use with other Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/Glucose-Dependent Insulintropic Polypeptide (GIP) Receptor Agonists. GLP-1/GIP receptor agonists were added to this condition not recommended for approval.	11/15/2023
Selected Revision	<u>Saxenda, Wegovy, and Zepbound</u> Weight Loss, Adult: Initial Therapy and Patient is Continuing on Therapy: The criterion for a patient with a BMI ≥ 27 kg/m² and at least one of the following weight-related comorbidities: hypertension, type 2 diabetes, dyslipidemia, obstructive sleep apnea, or cardiovascular disease was modified to expand the list of comorbid conditions to include knee osteoarthritis, asthma, chronic obstructive pulmonary disease, non-alcoholic fatty liver disease, polycystic ovarian syndrome, or coronary artery disease.	01/31/2024
DEU Revision	The revised and new indication for Wegovy was added to the overview of the document.	03/25/2024

Selected Revision	<u>Wegovy</u> Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease who is Either Obese or Overweight. A new condition of coverage was added to FDA-approved indications for Wegovy.	04/03/2024
Selected Revision	<u>Saxenda, Wegovy, and Zepbound</u> Weight Loss, Adult: <u>Initial Therapy and Patient is Continuing on Therapy:</u> Metabolic-dysfunction associated steatotic liver disease (new nomenclature for non-alcoholic fatty liver disease) was added to the list of one of the weight-related comorbidities for a patient with a BMI ≥ 27 kg/m ² . Additionally, for the one or more weight-related comorbidity, the criterion was modified to state that the comorbidity is at baseline or current.	05/08/2024
Annual Revision	No criteria changes. Concomitant Use with Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/ Glucose-Dependent Insulinotropic Polypeptide (GIP) Agonists. This condition not recommended for approval was reworded. Previously, the condition read "Concomitant Use with Other Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/Glucose Dependent Insulinotropic Polypeptide (GIP) Receptor Agonists.	07/17/2024
DEU Revision	10/24/2024: Updated Zepbound indication added to overview.	--
Selected Revision	<u>Saxenda</u> Weight Loss, Adult. <u>Patient is Continuing on Therapy with Saxenda:</u> Criterion that required the patient was able to tolerate the Saxenda maintenance dose of 3 mg once daily was removed. Weight Loss, Pediatric. <u>Patient is Continuing on Therapy with Saxenda:</u> Criterion that required the patient was able to tolerate the Saxenda maintenance dose of 2.4 mg once daily or 3 mg once daily was removed. <u>Wegovy</u> Weight Loss, Adult. <u>Patient is Continuing on Therapy with Wegovy:</u> Criteria related to dosing were removed. Specifically, the criteria that required the patient was able to tolerate the Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly OR if the patient had received < 12 consecutive months of Wegovy and was continuing to titrate the Wegovy dose to a target of 1.7 mg once weekly or 2.4 mg once weekly, according to the prescriber, was removed. The approval duration was changed to 1 year for a patient continuing on therapy with Wegovy; previously the approval duration was 1 year if the patient was able to tolerate the Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly and up to 5 months if the patient was continuing to titrate to the Wegovy target dose of 1.7 mg or 2.4 mg once weekly. Weight Loss, Pediatric. <u>Patient is Continuing on Therapy with Wegovy:</u> Criteria related to dosing were removed. Specifically, the criteria that required the patient was able to tolerate the Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly OR if the patient had received < 12 consecutive months of Wegovy and was continuing to titrate the Wegovy dose to a target of 1.7 mg once weekly or 2.4 mg once weekly, according to the prescriber, was removed. The approval duration was changed to 1 year for a patient continuing on therapy with Wegovy; previously the approval duration was 1 year if the patient was able to tolerate the Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly and up to 5 months if the patient was continuing to titrate to the Wegovy target dose of 1.7 mg or 2.4 mg once weekly.	01/08/2025

	Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease who is Either Obese or Overweight. <u>Initial Therapy.</u> The criterion requiring that the patient has a BMI ≥ 27 kg/m² was clarified to state that the patient has a “current” BMI ≥ 27 kg/m². <u>Patient is Continuing Therapy with Wegovy:</u> The criterion that required the patient was able to tolerate the Wegovy maintenance dose of 1.7 mg once weekly or 2.4 mg once weekly was removed. <u>Zepbound</u> Weight Loss, Adult. <u>Patient is Continuing Therapy with Zepbound:</u> Criteria related to dosing were removed. Specifically, the criteria that required the patient was able to tolerate the Zepbound maintenance dose of 5 mg, 10 mg, or 15 mg once weekly OR if the patient had received < 12 consecutive months of Zepbound and was continuing to titrate the Zepbound dose to a target of 10 mg once weekly or 15 mg once weekly, according to the prescriber, was removed. The approval duration was changed to 1 year for a patient continuing on therapy with Zepbound; previously the approval duration was 1 year if the patient was able to tolerate the Zepbound maintenance dose of 5 mg, 10 mg, or 15 mg once weekly and up to 4 months if the patient was continuing to titrate to the Zepbound target dose of 10 mg or 15 mg once weekly. Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity. A new FDA-approved indication was added to the Policy.	
Selected Revision	<u>Wegovy:</u> Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease who is Either Obese or Overweight. <u>Initial Therapy.</u> For the requirement that a patient has had a prior stroke, a note was added to clarify that this does not include a transient ischemic attack (TIA). <u>Zepbound:</u> Obstructive Sleep Apnea, Moderate to Severe, in a Patient with Obesity. <u>Initial Therapy.</u> The requirement that a patient had a sleep study was modified to remove the timeframe that the sleep study was within the past 1 year. A patient is still required to have a sleep study. Conditions Not Covered : Concomitant Use with Other Medications FDA-Approved for Weight Loss. This condition not recommended for approval was clarified to state that concomitant use with other medications <u>FDA-approved</u> for weight loss is not recommended. Previously, the requirement did not specify medications were “FDA-approved” for weight loss. The note with examples of weight loss medications was updated to reflect product availability for medications FDA-approved for weight loss. Concomitant Use with Glucagon-Like Peptide-1 (GLP-1) Agonists or GLP-1/ Glucose-Dependent Insulinotropic Polypeptide (GIP) Agonists. The note was updated to reflect availability for other GLP-1 or GLP-1/GIP agonists.	05/28/2025

Annual Revision	<u>Saxenda:</u> Weight Loss in an Adult with Overweight or Obesity. This condition of approval was modified to add "with overweight or obesity". Weight Loss in a Pediatric Patient with Obesity. This condition of approval was modified to add "with obesity". <u>Wegovy:</u> Weight Loss in an Adult with Overweight or Obesity. This condition of approval was modified to add "with overweight or obesity". Weight Loss in a Pediatric Patient with Obesity. This condition of approval was modified to add "with obesity". Major Adverse Cardiovascular Event(s) Risk Reduction in a Patient with Established Cardiovascular Disease in a Patient with Overweight or Obesity. This condition of approval was reworded from "in an overweight or obese patient" to "in a patient with overweight or obesity". <u>Zepbound:</u> Weight Loss in an Adult with Overweight or Obesity. This condition of approval was modified to add "with overweight or obesity".	07/09/2025
Selected Revision	Policy Statement: The following was added to the Policy Statement: In clinical criteria, as appropriate, an asterisk (*) is noted next to the specified gender. In this context, the specified gender is defined as follows: males are defined as individuals with the biological traits of a male, regardless of the individual's gender identity or gender expression; females are defined as individuals with the biological traits of a female, regardless of the individual's gender identity or gender expression. Because of the specialized skills required for evaluation and diagnosis of patients treated with Wegovy for (MASH)/non-alcoholic steatohepatitis (NASH) as well as the monitoring required for adverse events and long-term efficacy, approval requires Wegovy for MASH/NASH to be prescribed by or in consultation with a physician who specializes in the condition being treated. Documentation: A requirement for documentation was added for the use of Wegovy for metabolic dysfunction-associated steatohepatitis MASH/NASH. Documentation may include, but is not limited to, chart notes, laboratory results, medical test results, claims records, prescription receipts, and/or other information. All documentation must include patient-specific identifying information. <u>Wegovy:</u> Metabolic Dysfunction-Associated Steatohepatitis (MASH)/Non-Alcoholic Steatohepatitis (NASH). A new condition of approval was added to FDA-Approved Indications. Conditions Not Covered : Concomitant Use with Other Medications FDA-Approved for Weight Loss. This condition not recommended for approval was removed.	08/27/2025

Selected Revision	Liraglutide, generic to Saxenda, was added to the policy. Policy Statement: The Policy Statement was updated as follows to address the availability of liraglutide, generic to Saxenda: Prior Authorization is recommended for prescription benefit coverage of liraglutide (Saxenda, generic), Wegovy, and Zepbound. Of note, this policy targets liraglutide (Saxenda, generic), Wegovy, and Zepbound; other glucagon-like peptide-1 agonists that do not carry an FDA-approved indication for weight loss are not targeted in this policy. <u>Liraglutide (Saxenda, generic), Wegovy, and Zepbound:</u> Weight Loss in an Adult with Overweight or Obesity: <u>Initial Therapy and Patient is Continuing on Therapy:</u> The notes that define baseline were updated to include liraglutide, generic to Saxenda. <u>Liraglutide (Saxenda, generic) and Wegovy:</u> Weight Loss in a Pediatric Patient with Obesity: <u>Initial Therapy and Patient is Continuing on Therapy:</u> The notes that define baseline were updated to include liraglutide, generic to Saxenda.	09/10/2025
-------------------	--	------------

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.