



Medical Coverage Policy

Effective Date09/15/2025

Next Review Date5/15/2026

Coverage Policy Number..... 0053

Hyperbaric and Topical Oxygen Therapies

Table of Contents

Overview	2
Coverage Policy	2
Health Equity Considerations.....	4
General Background.....	4
Medicare Coverage Determinations	25
Coding Information	25
References.....	26
Revision Details.....	40

Related Coverage Resources

[Autism Spectrum Disorders/Pervasive Developmental Disorders: Assessment and Treatment](#)
[Bone Growth Stimulators: Electrical \(Invasive\), Ultrasound](#)
[Complementary and Alternative Medicine](#)
[COVID-19 Drug and Biologic Therapeutics](#)
[COVID-19: In Vitro Diagnostic Testing](#)
[Infertility Services](#)
[Negative Pressure Wound Therapy/Vacuum-Assisted Closure \(VAC\) for Nonhealing Wounds](#)
[Compression Devices](#)
[Tissue-Engineered Skin Substitutes](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only

be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses the use of systemic hyperbaric oxygen (HBO) therapy (HBOT), also called hyperbaric oxygen therapy (HOT), and topical oxygen therapy (TOT). Systemic HBO is proposed for the treatment of multiple conditions and involves the inhalation of 100% oxygen in a single or multiplace chamber. Topical oxygen therapy (TOT) delivers 100% oxygen to a localized area (e.g., over a wound).

Coverage Policy

Systemic hyperbaric oxygen therapy (HBO/HBOT/HOT) in single or multiplace chambers is considered medically necessary first-line treatment for ALL of the following conditions:

- acute carbon monoxide poisoning
- air or gas embolism
- decompression sickness
- exceptional blood loss when transfusion is not an option

Systemic hyperbaric oxygen therapy (HBO/HBOT/HOT) in single or multiplace chambers is considered medically necessary adjunctive treatment for ALL of the following conditions:

- acute cyanide poisoning, after administration of antidote
- acute traumatic peripheral ischemia/insufficiency (e.g., crush injuries, compartment syndrome, suturing of severed limbs)
- avascular necrosis
- central retinal artery occlusion
- clostridial myositis and myonecrosis (i.e., gas gangrene)
- compromised skin grafts and flaps (i.e., preexisting grafts or flaps that are showing signs of failure or necrosis)
- delayed osteoradionecrosis, including pre- and post-dental extraction(s) from an irradiated mandible
- idiopathic sudden sensorineural hearing loss (ISSHL) within four weeks of symptom onset
- intracranial abscess
- necrotizing soft tissue infections (e.g., necrotizing fasciitis, Meleney's ulcer)
- osteomyelitis unresponsive to conventional medical and surgical interventions
- radiation-induced cystitis or hemorrhagic cystitis (i.e., resulting from chemolytic response, graft-versus-host disease [GVHD])

- soft tissue radionecrosis, delayed (e.g., radiation-induced enterocolitis, proctitis, brain necrosis)
- thermal burns, acute, requiring inpatient hospitalization
- Wagner grade III or higher diabetic wounds/ulcers of the lower extremities that have failed standard wound therapy

Systemic hyperbaric oxygen therapy in single or multiplace chambers is considered not medically necessary for ANY of the following conditions:

- actinomycosis
- acute cerebral edema
- acute coronary syndrome (ACS)/myocardial ischemia/infarction (MI), cardiogenic shock/preconditioning for coronary artery bypass graft surgery
- acute or chronic cerebral vascular insufficiency
- acute thermal and chemical pulmonary damage (i.e., smoke inhalation with pulmonary insufficiency)
- acute wound, flap, and/or graft
- anorectal disorders (e.g., chronic anal fissure [CAF], internal hemorrhoids, infectious proctitis)
- autism spectrum disorders
- brain injury, closed head injury, traumatic brain injury (TBI), anoxic encephalopathy
- brown recluse spider bites
- cancer
- carbon tetrachloride poisoning
- cerebral palsy
- cerebral radionecrosis
- chronic fatigue syndrome
- chronic peripheral vascular insufficiency
- COVID-19
- Crohn's disease
- cutaneous decubitus/pressure ulcers
- dementia
- epilepsy
- fractures, acute, delayed union or nonunion
- headaches (e.g., cluster, migraine)
- hepatic necrosis
- human immunodeficiency virus (HIV)-fatigue
- in vitro fertilization
- Lyme disease
- lymphedema
- malignant otitis externa (e.g., necrotizing external otitis)
- multiple sclerosis
- mycoses
- nonvascular causes of chronic brain syndrome (e.g., Pick's disease, Alzheimer's disease, Korsakoff's disease)
- ophthalmologic conditions other than central retinal artery occlusion (e.g., optic neuropathy, glaucoma)
- organ storage
- organ transplantation
- penile glans necrosis
- pulmonary emphysema
- reflex sympathetic dystrophy/complex regional pain syndrome
- rheumatoid arthritis

- sepsis
- sickle cell disease
- soft tissue injury (e.g., delayed onset muscle soreness, sprains, strains)
- spinal cord injury
- stroke
- tetanus
- tinnitus
- venous stasis ulcers

Topical oxygen therapy (TOT) is considered medically necessary as an adjunct treatment of a diabetic foot ulcer that has failed to heal with optimal standard of care.

Topical oxygen therapy (TOT) is considered not medically necessary for any other indication.

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Diabetic foot ulcers (DFUs) are a common and serious complication in individuals with diabetes. These ulcers frequently become infected, significantly contribute to the risk of amputation, and are associated with increased mortality. Although systematic reviews rarely include detailed demographic data or conduct subgroup analyses, one review (Sun, et al., discussed below) did note that male participants outnumbered females by a ratio of 2.95 to 1.

General Background

Systemic hyperbaric oxygen therapy (HBO/HBOT/HOT) involves the inhalation of 100% oxygen under increased atmospheric pressure (e.g., 2 to 3 atmospheres absolute [ATA]) (Undersea and Hyperbaric Medicine Society [UHMS], 2023). A hyperbaric oxygen chamber (whether single or multiplace chamber [i.e., created to hold several people]) is a device intended to promote the movement of oxygen from the environment to the patient's tissues by means of pressurization. Forcing oxygen into the tissues, organs, brain, and fluids of the body is proposed to stimulate cell growth and regeneration, displace toxins and impurities, and stimulate the immune system. Treatment sessions may last for 30–120 minutes and may be given for up to five times per week. Some conditions may only require one or two treatments (e.g., cyanide poisoning) while others may require 10–40 treatments (e.g., osteonecrosis) depending on the severity of the illness and the clinical response of the patient (i.e., complete response occurs, or no improvement is being seen).

Topical oxygen therapy (TOT) consists of the direct application of oxygen to a wound site. Although TOT is sometimes referred to as topic hyperbaric oxygen, the kits/devices used for TOT apply oxygen to the wound site at slightly above atmospheric pressure. As noted above, systemic hyperbaric oxygen involves the inhalation of 100% oxygen under increased atmospheric pressure.

U.S. Food and Drug Administration (FDA): Mono- and multiplace hyperbaric chambers are approved by the FDA as a Class II, 510(k) device. Examples of these chambers include the OxyHeal 1000 Monoplace Hyperbaric Chambers (OxyHeal Health Group, LaJolla, CA) and the Multiplace Hyperbaric Chambers (Makai Marine Industries, Inc., Boca Raton, FL). The devices were approved for the treatment of the conditions recommended by the Undersea Hyperbaric Medicine Society at that time (FDA, 2005; FDA, 2004).

Various topical hyperbaric oxygen systems have received FDA clearance through the 510(k) process, including the following:

- Hyper-Box Topical Wound Oxygen System (AOTI, Inc. Qualtech House, Gateway, Ireland), includes either a reusable chamber or single use sleeve that encloses the limb and is connected to a controller unit and oxygen source.
- O2 Boot™ (GWR Medical, Inc., Chadds Ford, PA), a single use disposable device applied over the wound and sealed with a hypoallergenic adhesive, to deliver oxygen directly to the wound site
- Natrox Topical Oxygen Delivery System with IODP (Inotec AMD Ltd, Fairfax, VA), a wearable, portable device consisting of an oxygen generator, oxygen delivery system, and rechargeable batteries. It provides continuous oxygen delivery to the wound site.
- EPIFLO-28 (Neogenix, LLC dba Ogenix, Beachwood, Ohio), a small silent disposable oxygen concentrator that delivers oxygen to the wound via a long sterile tube.

Each of the above devices are indicated for the treatment of skin ulcerations due to diabetes, venous stasis, post-surgical infections and gangrenous lesions; decubitus ulcers; amputations/infected stumps; skin grafts; burns; and frostbite.

Systemic Hyperbaric Oxygen

Literature Review - HBO as Primary Therapy: Evidence in the published peer-reviewed literature and professional society guidelines support the safety and effectiveness of HBO as a primary treatment option for acute carbon monoxide poisoning; air or gas embolism; decompression sickness; and exceptional blood loss when transfusion is not an option (Undersea & Hyperbaric Medical Society [UHMS], 2023; Bennett, et al., 2012).

Literature Review - HBO as Adjunctive Therapy: HBO has been shown to be effective and is an established adjunctive therapy used in combination with other established therapies for the treatment of acute cyanide poisoning; acute traumatic peripheral ischemia/insufficiency (e.g., crush injuries, compartment syndrome, suturing of severed limbs); central retinal artery occlusion, clostridial myositis and myonecrosis (i.e., gas gangrene); compromised skin grafts and flaps (i.e., preexisting grafts or flaps that are showing signs of failure or necrosis); intracranial abscess; necrotizing soft tissue infections such as necrotizing fasciitis or Meleney's ulcer; osteomyelitis that is unresponsive to conventional medical and surgical interventions; delayed radiation damage of non-neurologic tissue (i.e., osteoradionecrosis, including pre- and post-dental extraction in an irradiated mandible, and mandibular radionecrosis), soft tissue radionecrosis (e.g., radiation-induced enterocolitis, cystitis, proctitis; laryngeal and brain necrosis) and acute thermal burns requiring hospitalization (Lin, et. al., 2023; UHMS, 2023; Bennett, et al., 2012; Eskes, et al., 2011; Nabil and Samman, 2011; Fritz, et al., 2010; Goldman, 2009).

HBO is also a recognized adjunctive therapy for the treatment of radiation-induced cystitis or hemorrhagic cystitis resulting from chemolytic response or graft-versus-host disease, and

radiation-induced enterocolitis (Cardinal, et al., 2018; Fink, 2006; Chong, 2005; Fine, 2005; El-Zimaity, 2004; Lazzarini, 2004; Hailey, 2003; Wang, 2003; Kalayoglu-Besisik, 2003; Cesaro, 2003).

Randomized controlled trials and prospective case series support the safety and efficacy of HBO as an effective adjunctive therapy for the treatment of Wagner grades III–V diabetic wounds/ulcers of the lower extremity that are refractory to aggressive medical management including wound care, glucose control and surgical debridement or surgical revascularization. A Wagner grade III wound involves a deep ulcer that contains an abscess, osteomyelitis, or both; grade IV is an ulcer that has led to gangrene of the toes and/or forefoot; and a grade V ulcer has caused gangrene of the entire foot or enough of the foot that it cannot be salvaged. Most study protocols utilizing hyperbaric oxygen for diabetic wound healing exposed patients daily at 2–2.5 ATA for 90–120 minutes for a total of 20–40 sessions (Huang, et al., 2015; Kranke, et al., 2015; Weaver, 2014; Goldman, 2009; Roeckl-Wiedmann, et al., 2005).

Sudden sensorineural hearing loss (SSHL) is an acute hearing impairment defined as a 30 decibel (dB) or greater hearing loss occurring in at least three contiguous audiometric frequencies over 72 hours or less. With sudden hearing loss, the loss is typically defined in relation to hearing in the opposite ear because pre-event audiometry is generally not available. Idiopathic means that there is no identifiable cause of the sudden hearing loss and 85%–90% of SHL is idiopathic. SHL is considered an emergency situation that requires immediate medical intervention. The standard treatment is systemic and/or intratympanic corticosteroids. Patients refractory to initial therapy may be given the addition of HBO as an adjuvant (Weber, 2024; UHMS, 2011). HBO is FDA approved for the treatment of “hearing loss (complete hearing loss that occurs suddenly and without any known cause)” (FDA, 2021). RCT and systematic reviews with small patient populations reported that HBO was an effective adjunctive therapy for the treatment of SSHL (Tong, et al., 2021; Cvorovic et al., 2013; Bennett et al., 2012). Therefore, HBO has evolved into an accepted treatment option for a small subset of patients.

Avascular necrosis (AVN), also called osteonecrosis or aseptic necrosis, is a disease in which there is a lack of blood supply to the bone causing death of bone tissue. Ultimately, AVN may lead to collapse of the bone and joint surface. AVN most often occurs in the hip joint in the femoral head and usually leads to osteoarthritis. Risk factors include hip injury, alcohol abuse and/or excessive corticosteroid use. AVN may be associated with other disease entities (e.g., Gaucher disease, sickle cell disease) and in some cases there may be no underlying disease (idiopathic AVN). Treatment depends on the severity of symptoms and may include limited weight bearing, physical therapy, cessation of alcohol usage, and/or surgical intervention. HBO has been proposed for the treatment of AVN. Studies are primarily in the form of randomized control trials (RCTs), systematic reviews, case series and retrospective reviews with small patient populations (n=12–109) and reported that HBO was an effective adjunctive therapy for the treatment of AVN (Paderno, et al., 2021; Li et al., 2017; Uzun et al. 2016; Camporesi et al., 2010; Reis, et al., 2003). Therefore, HBO has evolved into an accepted treatment option for a small subset of patients.

Professional Societies: The American College of Emergency Physicians (ACEP) (2017) recommended the use of HBO therapy or high flow normobaric therapy for acute carbon monoxide poisoned patients. However, it remains unclear whether HBO therapy is superior to normobaric oxygen therapy for improving long-term neurocognitive outcomes.

The Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine suggested HBO for the management of diabetic foot ulcers in patients who fail to respond to 4–6 weeks of conservative management. The clinical guideline

noted that patients should be selected for this therapy carefully considering the cost and the burden of prolonged daily treatment (Society for Vascular Surgery, 2016).

The Undersea and Hyperbaric Medical Society (UHMS) (2023) approved the following indications for systemic HBO:

- air or gas embolism
- carbon monoxide poisoning
- carbon monoxide poisoning complicated by cyanide poisoning
- central retinal artery occlusion
- clostridial myositis and myonecrosis (gas gangrene)
- crush injury, compartment syndrome, and other acute traumatic ischemias
- decompression sickness
- enhancement of healing in select problem wounds
- exceptional blood loss (severe anemia)
- intracranial abscess
- necrotizing soft tissue infections
- osteomyelitis (refractory)
- delayed radiation injury (soft tissue and bony necrosis)
- skin grafts and flaps (compromised)
- thermal burns (acute)
- idiopathic sudden sensorineural hearing loss
- avascular necrosis (aseptic osteonecrosis)

The Undersea and Hyperbaric Medical Society (2011) support HBO for the treatment of ISSHL stating that patients who meet the criteria for ISSHL may benefit from HBO. Candidates for HBO include those patients with “moderate to profound ISSHL (≥ 41 dB) who present within 14 days of symptom onset”. According to the UHMS, patients presenting after this time may experience improvement when treated with HBO, however, the medical literature suggests that early intervention is associated with improved outcomes. The best evidence supports the use of HBO within two weeks of symptom onset.

The American Academy of Otolaryngology – Head and Neck Surgery (AAO-HNS) (2012, updated 2019) developed evidence-based clinical practice guidelines for the management of sudden hearing loss with a special emphasis on managing sudden sensorineural hearing loss (SSNHL) in adult patients (aged 18 years and older). Following a systematic review of randomized controlled trials “with methodological limitations”, AAO-HNS stated that HBO, is an option when combined with steroid therapy in SSNHL as primary therapy within two weeks of onset of symptoms and as salvage therapy when used within 4 weeks of onset, with potentially more benefit noted in cases of severe to profound loss”. AAO-HNS’s stated that although the level of evidence was “modest and imprecise”, it was “sufficient to promote greater awareness of HBO for the treatment of SSNHL”. The report went on to say that the results should be interpreted with caution due to the small number of patients in the trials, methodological shortcomings, and poor reporting. The authors also noted that HBO is not recognized by many United States clinicians as an intervention for ISSNHL.

Literature Review - Other Proposed Indications for Systemic HBO: There is insufficient evidence in the published peer-reviewed scientific literature to support HBO as a primary or adjunctive treatment of the conditions discussed below (this list may not be all inclusive). HBO is not FDA approved for these other indications.

Actinomycosis: Actinomycosis is a rare chronic, indolent, suppurative, tissue-destructive infection presenting with lumps and sinus formation, usually involving the head and neck,

although it can affect other parts of the body, such as the abdomen and thorax. Adjunctive HBO has been proposed as a treatment option for patients who are unresponsive to medical and surgical intervention; however, studies are primarily in the form of case reports.

Acute Cerebral Edema: Cerebral edema accompanies a wide variety of pathologic processes and may be present in head/brain injury, stroke, brain tumor, cerebral infections (e.g., brain abscess, encephalitis and meningitis), lead encephalopathy, hypoxia, disequilibrium syndrome associated with dialysis and diabetic ketoacidosis, Reye's syndrome, fulminant hepatic encephalopathy, and hydrocephalus (Rowland, 2005). HBO has not been established as a treatment option for cerebral edema.

Acute Coronary Syndrome (ACS)/Myocardial Ischemia/Infarction (MI), Cardiogenic Shock/Preconditioning for Coronary Artery Bypass Graft Surgery): ACS includes acute MI and unstable angina. HBO therapy has been proposed as an adjunct to standard therapy to improve oxygen supply to the heart and possibly decrease the amount of myocardial ischemic death that could occur and/or to prevent cardiogenic shock. HBO has also been investigated for preconditioning coronary artery disease (CAD) patients prior to elective surgery to improve left ventricular stroke work postoperatively. However, there is insufficient evidence to support the effectiveness of HBO for these conditions.

Bennett et al. (2011) (updated 2015) conducted a systematic review of randomized controlled trials comparing the treatment of acute coronary syndrome (ACS) with HBO and without HBO. Six trials (n=665) met inclusion criteria. Overall, HBO resulted in a significant decrease in the risk of death ($p=0.02$), a significantly lower extent of heart muscle damage measured by lesser rise in muscle enzymes ($p=0.005$) and a significantly better left ventricular ejection fraction ($p=0.001$). Evidence from individual trials reported a reduction in the risk of major adverse coronary events (MACE) ($p=0.003$), re-infarction ($p=0.04$), dysrhythmias ($p=0.01$) and less time to relief of pain ($P<0.00001$). However, the authors warned that because of the "modest number of patients, methodological shortcomings and poor reporting, these results should be interpreted cautiously, and an appropriately powered trial of high methodological rigor is justified to define those patients (if any) who can be expected to derive the most benefit from HBOT. The routine application of HBOT to these patients cannot be justified from this review." No new trials were located in the most recent update (Bennett, et al., 2015).

Yogarathnam et al. (2010) conducted a randomized controlled trial (n=81) to determine if preconditioning coronary artery disease (CAD) patients with HBO prior to first-time, elective coronary artery bypass graft surgery (CABG) with on-pump cardiopulmonary bypass (CPB), would improve postoperative myocardial left ventricular stroke work (LVSW). Preoperatively, the study group (n=41) received HBO for two 30-minute intervals, five minutes apart. The control group (n=40) was not treated with HBO. Hemodynamic monitoring was performed on 22 HBO patients and 25 control group patients. Immediately following HBO, the study group had a significant reduction in pulmonary vascular resistance (PVD) ($p=0.03$), but the significant difference was not maintained. Intraoperatively, the HBO group had a significant reduction in blood loss ($p=0.05$). There was no significant difference in the rise in the serum troponin T level, but the rise was greater in the control group. This indicated that HBO-treated patients had less postoperative myocardial injury than the control group. Postoperatively, the HBO group had a significantly improved stroke volume ($p=0.01$) and LVSW ($p=0.05$), spent 24 minutes longer on mechanical ventilation and was intubated 36 minutes longer than the control group. The HBO group had a significantly shorter length of stay in the intensive care unit ($p=0.05$). The study group also had a reduction in blood loss (11.6%), blood transfusion (34%), low cardiac output syndrome (10.4%), inotrope use (8%), atrial fibrillation (11%), pulmonary complications (12.7%), and wound infections (7.6%), but the differences were not statistically significant. No renal or neurological complications were reported in the HBO group compared to 5% and 2.5%, respectively in the

control group. Author-noted limitations of the study included the small patient population, recruitment of low-risk patients, and lack of comparison to patients who underwent CAPG without the use of CPB and to patients with controlled ischemia. Another limitation of the study is that all patients were not hemodynamically monitored during the postoperative period.

In a randomized controlled trial by Dekleva et al. (2004), 74 patients were assigned to HBO and streptokinase treatment versus streptokinase treatment alone within the first 24 hours after diagnosis. This study was small in sample size, showed treatment effectiveness limited to the first three days following HBO, and excluded patients with significant electrical complications. Due to these limitations, the effectiveness of HBO for the treatment of acute MI cannot be determined.

Acute or Chronic Cerebral Vascular Insufficiency: Cerebral vascular insufficiency is defined as insufficient blood flow to the brain that can lead to a stroke or transient ischemic attack (TIA). Although HBO has been proposed as a treatment option for cerebral vascular insufficiency, there is insufficient evidence in the peer-reviewed scientific literature to support its use for this indication.

Acute Thermal and Chemical Pulmonary Damage: HBO for the treatment of acute thermal and chemical pulmonary damage including smoke inhalation and pulmonary insufficiency in the absence of acute carbon monoxide poisoning is not supported by the evidence in the peer-reviewed literature.

Acute Wound, Flap and/or Graft: HBO has been proposed for the treatment of acute wounds, flaps and grafts. Published studies have included randomized controlled trials, case series and retrospective reviews. Dauwe et al. (2014) conducted a systematic review of the literature to evaluate the role of HBO in the treatment of acute wounds, flaps and grafts. Four randomized controlled trials, three prospective studies and one retrospective review met inclusion criteria. The studies included treatment of burn patients, crush injuries, postoperative ecchymosis following face lift surgery, post mastectomy and free parascapular flaps for lower extremity reconstruction. Due to the heterogeneity of the small patient populations (n=5-125), the poor methodology of the studies and conflicting outcomes, the authors concluded that the data did not support HBO for these indications.

Anorectal Disorders: HBO has been proposed as a treatment option for anorectal disorders (e.g., chronic anal fissure, internal hemorrhoids, infectious proctitis). The efficacy of HBO as primary or adjunctive treatment for anorectal disorders has not been established. Randomized controlled trials comparing HBO to standard care (e.g., non-steroidal anti-inflammatory medications, steroid enemas, cauterization or surgical excision) are lacking (Rao, 2004; Schwartz, 2004).

Autism: Autism is the most common condition in the group of developmental disorders known as autism spectrum disorders (ASD). HBO has been proposed as a potential treatment modality for improving cognitive function by increasing tissue oxygenation and improving cerebral blood flow. There are a limited number of randomized controlled trials evaluating HBO for the treatment of autism. Published studies have been primarily in the form of case series with small, heterogeneous patient populations (n=6-18) and involved various HBO treatment regimens (Sakulchit, et al., 2017; Rossignol, et al, 2007; Rossignol and Rossignol, 2006).

Xiong et al. (2016) conducted a Cochrane review of randomized and quasi-randomized controlled trials to investigate hyperbaric oxygen therapy for the treatment of autism spectrum disorder (ASD). One trial with a total of 60 children met inclusion criteria. Subjects were randomized to hyperbaric oxygen therapy or sham treatment. The quality of evidence was rated as low due to the small sample size and wide confidence intervals. Other limitations of the evidence included selection bias, short duration of treatment and short-term follow-up. Overall, there was no

reported improvement in social interaction and communication, behavioral problems, communication and linguistic abilities, or cognitive function. Regarding safety of hyperbaric oxygen therapy, minor-grade ear barotrauma events were reported. The authors concluded that, there is no evidence that hyperbaric oxygen therapy improves core symptoms and associated symptoms of ASD.

In 2013 (updated 2021), the FDA posted a warning to consumers regarding HBO. The Agency stated that HBO has not been clinically proven to cure or be effective in the treatment of autism. The information warns against the use of HBO for indications that are not FDA approved.

Ghanizadeh (2012) conducted a systematic review of randomized controlled trials to evaluate the efficacy of HBO for the treatment of autism in children. Two randomized controlled trials met inclusion criteria. One study was the Rossignol et al. study discussed below. The second study (n=42) reported that HBO was not more effective than placebo.

Rossignol et al. (2009) conducted a multicenter, randomized, double-blind, controlled trial to evaluate the efficacy of HBO in the treatment of children (n=62), ages 2–7 years, diagnosed with autistic disorder. The children were randomly assigned to the study group (n=33) treated with HBO at 1.3 atmosphere and 24% oxygen or to the control group (n=29) treated with slightly pressurized room air and 21% oxygen. Forty, one-hour sessions (two sessions per day for five days) were administered over four consecutive weeks. Compared to the control group, the treatment group had significantly improved outcomes in the mean physician Clinical Global Impression (CGI) scale in overall functioning (p=0.0008), receptive language (p<0.0001), social interaction (p=0.0473), and eye contact (p=0.0102). Significantly more children in the treatment group were rated as “very much improved” (p=0.0471) or “much improved” (p=0.0024). Significant improvements were also reported by the treatment group in the parental CGI scores in overall functioning (p=0.0336), receptive language (p=0.0168), and eye contact (p=0.0322). Significant improvements were noted in total score, irritability, stereotypy, hyperactivity and speech (p<0.03 for each) on the Aberrant Behavior Checklist in the treatment group. The treatment group also showed significant improvement in the Autism Treatment Evaluation Checklist sensory/cognitive awareness score (p=0.0367) compared to the control group. Children over age five years with lower initial autism severity showed the most significant improvements. Due to the short-term duration of this study, the authors stated that studies with long-term outcomes were needed to formally validate the results. It is also unknown what the ideal HBO treatment regimen is for this patient population.

Following a review of the evidence, which included one randomized controlled trial and three case series, Undersea and Hyperbaric Medical Society (UHMS) (2009) concluded that although there is a strong case for further studies on the role of HBO in the treatment of autism, HBO cannot be recommended as a routine treatment option.

Brain Injury, Closed Head Injury, Traumatic Brain Injury (TBI), Anoxic Encephalopathy:

In patients with moderate or severe TBI, the goal is to resuscitate the patient adequately to prevent further brain injury. The available evidence on adjunctive HBO treatment for severe traumatic brain injury is limited, and patient outcomes following HBO therapy are uncertain (Rowland, 2005).

Crawford et al. (2017) conducted a systematic review to evaluate the efficacy of HBO for the treatment of traumatic brain injury (TBI). Twelve randomized controlled trials met inclusion criteria. Four studies (n=250) included patients with mild TBI suffering persistent symptoms over many months. Seven studies included patients with moderate-to-severe TBI treated acutely and one study did not clearly define the severity of the TBI. Overall, there were no statistically significant differences between HBO and sham. There were some statistical significant within

group differences reporting improvement with HBO regarding cognitive performance and post-concussion symptom severity. Minor adverse events included ear pain, nausea, sinus pain, headaches, tooth pain, transient worsening of myopia and musculoskeletal pain. No serious adverse events were reported. Eight studies were rated as acceptable methodology and four as low quality methodology. Limitations of the studies included the heterogeneity of the treatment regimens and outcomes measures. Studies used various exposure times to HBO (60–117 minutes), number of sessions (3–40), length of sessions (8–10 weeks), and amount of pressure used (1.5–2.4 ATA). The types of sham arms differed in terms of pressure and oxygen levels used. Additional research is needed to support the efficacy of HBO for the treatment of TBI.

Hawkins et al. (2017) conducted a systematic review of the literature to assess the effectiveness of HBO for the treatment of concussion in subjects who suffered from mild traumatic brain injury or post-concussion syndrome. Five randomized controlled trials met inclusion criteria. Studies included 50–61 subjects and compared HBO to sham and in two studies to sham, HBO and another oxygen fraction treatment group. Four studies reported no significant improvement with HBO.

Wang et al. (2016) conducted a systematic review and meta-analysis to investigate HBO for the treatment of traumatic brain injury (TBI). Eight studies (n=519) comparing hyperbaric oxygen therapy vs. control in patients with mild (Glasgow coma scale [GCS] 13–15) to severe (GCS 3–8) TBI were included. Studies were either randomized controlled trials or prospective two-arm studies. The primary outcome was the GCS. Secondary outcomes included the Glasgow outcome score (GOS), overall mortality, and changes in post-traumatic stress disorder (PTSD) score. Mean age of the subjects ranged from 23–41 years. Meta-analysis of two studies (n=120) revealed that the change in GCS score was significantly higher in the HBO group ($p<0.001$). Analysis of three studies (n=141) showed significantly higher rate of improvement in GOS ($p=0.020$) and a lower overall mortality rate ($p<0.001$). There was no significant change in the PTSD score between the control group and HBO group. The pooled odds ratio for the GOS improvement rate became insignificant with the removal of two studies indicating poor reliability of the meta-analysis. Limitations of the studies included: heterogeneity of pooled data; all the studies had incomplete outcome data; number of studies included in the final analysis were few (2–3); heterogeneity of treatment regimens (e.g., starting time of HBO, oxygen concentration, pressure of treatment protocol); and poor reliability of the GOS meta-analysis. A subgroup analysis of mild and severe TBI was not performed due to incomplete reporting of data and the limited number of eligible studies. The authors noted that whether HBO has a significantly favorable outcome in mild TBI patients as opposed to severe TBI patients is unknown.

Randomized controlled trials have reported that there was no significant improvement when HBO was used for the treatment of traumatic brain injury and post-concussion syndrome. Miller et al. (2015) conducted a multi-center, randomized controlled trial comparing standard care alone to standard care plus HBO or standard care plus sham for the treatment of mild traumatic brain injuries. Treatment regimens included 40 HBO sessions administered at 1.5 atmospheres absolute (ATA) or 40 sham sessions consisting of room air at 1.2 ATA. The primary outcome measure was the Rivermead Post-Concussion Symptoms Questionnaire (RPQ). The sham group and the HBO group showed improvement in the RPQ scores ($p=0.002$ and $p=0.008$, respectively) but there was no significant difference in clinical outcomes in the sham group vs. the HBO group ($p=0.70$). Cifu et al. (2014) (n=61) compared HBO to sham for the treatment of mild traumatic brain injury and post-concussion syndrome. Treatments included 40, once daily, 60-minute hyperbaric chamber compressions at 2.0 atmospheres absolute (ATA) at 1 of 3 randomly preassigned oxygen fractions which resulted in blinded groups. The primary outcome measure was the Rivermead Post-Concussion Questionnaire-16 (RPQ-16) taken before compressions and at one week and three months following HBO. At the three-month follow-up no significant improvements in symptoms, functional status, or cognitive or psychomotor performance were seen with HBO.

In a Cochrane review of randomized controlled trials, Bennett et al. (2012) evaluated the benefits and harms of adjunctive HBO for the treatment of patients with TBI. The authors concluded that the combined results of the studies, involving 571 patients, suggested that HBO may reduce the risk of death and improve the final Glasgow Coma Scale. However, there was little evidence of a good outcome and the routine use of HBO for this subpopulation was not supported by the evidence.

Brown Recluse Spider Bites: Brown recluse spider (i.e., *Loxosceles reclusa*) venom contains enzymes that cause local (e.g., dermonecrosis) and systemic toxicity. There are a limited number of case studies that administered HBO as a treatment option. The studies did not show that HBO therapy produced better patient outcomes than standard aggressive wound care and antibiotic administration (Ruha, 2024; Arnold, 2018, updated 2021; Wasserman, 2005).

Cancer: HBO therapy has been proposed for use as a cure for cancer and as a means of enhancing tumor response to chemotherapeutic treatment. The American Cancer Society, the National Cancer Institute and the National Comprehensive Cancer Network® do not discuss HBO as a treatment option for any cancers. In 2013 (updated 2021), the FDA posted a warning to consumers regarding HBO. The Agency stated that HBO has not been clinically proven to cure or be effective in the treatment of cancer.

Bennett et al. (2018) conducted a Cochrane review of randomized controlled trials to assess the safety and efficacy of administering radiotherapy for the treatment of malignant tumors while breathing HBO. Nineteen trials included 2286 subjects of which 1103 were allocated to HBOT and 1153 to control groups. For head and neck cancer, there was an overall reduction in the risk of dying at one year and five years following therapy. There was also some evidence of improved local tumor control immediately following irradiation, at year one and year five. The evidence was considered of moderate quality due to the inconsistency of outcomes between trials. No trials reported quality of life outcomes. It was noted that benefits came at the cost of an increased risk of severe local radiation reactions with HBOT (high level of evidence). There was no clear benefit of HBOT for the treatment of cervical cancer or bladder cancer. When all cancer types were combined, there was high-quality evidence for an increased risk of severe radiation tissue injury during the course of radiotherapy with HBOT and moderate quality of evidence of oxygen toxic seizures during treatment. The authors noted that given the methodological and reporting inadequacies of the studies, the results should be reviewed with caution. Additional research is needed to determine the benefits of HBO for head and neck cancer. Based on the evidence HBO is not supported for the treatment of uterine cervical or bladder cancer. There is little evidence available concerning malignancies at other anatomical sites.

Carbon Tetrachloride Poisoning: Poisoning from carbon tetrachloride, which is used in industrial solvents, grain fumigants, insecticides, and the production of fluorocarbons, may cause nausea, vomiting, abdominal pain, diarrhea, confusion, coma, respiratory depression, hypotension, convulsions and even death. Although HBO has been proposed as a treatment option for carbon tetrachloride poisoning, there is insufficient evidence to support its effectiveness.

Cerebral Palsy: Cerebral palsy (CP) is an umbrella term covering a group of nonprogressive, but often changing, motor-impairment syndromes secondary to lesions or anomalies of the brain arising in the early stages of development. The evidence in the peer-reviewed literature does not support HBO for the treatment of CP.

Laureau et al. (2022) conducted a systematic review (n=1008) of five randomized control trials (RCT), six observational studies and one retrospective review to evaluate the effectiveness and safety of HBO in the treatment of cerebral palsy (CP). In the RCTs, the treatment intervention was

100% O₂, 1.5 to 1.75 atmospheres absolute (ATA). The comparator was pressurized air in three RCTs and physical therapy in two RCTs. Similar improvements were observed regarding motor and/or cognitive functions in the HBOT and control groups. Most common adverse event was middle ear barotrauma (up to 50% of children). Other adverse events included seizures, confinement anxiety, pulmonary disorders, nausea, hypoglycemia, hypotension, visual disorders and dizziness. Author noted study limitations include variable control intervention, heterogeneity of clinical presentation of CP, and variable length of time period of treatments. In conclusion, HBOT does not improve motor function, cognition, and functional performance in children with CP.

Lacey, et al. (2012) conducted a randomized controlled trial (n=46) to determine if HBO would improve functional abilities in children (ages 3–8 years) with spastic CP. One group received HBO (n=24), 100% oxygen at a pressure (or depth) of 1.5atm; and the second group received hyperbaric air (HBA) (n=22), a mixture of gases (14% oxygen) at 1.5atm to simulate 21% oxygen at room air. Eighty-minute sessions took place once a day for eight weeks for a total of 40 treatments. At the six month follow-up there were no changes from baseline in the Gross Motor Function Measure (GMFM)-88 and GMFM-66 or dimension A-D scores (i.e., lying and rolling, sitting, crawling and kneeling, standing) in either group. There were no significant differences between groups. The HBO group showed a significant increase in dimension E score (walking, running and jumping). Although both groups showed improvement, there was also no significant difference between the groups in the Pediatric Evaluation of Disability Inventory (PEDI) scores. The study was stopped because “the calculated conditional probability of obtaining a difference between groups if the study continued to the end was only between 0.5% and 1.6%”. The results of the study do not support HBO in the treatment of this patient population.

In a 2007 systematic review including two randomized controlled trials and four observational studies evaluating the benefits and adverse effects of HBO for the treatment of CP, McDonagh et al., reported that the improvements in motor function when compared to baseline for both HBO and room air were not significantly different. The evidence to support HBO therapy for CP is insufficient at this time.

In a clinical report for the American Academy of Pediatrics (AAP) regarding the treatment of children and youth with CP, Liptak et al. (2011) listed HBO as a therapy for which some evidence exists to refute its effectiveness. “Because CP is so heterogeneous, it is unlikely that all children would improve with a single therapy; benefits have not been proven”.

Cerebral Radionecrosis: Cerebral radionecrosis is a complication of radiation therapy of intracranial and extracranial tumors. Delayed radionecrosis may appear as an intracranial mass and is typically surgically removed. Although HBO has been suggested as a treatment option when surgery is not feasible, clinical trials demonstrating the efficacy of HBO for this indication are lacking.

Chronic Fatigue Syndrome: Chronic fatigue syndrome (CFS) is a disorder of unknown etiology, which may have an infectious basis. It involves a state of chronic fatigue for more than six months and can be accompanied by cognitive difficulties. Because most cases of CFS may be based on a viral infection, no effective therapy exists (Roberts, 2024). Evidence supporting HBO for the treatment of CFS is lacking.

Chronic Peripheral Vascular Insufficiency: Peripheral vascular insufficiency is most commonly a disease of the arteries and is caused by atherosclerosis which results in insufficient tissue perfusion. Although HBO has been proposed as a treatment option for peripheral vascular insufficiency, there is insufficient evidence in the peer-reviewed literature to support HBO for this indication.

COVID-19

COVID-19 is the infectious disease caused by the coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). There is insufficient evidence to support HBO for the treatment of this condition. In 2021, the FDA posted a warning to consumers regarding HBO. The Agency stated that HBO has not been clinically proven to cure or be effective in the treatment of COVID-19.

The Undersea and Hyperbaric Medical Society (UHMS) Position Statement: Hyperbaric Oxygen (HBO2) for COVID-19 Patients (jointly approved by the American College of Hyperbaric Medicine [ACHM]).(revised August 2020) states :

- "The UHMS continues to advocate strongly for well-designed IRB-approved clinical trials of hyperbaric oxygen for COVID-19. Well-designed trials are necessary to establish a proper mechanistic and clinical foundation for COVID-19 treatment and increase our understanding of this disease and the potential role of hyperbaric oxygen as part of a multidisciplinary approach.
- The UHMS recognizes the special value of Phase III randomized controlled trials in providing Level I evidence and strongly supports funding and conduct of these definitive studies.
- The UHMS now recognizes that hyperbaric oxygen treatment on an off-protocol basis for COVID-19 at the physician's discretion may be appropriate in some cases and recognizes that community centers and free-standing facilities may have limited access to IRBs (Institutional Review Boards). The UHMS strongly encourages well-documented scientific observations of the impact, patient selection criteria, and treatment methodology for those utilizing HBO2 to treat COVID-19 patients off-protocol.
- The UHMS strongly encourages well-documented scientific observations of the impact, patient selection criteria, and treatment methodology for those utilizing HBO2 to treat COVID-19 patients. The UHMS Research Committee has published key outcome determinants and therapeutic guidance."

Literature Review: There is insufficient evidence in the published peer-reviewed scientific literature to support the effectiveness of HBO for the treatment of acute or long COVID-19. The available studies are primarily focused on the effects of HBO on hospitalized patients with pneumonia and/or on mechanical ventilation. Studies are in the form of retrospective reviews, case series, and case reports (Gorenstein, et al., 2020; Guo, et al., 2020; Thibodeaux, et al., 2020). Studies on long Covid consist of a randomized control trial by Zilberman-Itskovich et. al. (2022) and a retrospective review (n=10) by Robbins et al. (2021).

Zilberman-Itskovich et. al. (2022) conducted a randomized, sham control, double blind clinical trial (n=73) to evaluate the effects of HBO on patients with persistent symptoms of post-Covid for at least three months after confirmed infection. Patients were randomized to receive HBO daily for 40 sessions (n=37) or sham treatment (n=36). Follow-up assessments were performed at baseline and 1–3 weeks after the last treatment session. The primary outcome was the cognitive assessment as evaluated by the Mindstreams computerized cognitive testing battery. The cognitive domains assessed were memory, executive function, attention, information processing speed, and motor skills. Cognitive scores were normalized for age, gender and educational levels. Secondary outcomes were measured using self-reported questionnaires: short form-36 (SF-36) to assess quality of life, the Pittsburgh Sleep Quality Index (PSQI) to assess sleep quality, the Brief Symptom Inventory (BSI-18) to evaluate psychological distress, based on three subscales: depression, anxiety, and somatization, and the Brief Pain Inventory (BPI) to measure pain intensity and impact. Brain imaging MRI scans were performed to evaluate brain perfusion. Following HBOT, there was a significant group-by-time interaction (difference in the change over time) with a medium net effect size in global cognitive function (d=0.495, p=0.038), attention

($d=0.463$, $p=0.05$) and executive function ($d=0.477$, $p=0.04$ and) compared to the control group. Significant improvement was also demonstrated with the following net effect size in the energy domain ($d=0.522$, $p=0.029$), sleep ($d=0.48$, $p=0.042$), psychiatric symptoms ($d=0.636$, $p=0.008$), and pain interference ($d=0.737$, $p=0.001$). Increased brain MRI perfusion and microstructural changes were noted in the supramarginal gyrus, left supplementary motor area, right insula, left frontal precentral gyrus, right middle frontal gyrus, and superior corona radiate of the brain following HBO treatment. Adverse events were not reported. Study limitations include small patient population, short follow-up time, and the optimal number of sessions for maximal therapeutic effect is unknown. Hadanny et. al. (2024) reported on the results of 31 patients in the treatment arm 12 months post last treatment with HBO. The authors reported that the short-term improvements in quality of life, quality of sleep, psychiatric and pain symptoms persisted at 12 months. Additional well-designed RCTs with large patient populations and long-term follow up are needed to determine if HBO is effective in the treatment of long-Covid.

Oliaei et al. (2021) conducted a systematic review to evaluate the effects of hyperbaric oxygen therapy (HBO) on the treatment of COVID-19 pneumonia. The analysis included eight studies of which three were clinical trials and five were case reports and case series. Although the overall results reported improvement following HBOT, the limitations of the studies prevent strong conclusions to be drawn supporting HBOT for the treatment of Covid.

Crohn's Disease: Crohn's disease is a chronic inflammatory disease of the gastrointestinal tract, the cause of which remains unknown. The available evidence is limited and is considered insufficient to determine the effect of HBO treatment on the health outcomes of patients with Crohn's disease.

Dulai et al. (2014) conducted a systematic review to evaluate the safety and efficacy of HBO for Crohn's disease ($n=286$) and ulcerative colitis ($n=327$). A total of 17 studies met inclusion criteria. One study was a randomized controlled trial. The remaining studies were case reports and case series. Known grades ($n=40$) of ulcerative colitis ranged from mild to severe. Of the 44 Crohn's disease patients who had disease extent reported, 40 (91%) had perineal disease and 21 (48%) had fistulas. The overall response rate for patients with irritable bowel disease (IBD) was 86%. The response rate for ulcerative colitis patients who had endoscopic follow-up ($n=40$) was 100%. Six patients suffered serious adverse events necessitating discontinuation of therapy (6.7/10,000 treatments). Limitations of the studies noted by the authors included: high or uncertain risk of bias; small, heterogeneous patient populations; lack of endoscopic follow-up of disease activity; missing data; unclear study designs; poorly described research methods; heterogeneity of treatment regimens; short-term follow up; and inadequate description of outcomes.

Cutaneous, Decubitus/Pressure Ulcers: Cutaneous, decubitus (pressure) ulcers are typically localized to an area of tissue necrosis that develops when soft tissue is compressed between a bony prominence and an external surface. HBO for the treatment of decubitus or pressure ulcers has generally been considered ineffective or not extensively evaluated (Javier, 2024).

Dementia: Dementia is characterized by progressive deterioration that interferes with social or occupational functions, such as: memory, orientation, abstraction, ability to learn, visuospatial perception, language function, and constructional praxis. Alzheimer's disease accounts for more than 50% of cases of dementia (Rowland, 2005). There is insufficient evidence in the peer-reviewed literature to support the treatment of dementia with HBO.

Xiao et al. (2012) conducted a Cochrane systematic review to assess HBO for the treatment of vascular dementia. One randomized controlled trial ($n=64$), "of poor methodological quality" met inclusion criteria. There is insufficient evidence to support HBO for the treatment of this condition.

Epilepsy: Epilepsy, or seizure disorder, is characterized by the tendency to have recurring seizures. HBO is proposed for the treatment of this condition as a means to improve cerebral circulation to the brain and decrease cerebral edema. HBO for the treatment of epilepsy has not been established.

Fractures (e.g., Acute, Delayed Union and/or Nonunion): The primary goal in the treatment of fractures is the realignment and stabilization of the fractured bone and restoration of function. HBO has been proposed to assist in improving the healing outcomes in delayed or nonunion fractures, but improvement in clinical outcomes has not been established.

In a Cochrane systematic review, Bennett et al. (2012) concluded that, although HBO has been proposed for many years for the treatment of fractures, there is insufficient evidence within the literature to support or refute that it aids in the healing of acute injuries and fractures, and/or assists in the healing process of a nonunion fracture. No studies met inclusion criteria.

Headaches (Cluster and Migraine): Cluster headaches and migraine headaches are distinct primary headaches. Both are extremely painful with cluster headaches being less common than migraine headaches. According to the International Headache Society, a cluster headache is on one side of the head and lasts 30–90 minutes. A migraine headache is a chronic condition with recurrent, episodic attacks that last hours to days. Although HBO has been proposed as a treatment option for headaches, there is insufficient evidence in the peer-reviewed literature supporting the efficacy of HBO for the treatment of these conditions.

Bennett et al. (2008) (updated 2015) conducted a systematic review and meta-analysis to evaluate the safety and efficacy of HBO compared to normobaric oxygen therapy (NBOT) used for the prevention and treatment of migraine and cluster headaches. The review included 11 randomized controlled trials (n=209) including five trials (n=103) that compared HBO to sham for acute migraines, two (n=29) that compared HBO to sham therapy for cluster headaches and one evaluated NBOT (n=56) to sham for a mixed headache group. Pooled data from three trials (n=58) suggested that HBO was effective in relieving migraines compared to sham therapy, but provided no evidence that HBO could prevent migraines or reduce nausea, vomiting or medication requirements. There was no evidence that HBO was effective for the termination of cluster headache.

Hepatic Necrosis: Hepatic necrosis is a severe and progressive form of hepatitis associated with hepatocellular death and hepatic failure. Although HBO has been proposed as a treatment option for hepatic necrosis, there is insufficient evidence in the peer-reviewed literature to support its use for this condition.

Human Immunodeficiency Virus (HIV) – Fatigue: Fatigue is often a chronic, debilitating symptom of individuals infected with HIV. It has been hypothesized that increased oxygenation by HBO may help to relieve the fatigue. However, evidence in the peer-reviewed literature supporting this hypothesis is lacking.

In Vitro Fertilization (IVF): Infertility may be the result of endometriosis, or abnormalities in tubal, uterine, endometrial, cervical, or ovulatory functions. It has been proposed that increasing oxygenation by HBO may aid in egg maturation and alignment of chromosomes during meiosis but there is insufficient evidence to report this claim.

Lyme Disease: Lyme disease is a clinical diagnosis, and currently the early use of antibiotics can prevent persistent, recurrent, and refractory conditions. The duration of therapy is determined by each individual's clinical response, but the adjuvant use of HBO therapy is not recommended as part of this treatment.

Lymphedema: Approximately 10–38% of all women who have breast-conserving surgery (BCS) or modified radical mastectomy have postsurgical irradiation to the lymph nodes, and 10% of those women develop lymphedema. HBO has not been established as an effective adjunctive treatment for the reduction of lymphedema. Studies have primarily been in the form of case series with small patient populations (n=10) and reported that the total limb volume did not change significantly from baseline measurements (Teas, et al., 2004).

Gothard et al. (2010) conducted a randomized controlled trial (n=58) to investigate the effectiveness of HBO in the treatment of patients with ipsilateral arm lymphedema, $\geq 15\%$ increase in arm volume, following treatment for cancer. Diagnosis included breast cancer (n=56) and Hodgkin lymphoma (n=2). All patients had undergone surgery and radiation therapy. The average interval of time from radiation therapy to randomization was 2.1–21.5 years. Patients were randomized to HBO (n=38) or to the control group (n=20). The study group received 30 HBO treatments while the control group continued best standard care for lymphedema according to the 2006 Lymphoedema Framework Best Practice for the Management of Lymphoedema International Consensus. At the 12-month follow-up (n=46), there were no statistically significant differences from baseline to follow-up in the median volume of the ipsilateral limb (expressed as a percentage of contralateral limb volume) and change over time in either group. There was no clear within-patient improvement from baseline to 12 months with either group. Author-noted limitations of the study included the small patient population and the interval of time from radiation therapy to randomization.

Malignant Otitis Externa: Malignant otitis externa (i.e., necrotizing external otitis) is an uncommon, yet potentially fatal infection of the external auditory canal and may involve surrounding tissue and soft bone. HBO therapy has been proposed as an adjunct to traditional therapy (e.g., diabetic control, administration of antibiotics, repeat debridement and surgical resection). However, the efficacy of HBO for this condition has not been established.

Phillips et al. (2013) conducted a Cochrane systematic review to determine the effectiveness of HBO when used as an adjunct to the traditional treatment protocols for malignant otitis externa. The researchers could not locate any randomized controlled trials that had measured the effectiveness of HBO within this population. A small number of case reports and case series were found, but there was no clear evidence that demonstrated the effectiveness HBO therapy for this condition.

Multiple Sclerosis: Multiple sclerosis (MS) is a chronic neurological disease in which there is patchy inflammation, demyelination, and gliosis in the central nervous system. HBO has been proposed as a treatment modality for MS based on the demonstrated ability of HBO to produce vasoconstriction with increased oxygen delivery and some anecdotal evidence of efficacy.

In a Cochrane systematic review, Bennett and Heard (2011) investigated the use of HBO for the treatment of MS. Two randomized controlled trials reported generally positive results, but the remaining seven randomized trials reported no evidence of treatment effects. Due to the paucity of evidence to confirm beneficial effects of HBO, the authors did not believe that routine use of HBO was justified.

Mycoses: Mycosis is an infection, or a disease caused by a fungus (e.g., candidiasis, aspergillosis, cryptococcus). Zygomycosis (e.g., mucormycosis, phycomycosis) is an infection caused by “bread mold fungi” and can infect immunosuppressed individuals (e.g., HIV). HBO has been proposed as a treatment option for some forms of invasive mycosis (e.g., zygomycosis), but its efficacy remains unproven (McAdam and Sharpe, 2005).

Nonvascular Causes of Chronic Brain Syndrome (e.g., Pick's Disease, Alzheimer's Disease, Korsakoff's Disease): Chronic Brain Syndrome, also called dementia, is a loss of brain function. Alzheimer's disease and Pick's disease are forms of dementia. Alzheimer's is a primary degenerative dementia that typically involves diffuse atrophy of the brain, while Pick's disease is a classical frontotemporal dementia. Korsakoff's is a psychosis that results from a thiamine deficiency and is primarily a memory disorder. The efficacy of HBO for these conditions has not been established (Smith and Seirafi, 2006).

Ophthalmologic Conditions Other Than Central Retinal Artery Occlusion (e.g., Optic Neuropathy, Glaucoma): HBO has been proposed as an adjunctive treatment option for various ophthalmologic conditions, including, optic neuropathy, and glaucoma. There is insufficient evidence to determine the health outcomes of HBO for the treatment of ophthalmologic conditions other than central retinal artery occlusion.

Organ Transplant/Storage: Researchers have hypothesized that HBO may enhance the performance and growth in pancreatic islet grafts when they are subjected to high levels of oxygen prior to transplant. HBO has also been proposed for administration following organ transplantation to reduce the risk of organ rejection (e.g., liver) as well as, keeping donated organs viable for a longer period of time. However, additional research is required to establish the efficacy of HBO therapy on organ transplantation and storage (Muralidharan, et al., 2007; Juang, 2002).

Penile glans necrosis: Penile glans necrosis (glans penis necrosis) is a rare clinical condition caused by trauma, poorly controlled diabetes mellitus, adverse effect of vasoconstrictive solutions, end-stage renal disease due to calciphylaxis, obesity, and circumcision. HBO has been proposed as an adjunctive treatment option of glans penis necrosis after prostatic artery embolization. There is insufficient evidence in the published peer-reviewed medical literature demonstrating the safety, efficacy, and long-term outcomes of hyperbaric oxygen therapy for the treatment of glans penis necrosis. Studies consist of one small case series (n=6) (Chung, 2023) and case reports.

Pulmonary Emphysema: Emphysema is defined as an abnormal permanent enlargement of air spaces in the distal bronchioles that is associated with chronic bronchitis. HBO has been proposed as a treatment option for emphysema, however, improvements in health outcomes have not been established in clinical trials.

Reflex Sympathetic Dystrophy (RSD)/Complex Regional Pain Syndrome (CRPS): CRPS, also known as RSD or causalgia, is a neuropathic condition that causes intense pain primarily in the arms, hands, legs, or feet. HBO has been proposed as a treatment option for the pain associated with CRPS. Evidence in the peer-reviewed literature does not support the effectiveness of HBO for the treatment of CRPS.

Kiralp et al. (2004) conducted a double-blinded, randomized, placebo-controlled study (n=71) to assess the effectiveness of HBO for treating patients with CRPS. The patients were allocated alternately to receive fifteen, 90-minute therapy sessions of HBO therapy (n=37) or normal air therapy (n=34). The visual analog scale score indicated that pain decreased starting from the first day until day 45 of treatment. An increase in wrist flexion was observed with the HBO group after 15 therapy sessions. A decrease in wrist circumference in the HBO group was also reported. There was a statistically significant difference for all variables except wrist extension. The study is limited by the small patient population and short-term follow-up. Additional studies with larger populations and long-term follow-ups are needed to validate the results of this clinical trial.

Rheumatoid Arthritis: Rheumatoid arthritis (RA) is a chronic systemic inflammatory disease of unknown cause that primarily affects the peripheral joints leading to joint destruction and limited

mobility. Although HBO has been proposed for the treatment of RA to decrease pain and inflammation, there is insufficient evidence supporting its efficacy.

Sepsis: Sepsis is a group of disorders that result from infection by bacteria, viruses, fungi, or parasites or the toxic products of these microorganisms. Sepsis involves early signs of circulatory compromise to full-blown circulatory collapse with potentially multi-organ system failure and death. The role of HBO as an adjunctive therapy in the treatment of sepsis remains controversial.

Sickle-Cell Disease: Sickle-cell disease is a hereditary disorder of hemoglobin structure and function. The anemia of sickle-cell disease is due to both chronic and acute hemolysis. Several new approaches to treatment of sickle-cell disease are currently under evaluation; however, these approaches do not include HBO (Lodewijk, 2007). Studies supporting HBO for the treatment of sickle-cell anemia are lacking.

Soft Tissue Injury (e.g., Delayed Onset Muscle Soreness, Closed Soft Tissue Injury, Sprains, Strains): Soft tissue injuries can range from abrasions and bruising to disruptions of tendons, ligaments, and muscles. Muscle soreness and damage are commonly associated with athletic activity. HBO has been proposed as an adjunct to conventional therapies (e.g., rest, elevation, pharmacotherapy) to expedite the healing process, but its beneficial impact on health outcomes has not been established.

According to Bennett et al. (2005) in a Cochrane systematic review including nine randomized controlled trials (n=219), there was insufficient evidence to conclude that HBO in the treatment of delayed onset of muscle soreness or closed soft tissue injury is efficacious.

Spinal Cord Injuries: Bruising, pressure, cutting or severance of the spinal cord may result in partial or complete loss of sensation and movement below the site of injury. Studies investigating the adjunctive use of HBO for the treatment of spinal cord injuries are primarily in the form of small, uncontrolled case series with a range of spinal cord injuries. Overall, results were not favorable. HBO therapy for the management of spinal cord injury has not been widely accepted (Rowland, 2005).

Stroke: Medical therapies for stroke are designed to minimize or prevent ischemic brain infarction, optimize functional recovery, and avert stroke recurrence. Specific therapies depend on the stroke syndrome. Xu et al. (2018) conducted a randomized controlled trial (n=79) to investigate the safety and efficacy of HBO therapy on diabetic patients who had suffered an acute intracerebral hemorrhage. Patients were randomized to 60 min of HBO daily for 30 days in a monophasic chamber or to normobaric oxygen therapy with similar protocol. No significant differences were seen between the groups at the one-month follow-up. At the six months follow-up significant improvements in the HBO group were seen in the modified Rankin Scale (p=0.045) and National Institutes of Health Stroke Scale (p=0.035), but not in the Barthel Index (p=0.080) and Glasgow Outcome Scale (p=0.73). Limitations of the study include the small patient population, short-term follow-up, and the patient population of diabetics with acute intracerebral hemorrhage preventing generalization of the results. Additional studies are needed to support HBO for this indication.

In a Cochrane review conducted by Bennett et al. (2014), the authors assessed the safety and effectiveness of adjunctive HBO therapy in the treatment of acute ischemic stroke. Eleven randomized controlled trials (n=705) met inclusion criteria. The authors concluded that there was no good evidence to show that HBO improved clinical outcomes when used for the treatment of this subpopulation. Additional research is needed to identify the benefit of HBO for these patients.

Guidelines for the early management of acute ischemic stroke (AIS) by the American Heart Association and the American Stroke Association (2018) stated HBO is not recommended for patients with AIS except when caused by air embolization. The limited available data show no benefit from HBO.

Tetanus: Tetanus is caused by the bacteria *Clostridium tetani* and is characterized by an acute onset of hypertonia and generalized muscle spasms. Although HBO has been proposed as a treatment option for tetanus, there is insufficient evidence in the peer-reviewed literature to support its efficacy.

Tinnitus: Tinnitus, also commonly referred to as “ringing in the ears” or “head noise,” is defined as the perception of sound in the head when no external sound is present. This symptom can occur in one ear or bilaterally, as well as internal and external to the auricle. HBO has been investigated as a treatment option in order to increase the supply of oxygen to the ear and brain in an attempt to decrease the severity of hearing loss and tinnitus. Overall, improved clinical outcomes have not been reported following HBO.

Bennett, et al. (2012) conducted a systemic review of seven randomized controlled trials (n=392) to assess the benefits and harms of HBO for the treatment of tinnitus and/or sudden sensorineural hearing loss. The significance of any improvement in tinnitus could not be assessed by pooled data and the routine use of HBO for the treatment of tinnitus could not be recommended.

In a study to analyze the effectiveness of HBO treatment on tinnitus, Porubsky et al. (2007) randomized 360 patients into two HBO treatment protocols (2.2 bar vs. 2.5 bar). Twelve patients (3.3%) experienced complete remission of tinnitus, in 122 (33.9) the intensity lessened, and 44 (12.2%) had a subjectively agreeable change of noise characteristics. No change was found in 157 cases (43.6%) and 25 (6.9%) experienced deterioration. There was no statistically significant difference between the two groups ($p > 0.05$). Out of 68 patients with a positive expectation of HBO effects, 60.3% stated that the tinnitus had improved compared to 47.2% of patients (n=271) who underwent therapy with an indifferent expectation and 19% (n=21) of patients with a negative expectation. The influence of subjective expectation on the outcome was statistically significant ($p < 0.05$).

Venous Stasis Ulcers: Venous stasis ulcers are the result of chronic venous insufficiency and can lead to life-threatening infections of the lower extremities. Although HBO therapy has been proposed for the treatment of this population, its efficacy has not been established by clinical trials. A Cochrane systematic review of randomized controlled trials evaluating HBO for the treatment of chronic wounds (Kranke, et al., 2012; updated 2015) included one trial with 16 patients who had venous ulcers. At six weeks the author reported significant reduction in the ulcer area. Large randomized controlled trials with long-term follow-ups are needed to validate the results of this study.

Keohane et al. (2023) conducted a systematic review to evaluate the efficacy of Hyperbaric Oxygen Therapy (HBO) in the complete healing or reduction in size of venous leg ulcers (VLU) when compared to control group. Six studies (n=166) met inclusion criteria. Two studies reported no statistically significant difference between HBOT and controls for the outcome of complete ulcer healing ($p = 0.4478$) at 12 weeks. Four studies reported non-significant results of complete ulcer healing at 5–6 weeks follow-up ($p = 0.1136$). A change in size of VLU was reported in each study ($p = 0.0024$). There was significant variance across the studies, with different controls, reporting of outcomes and duration of follow-up. Additional limitations included small, heterogeneous patient populations and short-term follow-up. Although studies reported a reduction in area of VLU versus controls, this does not translate to a significant clinical benefit.

Other Indications: Studies, primarily in the form of case series (n=5-20), case reports and retrospective reviews have investigated HBO as a primary or adjunctive therapy for various other indications including: altitude sickness, Bell's palsy, chronic non-healing wounds, comatose patients, cutaneous polyarteritis nodosa lesions, diabetes, frostbite, femoral head necrosis, fibromyalgia, gastrointestinal ulcers, heat stroke, high altitude illness, keloid recurrence, myofascial pain, Parkinson disease, chronic periodontitis, radiation-induced xerostomia, scleroderma, venomous snake bites, and to improve the success of osseointegration following maxillofacial implants. Overall, improved health outcomes following HBO for the treatment of these conditions have not been established.

The FDA issued a warning (2013, updated 2021) regarding the use of HBO for indications that are not FDA approved ("off-label"). Per the FDA, the safety and effectiveness of HBO has not been established for the following diseases and conditions: AIDS/HIV, Alzheimer's disease, autism, asthma, Bell's palsy, brain Injury, cerebral palsy, depression, diabetes, heart disease, hepatitis, migraine, multiple sclerosis, Parkinson's disease, spinal cord Injury, sport's injury, and stroke.

Fox et al. (2015) conducted a systematic review to assess the efficacy of HBO for the treatment of radiation-induced xerostomia and related quality-of-life (QOL). Studies included patients who had received radiation therapy for head and neck cancer and had not previously been treated with HBO. Seven studies met the inclusion criteria. Two studies were randomized controlled trials (n=45), four were prospective case series (n=121) and one was retrospective in design (n=80). The average number of HBO treatments ranged from 20–42.7 and the average time between radiation therapy and HBO treatment ranged from two days to 6.42 years. Overall, patients had increased stimulated saliva output, decreased sensation of dry mouth, and trends toward improvement in QOL related to dry mouth and sticky saliva. However, no significant improvement in overall QOL was demonstrated. The studies were limited by the small, heterogeneous patient populations; heterogeneity of treatment regimens; and lack of a comparator in the majority of the studies.

Wu et al. (2014) conducted a randomized controlled trial (n=80) to evaluate adjunctive HBO following autologous bone marrow mononuclear cells (BM-MNCs) infusion for the treatment of type 2 diabetes mellitus. Patients were treated with standard care for the first four months to reach optimal glycemic control. Thereafter, patients were randomized into four groups: BM-MNCs plus HBO; BM-MNC only, HBO only and standard care. The primary end point was C-peptide area under the curve (AUC) of the oral glucose tolerance test. Following 12 months of treatment, the AUC was significantly improved in the BM-MNC group and the MB-MNC plus HBO group compared to standard care (p<0.01, ea.) but there was no significant improvement with the use of HBO and BM-MNC compared to BM-MNC alone.

In a 2012 (updated 2016) Cochrane systematic review of the literature, Holland et al. reported that one low quality randomized controlled trial (n=79) suggested that HBO may be effective for the treatment of Bell's palsy. Further randomized controlled trials are indicated.

Esposito et al. (2013) conducted a systematic review of randomized controlled trials to investigate the effectiveness of HBO administered with dental implants. Only one randomized controlled trial with 26 patients met inclusion criteria. One year after implantation, four patients died from each group. There were no statistically significant differences for prosthesis and implant failures, postoperative complications, and patient satisfaction between the two groups.

Topical Oxygen Therapy (TOT)

Literature Review: Topical oxygen therapy (TOT) is a relatively new therapy that involves the administration of oxygen topically over tissue by continuous diffusion or pressurized systems using

mechanical devices. It has been explored as an adjunctive treatment method for diabetic foot ulcers (DFU's). Oxygen is generally applied at levels slightly higher than atmospheric pressure for ninety minutes per day, three to five days per week, or continuously, twenty-four hours per day, seven days per week. Although TOT has been proposed for the treatment of other conditions, including pressure ulcers, venous leg ulcers, and surgical wounds, there is presently insufficient evidence to support its use for these indications.

There is a growing body of evidence to demonstrate that TOT, when added to standard of care, improves DFU healing when compared to standard of care alone. Standard treatment measures for DFU include debridement, dressing, offloading, vascular assessment, infection management, and blood glucose control. A DFU is one of the most common and most severe complications of diabetes, with approximately 19%-34% of individuals with diabetes experiencing DFU in their lifetime. Oxygen is a critical element in the wound healing process. Wounds and tissue injuries cause the area to become hypoxic, likely due to disruption of the vasculature and increased oxygen consumption. TOT can mitigate oxygen deficiency by delivering oxygen directly to the wound bed, avoiding the need to rely on an impaired vascular or respiratory system. (OuYang, 2024; Chen et al., International Working Group on the Diabetic Foot, 2023 update; Sun et al., 2022)

OuYang et al. (2024) conducted a systematic review and meta-analysis to compare methods for treating patients with diabetic foot ulcers and assess their impact on healing. Treatment measures included platelet-rich plasma, negative pressure wound therapy, HBOT, TOT, ultrasonic debridement, acellular dermal matrix, and stem cell transplantation. The analysis included five studies (n=461), including the Frykberg study discussed below, comparing TOT with standard of care; one eight-week and four twelve-week treatment periods. The included studies were published between 2016 and 2020. Treatment with TOT significantly increased complete wound healing rates compared with SOC (Odds ratio 0.25, 95% confidence interval: 0.078 to 0.69). There was no significant effect on reduction in wound area, amputation rate, or adverse event rates.

Chen et al. (2024) conducted a systematic review of ten (n=792) randomized controlled trials of interventions to enhance healing of chronic foot ulcers in diabetes, including only trials that compared interventions to standard of care (SOC). The Frykberg trial, discussed below, was included in the systematic review. Interventions in addition to topical oxygen therapy (TOT) included debridement methods, dressings, negative pressure wound therapy, oxygen and other gases, physical therapies, skin substitutes, human tissue, autologous products, pharmacologic interventions, and metabolic management. Outcome measures included complete wound healing, time to wound healing, reduction in wound area, sustained healing, amputation, quality of life, activities of daily living, new infection, and mortality.

Six included studies (n=636) reported a positive effect on wound healing at twelve weeks with the use of TOT, although the size of the effect was uncertain. Three studies reported that TOT shortens the time to healing, nine studies (n=662) reported reduced wound size with the use of TOT. and three studies reported that TOT had no effect on amputation rates. The authors noted that only a handful of studies were assessed as being at low risk for bias, and although many new RCTs have been published in the last four years, most were assessed to be at either unclear or high risk of bias, with trial design problems, including nonblinding, analyses limited to per-protocol only, lack of a description of the randomization method or baseline characteristics, or any description of the usual care in the study protocol. The authors also stated that when usual care was described it often fell short of standards suggested by the International Working Group on the Diabetic Foot (IWGDF) guidelines, making the magnitude of additional interventions unclear. The authors concluded that the evidence to support any other intervention to enhance wound healing is lacking, and further high-quality randomized controlled trials are encouraged.

An updated systematic review and meta-analysis conducted by Sun et al. (2022) evaluated the efficacy and safety of TOT compared to standard of care for treatment of diabetic foot ulcers. The review included seven randomized controlled trials including 614 participants. TOT significantly increased complete wound healing rates compared to standard of care at eight and twelve weeks (relative risk 1.63, 95% confidence interval:1.33to 2.00, $P<0.00001$). It was unclear whether TOT could shorten ulcer healing time; two studies reported a significant reduction in time to wound healing, while two reported no statistical difference. Three of the studies reported significant wound area reduction compared with SOC, while the remaining two studies favored TOT, but intervention durations were inconsistent.

Frykberg et al. (2020) conducted a randomized, double blind, placebo-controlled trial to evaluate the efficacy of multimodality cyclical pressure Topical Wound Oxygen (TWO2) home care therapy (HyperBox; AOTI Ltd., Galway, Ireland) in healing refractory diabetic foot ulcers (DFUs) that had failed to heal with standard of care (SOC) alone. Seventy-three patients were included from 17 diabetic foot centers from the United States, United Kingdom, France, Germany, and Luxembourg. The average age was 63.3 years, 86% male with 68.5% White/Hispanic, 14% Black, 4.1% Asian, 1.4% American Indian, and 12.3% did not report race. Patients were included if they were 18–89 years old, had a diagnosis of Type 1 or 2 diabetes with a non-healing full-thickness, grade 1 or 2 DFU measuring ≥ 1 cm² and ≤ 20 cm² post-debridement. All ulcers included were present between four weeks and one year duration and received SOC treatment for at least four weeks. Patients were excluded if they had evidence of active severe infection, gangrene, osteomyelitis, active Charcot, uncontrolled diabetes (HgbA1c $>12\%$), known malignancy, on renal dialysis or serum creatinine >2.5 mg/dL. All patients received concurrent SOC therapy and were randomized to receive active TWO2 therapy (n=37) or sham treatment (n=36). Patients treated themselves at home for 90 min daily five times per week with either the allocated TWO2 or sham therapy until the ulcer healed or for a total of 12 weeks. No study therapy was done at the study centers. The primary outcome was the percentage of ulcers in each group achieving 100% healing at 12 weeks. Secondary end points included wound area reduction, 12-month incidence of both recurrence and complete healing, incidence of amputation, Cardiff Wound Impact Schedule (CWIS) QOL assessment, and adverse events. Following the first analysis point at 12 weeks, the independent data monitoring committee recommended that enrollment should be concluded, per the predetermined stopping rules, since the active arm was shown to be superior to the sham arm for the primary outcome. The TWO2 arm had a closure rate of 41.7% (15/37) compared with 13.5% (5/36) in sham arm ($p=0.010$). For open wounds at 12 weeks, the wound area reduction was 1.97 cm² in TWO2 arm and 0.40 cm² in sham arm. At the 12 month follow up, 56% of ulcers in the TWO2 arm were closed versus 27% of sham treated ulcers ($p=0.013$). Recurrence occurred in 6.7% (1/15) of TWO2 arm and 40% (2/5) of sham arm. Two index limb amputations (5%) occurred in the active arm compared with three index limb amputations (8%) in the sham arm. QOL improved for those whose ulcers healed. There were equal numbers of adverse events in the study arm and the sham arm. Serious adverse events included wound infection, osteomyelitis, hypoglycemic event, urinary tract infection, significant necrotic tissue, cardiovascular event, UTC grade 2 ulceration, severe maceration/dermatitis, and pneumonia. No device related events occurred. The author noted limitations included the small number of patients included in the primary end point analysis, noting also that the group was similar in size to those in other wound care randomized controlled trials. The authors concluded that this sham-controlled, double-blind, randomized controlled trial demonstrated that at both twelve weeks and twelve months, adjunctive cyclical pressured TWO2 therapy was superior in healing chronic DRSs compared with optimal SOC alone.

ECRI Clinical Evidence Assessment

A 2025 ECRI Clinical Evidence Assessment, Topical Oxygen Therapy for Diabetic Foot Ulcers, evaluated the current state of published literature, professional society guidelines, and position

and consensus statements. The analysis included a systematic review (SR) of randomized controlled trials (Chen, et al. 2024), a SR and network meta-analysis (OuYang et al., 2024), a SR of randomized controlled trials (Sun et al., 2023), all discussed above. The assessment also considered a single center randomized controlled trial (Pacek et al. 2023), and long-term follow-up to a randomized controlled trial (Al-Jalodi et al., 2022). Relevant published guidelines and position statements (e.g., American Diabetes Association, the Wound Healing Society, the International Working Group on the Diabetic Foot) were also considered.

The assessment concluded that TOT, when added to standard of care, improves DFU healing when compared to SOC alone, according to evidence from three SR's. One SR, however, concluded that TOT may not be as effective as platelet rich plasma, stem cells, and the combination of platelet rich plasma and negative pressure wound therapy, and that additional head-to-head comparisons are needed to enable firm conclusions.

After assessing the available published clinical evidence in light of key outcomes and comparisons of interest, the report determined that the evidence for the use of TOT when added to standard of care is favorable. Additional randomized controlled trials are needed to determine the best TOT application method, and additional randomized controlled trials that compare TOT with other treatments are needed to address evidence gaps.

Professional Societies/Organizations

American Diabetes Society

The American Diabetes Association guideline, Retinopathy, Neuropathy, and Foot Care: Standards of Care in Diabetes-2025, includes the ADA's current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. The guideline states that TOT has been studied rather vigorously in recent years, with several high-quality randomized controlled trials and at least five systematic reviews, all supporting its efficacy in healing chronic DRU's at twelve weeks. The authors note that, importantly, TOT devices provide home-based therapy, replacing the need for daily visits to specialized centers. The guideline further states that very high participation with very few reported adverse events combined with improved healing rates makes this therapy an attractive option for advanced wound care. The guideline includes the following recommendation:

- For chronic diabetic foot ulcers that have failed to heal with optimal standard care alone, adjunctive treatment with randomized controlled trial-proven advanced agents should be considered. Considerations might include negative pressure wound therapy, placental membranes, bioengineered skin substitutes, several acellular matrices, autologous fibrin and leukocyte platelet patches, and topical oxygen therapy. (Evidence grade: A)

Wound Healing Society

Guidelines from the Wound Healing Society, updated in 2023, state that topical oxygen therapy has been shown to increase the incidence of healing and decrease the time to heal (Level I). This is based on the principle that oxygen is essential to promote wound healing. Topical oxygen diffuses oxygen into the ulcer wound bed. The rationale for the use of topical oxygen is to eliminate hypoxia and stimulate growth factors that contribute to angiogenesis.

International Working Group on Diabetic Foot (IWGDF)

IWGDF Guidelines on Interventions to Enhance the Healing of Foot Ulcers in People with Diabetes, part of the 2023 IWGDF Guidelines on the Prevention and Management of Diabetes-Related Foot Disease, include a list of recommendations that should be considered to be adjunctive to best standard of care when best standard of care alone has failed to heal the ulcers. This should include sharp debridement and basic wound dressings, which according to IWGDF Practical

Guidelines, should be dressings to absorb exudate and maintain a moist wound healing environment. The guideline includes the following recommendation:

- Consider the use of topical oxygen as an adjunct therapy to standard of care for wound healing in people with diabetes-related foot ulcers where standard of care alone has failed and resources exist to support this intervention. (Conditional; Low)

The strength of recommendations are ranked as either strong or weak/conditional. A strong recommendation indicates that most individuals should receive the recommended course of action. A conditional recommendation indicates that the best action may differ depending on patient values or circumstances. The quality of evidence is ranked as high, moderate, low or very low.

Undersea and Hyperbaric Medical Society (UHMS)

Regarding topical oxygen, UHMS (Meng et al., 2018; Feldmeier, et al., 2005) stated that topical oxygen is not hyperbaric nor is it equivalent to HBO. Outcomes from clinical studies evaluating HBO cannot be applied to topical oxygen. UHMS does not recommend the use of THBO outside of clinical trials.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD	National	Hyperbaric Oxygen Therapy (20.29)	12/18/2017
LCD	CGS Administrators	Oxygen and oxygen equipment (L33797).	4/01/2023

Note: Please review the current Medicare Policy for the most up-to-date information.
(NCD = National Coverage Determination; LCD = Local Coverage Determination)

Coding Information

Notes:

- This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare & Medicaid Services (CMS) code updates may occur more frequently than policy updates.
- Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Systemic Hyperbaric Oxygen Therapy

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
99183	Physician or other qualified health care professional attendance and supervision of hyperbaric oxygen therapy, per session

HCPCS Codes	Description
G0277	Hyperbaric oxygen under pressure, full body chamber, per 30 minute interval

Topical Oxygen Therapy

Considered Medically Necessary when topical oxygen therapy (TOT) is used as an adjunct treatment of a diabetic foot ulcer that has failed to heal with optimal standard of care:

HCPCS Codes	Description
A4575	Topical hyperbaric oxygen chamber, disposable

***Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.**

References

1. Ajduk J, Ries M, Trotic R, Marinac I, Vlatka K, Bedekovic V. Hyperbaric Oxygen Therapy as Salvage Therapy for Sudden Sensorineural Hearing Loss. J Int Adv Otol. 2017 Apr;13(1):61-64.
2. Al-Jalodi O, Kupcella M, Breisinger K, Serena TE. A Multicenter clinical trial evaluating the durability of diabetic foot ulcer healing in ulcers treated with topical oxygen and standard of care versus standard of care alone 1 year post-healing. Int Wound J. 2022;19(7):1838-1842. doi:10.1111/wj.13789
3. Alimoglu Y, Inci E, Edizer DT, Ozdilek A, Aslan M. Efficacy comparison of oral steroid, intratympanic steroid, hyperbaric oxygen and oral steroid + hyperbaric oxygen treatments in idiopathic sudden sensorineural hearing loss cases. Eur Arch Otorhinolaryngol. 2011 Dec;268(12):1735-41.
4. American Academy of Otolaryngology – Head and Neck Surgery (AAO-HNS). Clinical practice guideline: sudden hearing loss. Mar 2012, updated Aug 2019. Accessed Apr 3, 2025. Available at URL address: <https://www.entnet.org/content/clinical-practice-guideline-sudden-hearing-loss-update-2019>
5. American College of Emergency Physicians (ACEP). ACEP current clinical policies. Carbon monoxide poisoning. 2017. Accessed Apr 3, 2025. Available at URL address: <https://www.acepnow.com/article/acep-clinical-policy-acute-carbon-monoxide-poisoning/>
6. American Diabetes Association Professional Practice Committee. 12. Retinopathy, neuropathy, and foot care: Standards of Care in Diabetes-2025. Diabetes Care 2025;48(Suppl. 1):5552-5265.
7. American Heart Association and the American Stroke Association. Guidelines for the early management of patients with acute ischemic stroke. 2018. Accessed Apr 7, 2025. Available at URL address: <http://stroke.ahajournals.org/content/49/3/e46>
8. Andreyev HJ, Davidson SE, Gillespie C, Allum WH, Swarbrick E; British Society of Gastroenterology; Association of Colo-Proctology of Great Britain and Ireland; Association of Upper Gastrointestinal Surgeons; Faculty of Clinical Oncology Section of the Royal College of Radiologists. Practice guidance on the management of acute and chronic gastrointestinal problems arising as a result of treatment for cancer. Gut. 2012 Feb;61(2):179-92.

9. Annane D, Chadda K, Gajdos P, Jars-Guincestre MC, Chevret S, Raphael JC. Hyperbaric oxygen therapy for acute domestic carbon monoxide poisoning: two randomized controlled trials. *Intensive Care Med.* 2011 Mar;37(3):486-92.
10. Arnold T. Spider envenomation, brown recluse. Sep 18, 2018, updated Oct 14, 2021. Accessed Apr 3, 2025. Available at URL address: <http://emedicine.medscape.com/article/772295-overview>
11. Bennett MH, Best TM, Babul-Wellar S, Taunton JE. Hyperbaric oxygen therapy for delayed onset muscle soreness and closed soft tissue injury. *Cochrane Database of Systematic Reviews* 2005, Issue 4. Art. No.: CD004713. DOI: 10.1002/14651858.CD004713.pub2.
12. Bennett MH, Feldmeier J, Smee R, Milross C. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *Cochrane Database of Systematic Reviews* 2018, Issue 4. Art. No.: CD005007. DOI: 10.1002/14651858.CD005007.pub4.
13. Bennett MH, French C, Schnabel A, Wasiak J, Kranke P, Weibel S. Normobaric and hyperbaric oxygen therapy for the treatment and prevention of migraine and cluster headache. *Cochrane Database of Systematic Reviews* 2015, Issue 12. Art. No.: CD005219. DOI: 10.1002/14651858.CD005219.pub3.
14. Bennett MH, Heard R. Hyperbaric oxygen therapy for multiple sclerosis. *Cochrane Database of Systematic Reviews* 2004, Issue 1. Art. No.: CD003057. DOI: 10.1002/14651858.CD003057.pub2.
15. Bennett M, Kertesz T, Perleth M, Yeung P, Lehm JP. Hyperbaric oxygen therapy for idiopathic sudden sensorineural hearing loss and tinnitus. *Cochrane Database of Systematic Reviews* 2012, Issue 10. Art. No.: CD004739. DOI: 10.1002/14651858.CD004739.pub4.
16. Bennett MH, Lehm JP, Jepson N. Hyperbaric oxygen therapy for acute coronary syndrome. *Cochrane Database of Systematic Reviews* 2015, Issue 7. Art. No.: CD004818. DOI: 10.1002/14651858.CD004818.pub4.
17. Bennett MH, Lehm JP, Mitchell SJ, Wasiak J. Recompression and adjunctive therapy for decompression illness. *Cochrane Database of Systematic Reviews* 2012, Issue 5. Art. No.: CD005277. DOI: 10.1002/14651858.CD005277.pub3.
18. Bennett MH, Stanford R, Turner R. Hyperbaric oxygen therapy for promoting fracture healing and treating fracture non-union. *Cochrane Database of Systematic Reviews* 2012, Issue 11. Art. No.: CD004712. DOI: 10.1002/14651858.CD004712.pub4.
19. Bennett MH, Trytko B, Jonker B. Hyperbaric oxygen therapy for the adjunctive treatment of traumatic brain injury. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No.: CD004609. DOI: 10.1002/14651858.CD004609.pub3
20. Bennett MH, Weibel S, Wasiak J, Schnabel A, French C, Kranke P. Hyperbaric oxygen therapy for acute ischaemic stroke. *Cochrane Database Syst Rev.* 2014 Nov 12;11:CD004954. doi: 10.1002/14651858.CD004954.pub3.
21. Blackman E, Moore C, Hyatt J, Railton R, Frye C. Topical wound oxygen therapy in the treatment of severe diabetic foot ulcers: a prospective controlled study. *Ostomy Wound Manage.* 2010 Jun;56(6):24-31.

22. Boulton AJM, Armstrong D, Londahl M, Frykberg RG, Game, FL, Edmonds ME, et al. New Evidence-Based Therapies for Complex Diabetic Foot Wounds. Arlington, VA, American Diabetes Association, 2022.
23. Buckley NA, Juurlink DN, Isbister G, Bennett MH, Lavonas EJ. Hyperbaric oxygen for carbon monoxide poisoning. Cochrane Database of Systematic Reviews 2011, Issue 4. Art. No.: CD002041. DOI: 10.1002/14651858.CD002041.pub3
24. Camporesi EM, Vezzani G, Bosco G, Mangar D, Bernasek TL. Hyperbaric oxygen therapy in femoral head necrosis. J Arthroplasty. 2010 Sep;25(6 Suppl):118-23.
25. Capuano L, Cavaliere M, Parente G, Damiano A, Pezzuti G, Lopardo D, Iemma M. Hyperbaric oxygen for idiopathic sudden hearing loss: is the routine application helpful? Acta Otolaryngol. 2015 Jul;135(7):692-7.
26. Cardinal J, Slade A, McFarland M, Keihani S, Hotaling JN, Myers JB. Scoping Review and Meta-analysis of Hyperbaric Oxygen Therapy for Radiation-Induced Hemorrhagic Cystitis. Curr Urol Rep. 2018 Apr 13;19(6):38.
27. Centers for Medicare and Medicaid Services (CMS). Medicare Coverage Database. Accessed Apr 7, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/indexes/lcd-alphabetical-index.aspx>
28. Cesaro S, Brugiolo A, Faraci M, Uderzo C, Rondelli R, Favre C, et al. Incidence and treatment of hemorrhagic cystitis in children given hematopoietic stem cell transplantation: a survey from the Italian association of pediatric hematology oncology-bone marrow transplantation group. Bone Marrow Transplant. 2003;32:925-31.
29. Chambrone L, Mandia J Jr, Shibli JA, Romito GA, Abrahao M. Dental implants installed in irradiated jaws: a systematic review. J Dent Res. 2013 Dec;92(12 Suppl):119S-30S. doi: 10.1177/0022034513504947.
30. Chen P, Vilorio N, Dhatariya K, Jeffcoate W, Lobmann R, McIntosh C, et al., on behalf of the International Working Group on Diabetic Foot. Guidelines on interventions to enhance healing of foot ulcers in people with Diabetes, IWGDF 2023 update. Accessed July 10, 2025. Available at URL address: <https://iwgdfguidelines.org/>
31. Chen P, Vilorio N, Dhatariya K, Jeffcoate W, Lobmann R, McIntosh C, et al. Effectiveness of interventions to enhance healing of chronic foot ulcers in diabetes: a systematic review. Diabetes Metab Res Rev. 2024;40(3):e3786.
32. Chi TH, Chiang MC, Chen RF, Yuan CH. Does the addition of hyperbaric oxygen therapy to conventional treatment modalities influence the outcome of soldiers with idiopathic sudden sensorineural hearing loss? J R Army Med Corps. 2018 May;164(2):69-71.
33. Chong KT, Hampson NB, Corman JM. Early hyperbaric oxygen therapy improves outcome for radiation-induced hemorrhagic cystitis. Urology. 2005;65:649-53.
34. Chung E. Penile Glans Necrosis Following Prostatic Artery Embolization for Benign Prostatic Hyperplasia: Case Series and Review of Current Literature. World J Mens Health. 2023 Apr;41(2):396-402. doi: 10.5534/wjmh.210244. Epub 2022 May 20. PMID: 35791297; PMCID: PMC10042657.

35. Cifu DX1, Hart BB, West SL, Walker W, Carne W. The effect of hyperbaric oxygen on persistent postconcussion symptoms. *J Head Trauma Rehabil.* 2014 Jan-Feb;29(1):11-20.
36. Conlin AE, Parnes LS. Treatment of sudden sensorineural hearing loss: I. A systematic review. *Arch Otolaryngol Head Neck Surg.* 2007 Jun;133(6):573-81.
37. Cooley WC, American Academy of Pediatrics Committee on Children with Disabilities. Providing a primary care medical home for children and youth with cerebral palsy. *Pediatrics.* 2004 Oct;114(4):1106-13.
38. Cornely OA, Arikan-Akdagli S, Dannaoui E, et al; European Society of Clinical Microbiology and Infectious Diseases Fungal Infection Study Group; European Confederation of Medical Mycology. ESCMID and ECMM joint clinical guidelines for the diagnosis and management of mucormycosis 2013. *Clin Microbiol Infect.* 2014;20 Suppl 3:5-26.
39. Crawford C, Teo L, Yang E, Isbister C, Berry K. Is Hyperbaric Oxygen Therapy Effective for Traumatic Brain Injury? A Rapid Evidence Assessment of the Literature and Recommendations for the Field. *J Head Trauma Rehabil.* 2017 May/Jun;32(3):E27-E37.
40. Crawford PE, Fields-Varnado M. Guideline for the management of wounds in patients with lower-extremity neuropathic disease: an executive summary. *J Wound Ostomy Continence Nurs.* 2013 Jan-Feb;40(1):34-45.
41. Cvorovic L, Jovanovic MB, Milutinovic Z, Arsovic N, Djerić D. Randomized prospective trial of hyperbaric oxygen therapy and intratympanic steroid injection as salvage treatment of sudden sensorineural hearing loss. *Otol Neurotol.* 2013 Aug;34(6):1021-6.
42. Dauwe PB1, Pulikkottil BJ, Lavery L, Stuzin JM, Rohrich RJ. Does hyperbaric oxygen therapy work in facilitating acute wound healing: a systematic review. *Plast Reconstr Surg.* 2014 Feb;133(2):208e-15e.
43. Dekleva M, Neskovic A, Vlahovic A, Putnikovic B, Beleslin B, Ostojic M. Adjunctive effect of hyperbaric oxygen treatment after thrombolysis on left ventricular function in patients with acute myocardial infarction. *AHJ.* 2004 Oct;148(4).
44. Dulai PS, Gleeson MW, Taylor D, et al. Systematic review: The safety and efficacy of hyperbaric oxygen therapy for inflammatory bowel disease. *Aliment Pharmacol Ther.* 2014;39(11):1266-1275.
45. ECRI. Topical oxygen therapy for diabetic foot ulcers. Clinical evidence assessment. 2025 June.
46. Edsberg LE, Brogan MS, Jaynes CD and Fries K. Topical hyperbaric oxygen and electrical stimulation: exploring potential synergy. *Ostomy/Wound Management.* 2002;48(11):42-50.
47. Efrati S, Fishlev G, Bechor Y, Volkov O, Bergan J, Kliakhandler K, Kamiager I, Gal N, Friedman M, Ben-Jacob E, Golan H. Hyperbaric oxygen induces late neuroplasticity in post stroke patients--randomized, prospective trial. *PLoS One.* 2013;8(1):e53716.

48. Eftedal OS, Lydersen S, Helde G, White L, Brubakk AO, Stovner LJ. A randomized, double blind study of the prophylactic effect of hyperbaric oxygen therapy on migraine. *Cephalalgia*. 2004 Aug;24(8):639-44.
49. El-Zimaity M, Saliba R, Chan K, Shahjahan M, Carrasco A, Khorshid O, et al. Hemorrhagic cystitis after allogeneic hematopoietic stem cell transplantation: donor type matters. *Blood*. 2004 Jun;103(12):4674-80.
50. Eryigit B, Ziyilan F, Yaz F, Thomeer HGXM. The effectiveness of hyperbaric oxygen in patients with idiopathic sudden sensorineural hearing loss: a systematic review. *Eur Arch Otorhinolaryngol*. 2018 Dec;275(12):2893-2904.
51. Eskes A, Vermeulen H, Lucas C, Ubbink DT. Hyperbaric oxygen therapy for treating acute surgical and traumatic wounds. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No.: CD008059. DOI: 10.1002/14651858.CD008059.pub3.
52. Eskes AM, Ubbink DT, Lubbers MJ, Lucas C, Vermeulen H. Hyperbaric oxygen therapy: solution for difficult to heal acute wounds? Systematic review. *World J Surg*. 2011 Mar;35(3):535-42.
53. Esposito M, Worthington HV. Interventions for replacing missing teeth: hyperbaric oxygen therapy for irradiated patients who require dental implants. *Cochrane Database of Systematic Reviews* 2013, Issue 9. Art. No.: CD003603. DOI: 10.1002/14651858.CD003603.pub3
54. Feldmeier JJ, Hopf HW, Warriner RA 3rd, Fife CE, Gesell LB, Bennett M. UHMS position statement: topical oxygen for chronic wounds. *Undersea Hyperb Med*. 2005 May-Jun;32(3):157-68.
55. Feng JJ, Li YH. Effects of hyperbaric oxygen therapy on depression and anxiety in the patients with incomplete spinal cord injury (a STROBE-compliant article). *Medicine (Baltimore)*. 2017 Jul;96(29):e7334.
56. Fine HA, Barker II FG, Markert JM, Loeffler JS. (authors). Chapter 39: Neoplasms of the central nervous system. In: DeVita, *CANCER Principles & Practice of Oncology*. Philadelphia, PA: Lippincott Williams & Wilkins; 2005.
57. Fink D, Chetty N, Lehm JP, Marsden DE, Hacker NF. Hyperbaric oxygen therapy for delayed radiation injuries in gynecological cancers. *Int J Gynecol Cancer*. 2006;16:638-42.
58. Fox NF, Xiao C, Sood AJ, Lovelace TL, Nguyen SA, Sharma A, Day TA. Hyperbaric oxygen therapy for the treatment of radiation-induced xerostomia: a systematic review. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2015 Jul;120(1):22-8.
59. Fritz GW, Gunsolley JC, Abubaker O, Laskin DM. Efficacy of pre- and postirradiation hyperbaric oxygen therapy in the prevention of postextraction osteoradionecrosis: a systematic review. *J Oral Maxillofac Surg*. 2010 Nov;68(11):2653-60.
60. Frykberg RG, Franks PJ, Edmonds M, Brantley JN, Téot L, Wild T, Garoufalidis MG, Lee AM, Thompson JA, Reach G, Dove CR, Lachgar K, Grotemeyer D, Renton SC; TWO2 Study Group. A Multinational, Multicenter, Randomized, Double-Blinded, Placebo-Controlled Trial to Evaluate the Efficacy of Cyclical Topical Wound Oxygen (TWO2) Therapy in the

Treatment of Chronic Diabetic Foot Ulcers: The TWO2 Study. *Diabetes Care*. 2020 Mar;43(3):616-624. doi: 10.2337/dc19-0476. Epub 2019 Oct 16. PMID: 31619393.

61. Gaitanou K, Fildissis G, Vavasis P, Kalentzos V, Baltopoulos G. Management of sudden hearing loss with hyperbaric oxygen therapy. *Undersea Hyperb Med*. 2014 Sep-Oct;41(5):363-70.
62. Ghanizadeh A. Hyperbaric oxygen therapy for treatment of children with autism: a systematic review of randomized trials. *Med Gas Res*. 2012 May 11;2:13. doi: 10.1186/2045-9912-2-13.
63. Glover M, Smerdon GR, Andreyev HJ, Benton BE, Bothma P, Firth O, Gothard L, Harrison J, Ignatescu M, Laden G, Martin S, Maynard L, McCann D, Penny CE, Phillips S, Sharp G, Yarnold J. Hyperbaric oxygen for patients with chronic bowel dysfunction after pelvic radiotherapy (HOT2): a randomised, double-blind, sham-controlled phase 3 trial. *Lancet Oncol*. 2016 Feb;17(2):224-33.
64. Goldman RJ. Hyperbaric oxygen therapy for wound healing and limb salvage: a systematic review. *PM R*. 2009 May;1(5):471-89.
65. Gordillo GM, Roy S, Khanna S, Schlanger R, Khandelwal S, Phillips G, et al. Topical oxygen therapy induces vascular endothelial growth factor expression and improves closure of clinically presented chronic wounds. *Clin Exp Pharmacol Physiol*. 2008 Apr 21.
66. Gorenstein SA, Castellano ML, Slone ES, Gillette B, Liu H, Alsamarraie C, Jacobson AM, Wall SP, Adhikari S, Swartz JL, McMullen JJS, Osorio M, Koziatsek CA, Lee DC. Hyperbaric oxygen therapy for COVID-19 patients with respiratory distress: treated cases versus propensity-matched controls. *Undersea Hyperb Med*. 2020 Third-Quarter;47(3):405-413. doi: 10.22462/01.03.2020.1. PMID: 32931666.
67. Gothard L, Haviland J, Bryson P, Laden G, Glover M, Harrison S, Woods M, Cook G, Peckitt C, Pearson A, Somaiah N, Stanton A, Mortimer P, Yarnold J. Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer. *Radiother Oncol*. 2010 Oct;97(1):101-7.
68. Guo D, Pan S, Wang M, Guo Y. Hyperbaric oxygen therapy may be effective to improve hypoxemia in patients with severe COVID-2019 pneumonia: two case reports. *Undersea Hyperb Med*. 2020 Second-Quarter;47(2):181-187. doi: 10.22462/04.06.2020.2. PMID: 32574433.
69. Hadanny A, Zilberman-Itskovich S, Catalogna M, Elman-Shina K, Lang E, Finci S, Polak N, Shorer R, Parag Y, Efrati S. Long term outcomes of hyperbaric oxygen therapy in post covid condition: longitudinal follow-up of a randomized controlled trial. *Sci Rep*. 2024 Feb 15;14(1):3604. doi: 10.1038/s41598-024-53091-3. PMID: 38360929; PMCID: PMC10869702.
70. Hailey D. Hyperbaric oxygen therapy: recent findings on evidence for its effectiveness. Alberta Heritage Foundation for Medical Research (AHFMR). IP 13: Information paper. 2003 Mar.
71. Hawkins JR, Gonzalez KE, Heumann KJ. The Effectiveness of Hyperbaric Oxygen Therapy as a Treatment for Postconcussion Symptoms. *J Sport Rehabil*. 2017 May;26(3):290-294.

72. Headache Classification Committee of the International Headache Society (IHS) The International Classification of Headache Disorders, 3rd edition. Cephalalgia. 2018 Jan;38(1):1-211. doi: 10.1177/0333102417738202. PMID: 29368949.
73. Health Quality Ontario. Hyperbaric Oxygen Therapy for the Treatment of Diabetic Foot Ulcers: A Health Technology Assessment. Ont Health Technol Assess Ser. 2017 May 12;17(5):1-142. eCollection 2017.
74. Holland NJ, Bernstein JM, Hamilton JW. Hyperbaric oxygen therapy for Bell's palsy. Cochrane Database of Systematic Reviews 2012, Issue 2. Art. No.: CD007288. DOI: 10.1002/14651858.CD007288.pub2. Assessed as up to date July 6, 2016.
75. Holy R, Navara M, Dosel P, Fundova P, Prazenica P, Hahn A. Hyperbaric oxygen therapy in idiopathic sudden sensorineural hearing loss (ISSNHL) in association with combined treatment. Undersea Hyperbaric Med. 2011 Mar-Apr;38(2):137-42.
76. Horn CE, Himel HN, Selesnick SH. Hyperbaric oxygen therapy for sudden sensorineural hearing loss: a prospective trial of patients failing steroid and antiviral treatment. Otol Neurotol. 2005 Sep;26(5):882-9.
77. Huang ET, Mansouri J, Murad MH, Joseph WS, Strauss MB, Tettelbach W, Worth ER; UHMS CPG Oversight Committee. A clinical practice guideline for the use of hyperbaric oxygen therapy in the treatment of diabetic foot ulcers. Undersea Hyperb Med. 2015 May-Jun;42(3):205-47.
78. Javier NM. Pressure Injury. In: Ferri's Clinical Advisor 2024. Philadelphia, PA: Elsevier, 2024.
79. Juang J-H, Hsu BR-S, Kuo C-H, Ueng SW-N. Beneficial effects of hyperbaric oxygen therapy on Islet Transplantation. Cell Transplantation. 2002;11:95-101.
80. Kalayoglu-Besisik A, Abdul-Rahman IS, Erer B, Yenerel MN, Oguz FS, Tunc M, et al. Outcome after hyperbaric oxygen treatment for cyclophosphamide-induced refractory hemorrhagic cystitis. J Urol. 2003 Sep;170:922.
81. Kalliainen LK, Gordillo GM, Schlanger R, Sen CK. Topical oxygen as an adjunct to wound healing: a clinical case series. Pathophysiology. 2003 Jan;9(2):81-87.
82. Keohane C, Westby D, Nolan FC, Twyford M, Tawfick W, Walsh SR. Hyperbaric Oxygen as an Adjunct in the Treatment of Venous Ulcers: A Systematic Review. Vasc Endovascular Surg. 2023 Mar 8;15385744231162924. doi: 10.1177/15385744231162924. Epub ahead of print. PMID: 36891617.
83. Kiralp MZ, Yildiz S, Vural D, Keskin I, Ay H, Dursun H. Effectiveness of hyperbaric oxygen therapy in the treatment of complex regional pain syndrome. J Int Med Res. 2004 May-Jun;32(3):258-62.
84. Kranke P, Bennett MH, Martyn-St James M, Schnabel A, Debus SE, Weibel S. Hyperbaric oxygen therapy for chronic wounds. Cochrane Database of Systematic Reviews 2015, Issue 6. Art. No.: CD004123. DOI: 10.1002/14651858.CD004123.pub4.
85. Lamm K, Lamm H, Arnold W. Effect of hyperbaric oxygen therapy in comparison to conventional or placebo therapy or no treatment in idiopathic sudden hearing loss, acoustic

- trauma, noise-induced hearing loss and tinnitus. A literature survey. *Adv Otorhinolaryngol*. 1998;54:86-99.
86. Lacey DJ, Stolfi A, Pilati LE. Effects of hyperbaric oxygen on motor function in children with cerebral palsy. *Ann Neurol*. 2012 Nov;72(5):695-703. doi: 10.1002/ana.23681.
 87. Laureau J, Pons C, Letellier G, Gross R. Hyperbaric oxygen in children with cerebral palsy: A systematic review of effectiveness and safety. *PLoS One*. 2022 Oct 14;17(10):e0276126. doi: 10.1371/journal.pone.0276126. PMID: 36240157; PMCID: PMC9565562.
 88. Lavery LA, Suludere MA, Attinger CE, et al. WHI (Wound Healing Society) guidelines update: Diabetic foot ulcer treatment guidelines. *Wound Rep Reg*. 2024;32(1):34-46. Doi:10.1111/wrr.13133
 89. Lawrence R, Thevasagayam R. Controversies in the management of sudden sensorineural hearing loss: an evidence-based review. *Clin Otolaryngol*. 2015 Jun;40(3):176-82.
 90. Lazzarini L, Mader JT, Calhoun JH. Current concepts review: Osteomyelitis in long bones. *JBJS*. 2004;86(10):2305-18.
 91. Levett D, Bennett MH, Millar I. Adjunctive hyperbaric oxygen for necrotizing fasciitis. *Cochrane Database of Systematic Reviews* 2015, Issue 1. Art. No.: CD007937. DOI: 10.1002/14651858.CD007937.pub2.
 92. Li W, Ye Z, Wang W, Wang K, Li L, Zhao D. Clinical effect of hyperbaric oxygen therapy in the treatment of femoral head necrosis : A systematic review and meta-analysis. *Orthopade*. 2017 May;46(5):440-446. English. doi: 10.1007/s00132-016-3360-8. PMID: 27928615.
 93. Lin ZC, Bennett MH, Hawkins GC, Azzopardi CP, Feldmeier J, Smee R, Milross C. Hyperbaric oxygen therapy for late radiation tissue injury. *Cochrane Database Syst Rev*. 2023 Aug 15;8(8):CD005005. doi: 10.1002/14651858.CD005005.pub5. PMID: 37585677; PMCID: PMC10426260.
 94. Liptak GS, Murphy NA; Council on Children With Disabilities. Providing a primary care medical home for children and youth with cerebral palsy. *Pediatrics*. 2011 Nov;128(5):e1321-9. doi: 10.1542/peds.2011-1468.
 95. Liu SC, Kang BH, Lee JC, Lin YS, Huang KL, Liu DW, Su WF, Kao CH, Chu YH, Chen HC, Wang CH. Comparison of therapeutic results in sudden sensorineural hearing loss with/without additional hyperbaric oxygen therapy: a retrospective review of 465 audiologically controlled cases. *Clin Otolaryngol*. 2011 Apr;36(2):121-8.
 96. Liu R, Li L, Yang M, Boden G, Yang G. Systematic review of the effectiveness of hyperbaric oxygenation therapy in the management of chronic diabetic foot ulcers. *Mayo Clin Proc*. 2013 Feb;88(2):166-75.
 97. Lodewijk W, van Eps A, de Jong PE. (authors). Chapter 77: Sick Cell Disease. In: Freedberg E, Wolff A, Goldsmith K. (editors). *Schrier: Diseases of the Kidney and Urinary Tract*. Philadelphia, PA: Lippincott Williams & Williams; 2007.

98. Markt JC, Salinas TJ, Gay WD. Ch 98 Prosthetic management of head and neck defects. In: Flint: Cummings Otolaryngology: Head & Neck Surgery, 5th ed. Mosby Maryland Heights MO. 2010.
99. Mathieu D, Marroni A, Kot J. Tenth European Consensus Conference on Hyperbaric Medicine: recommendations for accepted and non-accepted clinical indications and practice of hyperbaric oxygen treatment. *Diving Hyperb Med*. 2017 Mar;47(1):24-32.
100. McAdam AJ, Sharpe, AH. Ch 8: infectious disease. Fungal infections. In: Kumar: Robbins and Cotran: Pathologic Basis of Disease, 7th ed. St. Louis, MO: W. B. Saunders, 2005.
101. McDonagh MS, Morgan D, Carson S, Russman BS. Systematic review of hyperbaric oxygen therapy for cerebral palsy: the state of the evidence. *Dev Med Child Neurol*. 2007 Dec;49(12):942-7.
102. Mechem CC and Manaker S. Hyperbaric oxygen therapy. In: UpToDate, Schwarz E, ed. Dec 11, 2023. UpToDate, Waltham, MA. Accessed Mar 25, 2024.
103. Meng X, Sunn Q, Guo Y. UHMS Position Statement: Topical Oxygen for Chronic Wounds. *Undersea Hyperb Med*. 2018 May-Jun;45(3):379-380. PMID: 30028926.
104. Miller RS, Weaver LK, Bahraini N, Churchill S, Price RC, Skiba V, Caviness J, Mooney S, Hetzell B, Liu J, Deru K, Ricciardi R, Fracisco S, Close NC, Surrent GW, Bartos C, Ryan M, Brenner LA; HOPPS Trial Team. Effects of hyperbaric oxygen on symptoms and quality of life among service members with persistent postconcussion symptoms: a randomized clinical trial. *JAMA Intern Med*. 2015 Jan;175(1):43-52.
105. Mukherjee A, Raison M, Sahni T, Arya A, Lambert J, Marois P, James PB, Parent A, Ballaz L. Intensive rehabilitation combined with HBO2 therapy in children with cerebral palsy: a controlled longitudinal study. *Undersea Hyperb Med*. 2014 Mar-Apr;41(2):77-85.
106. Murad MH, Altayar O, Bennett M, Wei JC, Claus PL, Asi N, Prokop LJ, Montori VM, Guyatt GH. Using GRADE for evaluating the quality of evidence in hyperbaric oxygen therapy clarifies evidence limitations. *J Clin Epidemiol*. 2014 Jan;67(1):65-72.
107. Muralidharan V, Christophi C. Hyperbaric oxygen therapy and liver transplantation. *HPB (Oxford)*. 2007;9(3):174-82.
108. Nabil S, Samman N. Incidence and prevention of osteoradionecrosis after dental extraction in irradiated patients: a systematic review. *Int J Oral Maxillofac Surg*. 2011 Mar;40(3):229-43.
109. National Institute on Deafness and Other Communication Disorders (NICDC). Sudden deafness. Sept 2018. Accessed Apr 7, 2025. Available at URL address: <http://www.nidcd.nih.gov/health/hearing/pages/sudden.aspx>
110. National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers: Quick Reference Guide. Emily Haesler (Ed.). Cambridge Media: Osborne Park, Western Australia; 2019. Accessed Apr 7, 2025. Available at URL address: <https://npiap.com/news/news.asp?id=478409&hhSearchTerms=%22Prevention+and+Treatment+of+Pressure+Ulcers%22>

111. Nilsson Renahl AI, Ansjon R, Lind F, Waldenlind E. Hyperbaric oxygen treatment of active cluster headache: a double-blind placebo-controlled cross-over study. *Cephalalgia*. 2002 Nov;22(9):730-9.
112. Nogueira-Filho GR, Rosa BT, David-Neto JR. Effects of hyperbaric oxygen therapy on the treatment of severe cases of periodontitis. *Undersea Hyperb Med*. 2010 Mar-Apr;37(2):107-14.
113. Oliaei S, SeyedAlinaghi S, Mehrtak M, Karimi A, Noori T, Mirzapour P, Shojaei A, MohsseniPour M, Mirghaderi SP, Alilou S, Shobeiri P, Azadi Cheshmekabodi H, Mehraeen E, Dadras O. The effects of hyperbaric oxygen therapy (HBOT) on coronavirus disease-2019 (COVID-19): a systematic review. *Eur J Med Res*. 2021 Aug 19;26(1):96. doi: 10.1186/s40001-021-00570-2. PMID: 34412709; PMCID: PMC8374420.
114. Oropallo A, Armstrong DG. Local care of diabetic foot ulcers. In: UpToDate, Coollins KA, (Ed), UpToDate, Waltham, MA. Topic last updated Mar 20, 2025. Accessed Jul 7, 2025.
115. OuYang H, Yang J, Wan H, Huang J, Yin Y (2024) Effects of different treatment measures on the efficacy of diabetic foot ulcers: a network meta-analysis. *Front. Endocrinol*. 15:1452192. doi: 10.3389/fendo.2024.145192
116. Pacek J, Szajkowski S, Cieslar G. Physical treatment of diabetic foot ulcers-preliminary study for topical application of oxygen or ozone auxiliary treatment of diabetic foot ulcers. *Dermatologic Therapy*. 2023; 2023:1 article number: 1843995
117. Paderno E, Zanon V, Vezzani G, Giacon TA, Bernasek TL, Camporesi EM, Bosco G. Evidence-Supported HBO Therapy in Femoral Head Necrosis: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health*. 2021 Mar 12;18(6):2888. doi: 10.3390/ijerph18062888. PMID: 33808951; PMCID: PMC7999152.
118. Pezzoli M1, Magnano M, Maffi L, Pezzoli L, Marcato P, Orione M, Cupi D, Bongioannini G. Hyperbaric oxygen therapy as salvage treatment for sudden sensorineural hearing loss: a prospective controlled study. *Eur Arch Otorhinolaryngol*. 2015 Jul;272(7):1659-66.
119. Phillips JS, Jones SEM. Hyperbaric oxygen as an adjuvant treatment for malignant otitis externa. *Cochrane Database of Systematic Reviews* 2013, Issue 5. Art. No.: CD004617. DOI: 10.1002/14651858.CD004617.pub3
120. Porubsky C, Stiegler P, Matzi V, Lipp C, Kontaxis A, Klemen H, Walch C, Smolle-Juttner F. Hyperbaric oxygen in tinnitus: influence of psychological factors on treatment results? *ORL J Otorhinolaryngol Relat Spec*. 2007;69(2):107-12.
121. Rao SSC. Practice Guidelines: Diagnosis and management of fecal incontinence. *Am J Gastro*.2004: 1585-1604.
122. Reis ND, Schwartz O, Militianu D, Ramon Y, Levin D, Norman D, Melamed Y, Shupak A, Goldsher D, Zinman C. Hyperbaric oxygen therapy as a treatment for stage-I avascular necrosis of the femoral head. *J Bone Joint Surg Br*. 2003 Apr;85(3):371-5.
123. Robbins T, Gonevski M, Clark C, Baitule S, Sharma K, Magar A, Patel K, Sankar S, Kyrou I, Ali A, Randeva HS. Hyperbaric oxygen therapy for the treatment of long COVID: early

evaluation of a highly promising intervention. *Clin Med (Lond)*. 2021 Nov;21(6):e629-e632. doi: 10.7861/clinmed.2021-0462. PMID: 34862223; PMCID: PMC8806311.

124. Roberts, JR. Chronic Fatigue Syndrome. Sep 7, 2020, updated Sep 6, 2024. Accessed Apr 3, 2025. Available at URL address: <http://emedicine.medscape.com/article/235980-overview>
125. Rockswold SB, Rockswold GL, Zaun DA, Liu J. A prospective, randomized Phase II clinical trial to evaluate the effect of combined hyperbaric and normobaric hyperoxia on cerebral metabolism, intracranial pressure, oxygen toxicity, and clinical outcome in severe traumatic brain injury. *J Neurosurg*. 2013 Jun;118(6):1317-28.
126. Roeckl-Wiedmann I, Bennett M, Kranke P. Systematic review of hyperbaric oxygen in the management of chronic wounds. *Br J Surg*. 2005 Jan;92(1):24-32.
127. Rollason V, Laverrière A, MacDonald LCI, Walsh T, Tramèr MR, Vogt-Ferrier NB. Interventions for treating bisphosphonate- related osteonecrosis of the jaw (BRONJ). *Cochrane Database of Systematic Reviews* 2016, Issue 2. Art. No.: CD008455. DOI: 10.1002/14651858.CD008455.pub2.
128. Rossignol DA, Rossignol LW. Hyperbaric oxygen therapy may improve symptoms in autistic children. *Med Hypotheses*. 2006;67(2):216-28.
129. Rossignol DA, Rossignol LW, James SJ, Melnyk S, Mumper E. The effects of hyperbaric oxygen therapy on oxidative stress, inflammation, and symptoms in children with autism: an open-label pilot study. *BMC Pediatr*. 2007 Nov 16;7:36.
130. Rossignol DA, Rossignol LW, Smith S, Schneider C, Logerquist S, Usman A, Neubrandner J, Madren EM, Hintz G, Grushkin B, Mumper EA. Hyperbaric treatment for children with autism: a multicenter, randomized, double-blind, controlled trial. *BMC Pediatr*. 2009 Mar 13;9:21.
131. Rowland LP, editor. *Merritt's neurology*. 11th ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2005.
132. Ruha AM. Spider bites and scorpion stings. In: Kellerman and Rakel: *Conn's Current Therapy*. Conn's Current Therapy 2024, 1479-1481.
133. Sakulchit T, Ladish C, Goldman RD. Hyperbaric oxygen therapy for children with autism spectrum disorder. *Can Fam Physician*. 2017 Jun;63(6):446-448.
134. Schwartz DA, Herdman CR. Review article: the medical treatment of Crohn's perianal fistulas. *Aliment Pharmacol Ther*. 2004;19:953-67.
135. Sevil E, Bercin S, Muderris T, Gul F, Kiris M. Comparison of two different steroid treatments with hyperbaric oxygen for idiopathic sudden sensorineural hearing loss. *Eur Arch Otorhinolaryngol*. 2016 Sep;273(9):2419-26. doi: 10.1007/s00405-015-3791-6.
136. Sideras K, Loprinzi CL, Foote RL. Ch 40 Oral complications. In: *Abeloff: Abeloff's Clinical Oncology*, 4th ed. Churchill Livingstone. 2008.

137. Simancas-Racines D, Arevalo-Rodriguez I, Osorio D, Franco JVA, Xu Y, Hidalgo R. Interventions for treating acute high altitude illness. Cochrane Database of Systematic Reviews 2018, Issue 6. Art. No.: CD009567. DOI: 10.1002/14651858.CD009567.pub2.
138. Smith J, Seirafi J. Ch 102 – Delirium and dementia. Dementia. In: Marx: Rosen's Emergency Medicine: Concepts and Clinical Practice, 6th ed. St. Louis, MO. Elsevier; 2006.
139. Society for Vascular Surgery. Clinical practice guidelines. The Management of Diabetic Foot: A Clinical Practice Guideline by the Society for Vascular Surgery in Collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. 2016. Accessed Apr 3, 2025. Available at URL address: [https://www.jvascsurg.org/article/S0741-5214\(15\)02025-X/fulltext](https://www.jvascsurg.org/article/S0741-5214(15)02025-X/fulltext)
140. Song KX, Liu S, Zhang MZ, Liang WZ, Liu H, Dong XH, Wang YB, Wang XJ. Hyperbaric oxygen therapy improves the effect of keloid surgery and radiotherapy by reducing the recurrence rate. J Zhejiang Univ Sci B. 2018 Nov.;19(11):853-862.
141. Sun X-K, Li R, Yang X-L, Yuan L. Efficacy and safety of topical oxygen therapy for diabetic foot ulcers: an updated systematic review and meta-analysis. Wound J. 2022;19(8):2200-2209.
142. Sutherland AM, Clarke HA, Katz J, Katznelson R. Hyperbaric Oxygen Therapy: A New Treatment for Chronic Pain Pain Pract. 2016 Jun;16(5):620-8.
143. Suzuki H, Hashida K, Nguyen KH, Hohchi N, Katoh A, Koizumi H, Ohbuchi T. Efficacy of intratympanic steroid administration on idiopathic sudden sensorineural hearing loss in comparison with hyperbaric oxygen therapy. Laryngoscope. 2012 May;122(5):1154-7. doi: 10.1002/lary.23245.
144. Tang X, Yin X, Zhang T, Peng H. The effect of hyperbaric oxygen on clinical outcome of patients after resection of meningiomas with conspicuous peritumoral brain edema. Undersea Hyperb Med. 2011 Mar-Apr;38(2):109-15.
145. Tawfick W, Sultan S. Does topical wound oxygen (TWO2) offer an improved outcome over conventional compression dressings (CCD) in the management of refractory venous ulcers (RVU)? A parallel observational comparative study. Eur J Vasc Endovasc Surg. 2009 Jul;38(1):125-32.
146. Teas J, Cunningham JE, Cone L, Jansen K, Raghavan SK, Nitcheva DK, et al. Can hyperbaric oxygen therapy reduce breast cancer treatment-related lymphedema? A pilot study. J Women's Health. 2004;13(9):1008-18.
147. Thibodeaux K, Speyrer M, Raza A, Yaakov R, Serena TE. Hyperbaric oxygen therapy in preventing mechanical ventilation in COVID-19 patients: a retrospective case series. J Wound Care. 2020 May 1;29(Sup5a):S4-S8. doi: 10.12968/jowc.2020.29.Sup5a.S4. PMID: 32412891.
148. Thom SR. Hyperbaric oxygen: its mechanisms and efficacy. Plast Reconstr Surg. 2011 Jan;127 Suppl 1:131S-141S.
149. Tong B, Niu K, Ku W, et al. Comparison of therapeutic results with/without additional hyperbaric oxygen therapy in idiopathic sudden sensorineural hearing loss: A randomized prospective study. Audiology & neuro-otology. 2021;26(1):11-16.

150. Undersea and Hyperbaric Medicine Society (UHMS). Position statements. Hyperbaric Oxygen (HBO2) for COVID-19 Patients. Aug 22, 2020. Accessed Apr 3, 2025. Available at URL address: <https://www.uhms.org/resources/featured-resources/position-statements.html>
151. Undersea and Hyperbaric Medicine Society (UHMS). Idiopathic sudden sensorineural hearing loss (ISSHL). 2011. Accessed Apr 3, 2025. Available at URL address: <https://www.uhms.org/14-idiopathic-sudden-sensorineural-hearing-loss-new-approved-on-october-8-2011-by-the-uhms-board-of-directors.html>
152. Undersea and Hyperbaric Medicine Society (UHMS). Indications for hyperbaric oxygen therapy, 15 ed. 2023. Accessed Apr 3, 2025. Available at URL address: <https://www.uhms.org/images/indications/UHMS%20Indications%2015th%20Ed%20Front%20Matter%20and%20References.pdf>
153. Undersea and Hyperbaric Medicine Society (UHMS). Position statements. The treatment of autism spectrum disorder with HBOT. Dec 5, 2009. Accessed Apr 3, 2025. Available at URL address: https://www.uhms.org/images/Position-Statements/autism_position_paper.pdf
154. United States Food and Drug Administration (FDA). 510(k) summary K030418. Makai Marine Multiplace Hyperbaric Chamber. Jan 29, 2004. Accessed Apr 7, 2025. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf3/K030418.pdf
155. United States Food and Drug Administration (FDA). 510(k) summary (K060739). OxyHeal 1000 Monoplace Hyperbaric Chamber. Dec 13, 2005. Accessed Apr 3, 2025. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf5/K052866.pdf
156. United States Food and Drug Administration (FDA). 510(k) summary (K080966). Hyper-Box Topical Wound Oxygen System. Aug 6, 2008. Accessed Apr 3, 2025. Available at URL address: http://www.accessdata.fda.gov/cdrh_docs/pdf8/K080966.pdf
157. United States Food and Drug Administration (FDA). 510(k) summary (K971507). O2Boot™. Nov 7, 1997. Accessed Mar 31, 2025. Available at URL address: https://www.accessdata.fda.gov/cdrh_docs/pdf/K971507.pdf
158. United States Food and Drug Administration (FDA). Hyperbaric oxygen therapy: Get the facts. July 26, 2021. Accessed Mar 31, 2025. Available at URL address: https://www.fda.gov/consumers/consumer-updates/hyperbaric-oxygen-therapy-get-facts?utm_medium=email&utm_source=govdelivery
159. Uzun G, Mutluoglu M, Ersen O, Yildiz S. Hyperbaric oxygen therapy in the treatment of osteonecrosis of the femoral head: a review of the current literature. *Undersea Hyperb Med.* 2016 May-Jun;43(3):189-99.
160. Van Hoesen KB. Ch 78 Hyperbaric medicine. In: Auerbach: Wilderness Medicine, 6th ed. Mosby. Maryland Heights Mo. 2011.
161. Van Voorhis BJ, Greensmith JE, Dokras A, Sparks AET, Simmons ST, Syrop CH. Hyperbaric oxygen and ovarian follicular stimulation for in vitro fertilization: a pilot study. *Fertil Steril.* 2005;83:226-8.

162. Wang C, Schwaitzberg S, Berliner E, Zarin DA, Lau J. Hyperbaric oxygen for treating wounds. *Arch Surg*. 2003;138:272-9.
163. Wang F, Wang Y, Sun T, Yu HL. Hyperbaric oxygen therapy for the treatment of traumatic brain injury: a meta-analysis. *Neurol Sci*. 2016 May;37(5):693-701.
164. Wasserman GS. Chapter 348: Brown Recluse Spider Envenomation. In: Harwood-Nuss' *Clinical Practice of Emergency Medicine*. 4th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2005.
165. Weaver LK, editor. Undersea and Hyperbaric Medical Society. *Hyperbaric oxygen indications*. 13th ed. Florida: Best Publishing Company; 2014.
166. Weber PC. Sudden sensorineural hearing loss in adults. In: UpToDate, Deschler DG, ed. Oct 2, 2024. UpToDate, Waltham, MA. Accessed Apr 3, 2025.
167. Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, Steere AC, Klempner MS, Krause PJ, Bakken JS, Strle F, Stanek G, Bockenstedt L, Fish D, Dumler JS, Nadelman RB. The clinical assessment, treatment, and prevention of lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 2006 Nov 1;43(9):1089-134.
168. Wu Z, Cai J, Chen J, Huang L, Wu W, Luo F, Wu C, Liao L, Tan J. Autologous bone marrow mononuclear cell infusion and hyperbaric oxygen therapy in type 2 diabetes mellitus: an open-label, randomized controlled clinical trial. *Cytotherapy*. 2014 Feb;16(2):258-65.
169. Xiao Y, Wang J, Jiang S, Luo H. Hyperbaric oxygen therapy for vascular dementia. *Cochrane Database of Systematic Reviews* 2012, Issue 7. Art. No.: CD009425. DOI: 10.1002/14651858.CD009425.pub2
170. Xiong T, Chen H, Luo R, Mu D. Hyperbaric oxygen therapy for people with autism spectrum disorder (ASD). *Cochrane Database of Systematic Reviews* 2016, Issue 10. Art. No.: CD010922. DOI: 10.1002/14651858.CD010922.pub2.
171. Xu Q, Fan SB, Wan YL, Liu XL, Wang L. The potential long-term neurological improvement of early hyperbaric oxygen therapy on hemorrhagic stroke in the diabetics. *Diabetes Res Clin Pract*. 2018 Apr;138:75-80.
172. Yildiz H, Senol L, Ercan E, Bilgili ME, Karabudak Abuaf O. A prospective randomized controlled trial assessing the efficacy of adjunctive hyperbaric oxygen therapy in the treatment of hidradenitis suppurativa. *Int J Dermatol*. 2016 Feb;55(2):232-7.
173. Yogaratnam JZ, Laden G, Guvendik L, Cowen M, Cale A, Griffin S. Hyperbaric oxygen preconditioning improves myocardial function, reduces length of intensive care stay, and limits complications post coronary artery bypass graft surgery. *Cardiovasc Revasc Med*. 2010 Jan-Mar;11(1):8-19.
174. You Q, Li L, Xiong SQ, Yan YF, Li D, Yan NN, Chen HP, Liu YP. Meta-analysis on the efficacy and safety of hyperbaric oxygen as adjunctive therapy for vascular dementia. *Front Aging Neurosci*. 2019 Apr 17;11:86. doi: 10.3389/fnagi.2019.00086. eCollection 2019.
175. Zilberman-Itskovich S, Catalogna M, Sasson E, Elman-Shina K, Hadanny A, Lang E, Finci S, Polak N, Fishlev G, Korin C, Shorer R, Parag Y, Sova M, Efrati S. Hyperbaric oxygen therapy

improves neurocognitive functions and symptoms of post-COVID condition: randomized controlled trial. Sci Rep. 2022 Jul 12;12(1):11252. doi: 10.1038/s41598-022-15565-0. PMID: 35821512; PMCID: PMC9276805.

Revision Details

Type of Revision	Summary of Changes	Date
Focused Review	Changed from not covered to covered: topical oxygen therapy as adjunct treatment for a diabetic foot ulcer that has failed to heal with optimal standard of care.	9/15/2025
Annual review	- Changed from not covered to covered: avascular necrosis - Added not covered: penile glans necrosis	5/15/2025
Annual review	No clinical policy statement changes.	5/15/2024

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.