



Medical Coverage Policy

Effective Date11/15/2025

Next Review Date11/15/2026

Coverage Policy Number..... 0103

Gastric Pacing/Gastric Electrical Stimulation (GES)

Table of Contents

Overview	2
Coverage Policy	2
Coding Information	2
General Background	3
Health Equity Considerations.....	10
Medicare Coverage Determinations	11
References.....	11
Revision Details.....	15

Related Coverage Resources

[Bariatric Surgery and Procedures](#)
[Electrical Stimulation Therapy and Devices in a Home Setting](#)
[Minimally Invasive Anti-Reflux Procedures and Peroral Endoscopic Myotomy \(POEM\) Procedures](#)
[Infant Nutritional Formula](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers

must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses gastric electrical stimulation (GES) for the treatment of intractable nausea and vomiting secondary to gastroparesis.

Coverage Policy

Permanent gastric electrical stimulation (GES) or gastric pacing (e.g., Enterra™ Therapy) is considered medically necessary when provided in accordance with the Humanitarian Device Exemption (HDE) specifications of the U.S. Food and Drug Administration (FDA) for intractable nausea and vomiting secondary to gastroparesis with failure, contraindication, or intolerance of pharmaceutical therapy.

Gastric electrical stimulation (GES) or gastric pacing for any other indication is not covered or reimbursable.

Temporary gastric electrical stimulation (GES) is considered experimental, investigational or unproven.

Coding Information

Notes:

1. This list of codes may not be all-inclusive since the American Medical Association (AMA) and Centers for Medicare & Medicaid Services (CMS) code updates may occur more frequently than policy updates.
2. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
43881	Implantation or replacement of gastric neurostimulator electrodes, antrum, open
64590	Insertion or replacement of peripheral or gastric neurostimulator pulse generator or receiver, direct or inductive coupling

HCPCS Codes	Description
C1767	Generator, neurostimulator (implantable), nonrechargeable
C1778	Lead, neurostimulator (implantable)
L8679	Implantable neurostimulator, pulse generator, any type
L8680	Implantable neurostimulator electrode, each

HPCS Codes	Description
L8688	Implantable neurostimulator pulse generator, dual array, nonrechargeable, includes extension

Considered Experimental/Investigational/Unproven when used to report temporary gastric electric stimulation:

CPT®* Codes	Description
43659	Unlisted laparoscopy procedure, stomach
43999	Unlisted procedure, stomach

***Current Procedural Terminology (CPT®) ©2024 American Medical Association: Chicago, IL.**

General Background

Gastric electrical stimulation (GES) delivers electrical stimuli to the musculature of the gastric wall by means of electrodes which are connected to a stimulator device. The intent is to restore effective gastric contractions. GES has been proposed for patients with gastroparesis who are refractory to medical treatment. There are two principal types of GES devices that are available: (1) low-frequency/high-energy GES with long pulse stimulation; and (2) high-frequency/low-energy GES with short pulse stimulation (Lal, et al., 2015; Bortolotti, 2011).

Low frequency/high-energy GES with long pulse stimulation, also called gastric pacing, uses frequencies close to or above the normal gastric slow wave cycle to reset regular slow wave rhythm. This type of GES involves heavy batteries, is not suitable for implantation and has a variable effect on the symptoms of gastroparesis (Lal, et al., 2015).

High-frequency/low-energy GES with short pulse stimulation (e.g., Enterra™ Therapy) is a type of gastric neurostimulation or neuromodulation (Lal, et al., 2015; Bortolotti, 2011). The device is implanted in the body and delivers high-frequency electrical stimulation at four times the basal rate (12 cycles per minute [cpm]) to the stomach. It is proposed that use of this device reduces the symptoms of gastroparesis such as nausea and vomiting and fosters improved gastric emptying.

Gastric Electrical Stimulation for Gastroparesis

Gastroparesis is a chronic motility disorder of the stomach characterized by gastric retention in the absence of mechanical obstruction. The main causes of gastroparesis are idiopathic, diabetic, iatrogenic (e.g., medication-induced) and postsurgical. Idiopathic gastroparesis refers to gastroparesis of unknown etiology. Diabetic gastroparesis is believed to be caused by chronic hyperglycemia which damages the vagus nerve. Iatrogenic gastroparesis can be caused by medications that delay gastric emptying, such as narcotics and tricyclic antidepressants. Gastroparesis that develops after surgery is called postsurgical gastroparesis (Camilleri, 2025; Zoll, et al., 2019; Pasricha, et al., 2017; Parkman, 2015).

Symptoms of gastroparesis include early satiety, nausea, vomiting, bloating, and upper abdominal discomfort. Postprandial vomiting (1–3 hours after meals) of undigested food is typical. Abdominal discomfort is of varying degrees and is not usually the predominant symptom. Symptoms may be persistent or present as episodic flares. Due to the symptoms, some patients will experience

weight loss and malnutrition and, in severe cases, dehydration. There is also an overlap of symptoms with functional dyspepsia (Camilleri, 2025; Zoll, et al., 2019; Parkman, 2015).

The treatment of gastroparesis is guided by the goals of correcting fluid, electrolyte, and nutritional deficiencies; identifying and treating the cause of delayed gastric emptying (e.g., diabetes); and suppressing or eliminating symptoms. Primary medical management for gastroparesis includes dietary modification and pharmacologic therapy with prokinetic (metoclopramide and erythromycin) and antiemetic agents. Patients refractory to treatment are difficult to manage. Treatment may involve changing or combining medications; placement of a gastrostomy or jejunostomy tube for enteral feedings; or in severe cases, total parenteral nutrition (TPN) for brief periods (Camilleri, 2025; Zoll, et al., 2019; Parkman, 2015). Some patients, however, remain refractory to gastroparesis treatment.

Although proposed as a treatment for refractory gastroparesis, the exact mechanism of action of GES is not clearly known. The Enterra™ Therapy System (Enterra Medical, Inc., St. Louis Park, MN) is a gastric electrical stimulator. Enterra Medical was formed in 2022 and will begin assuming the responsibility of Enterra Therapy from Medtronic (Medtronic Inc., 2025). According to the manufacturer, the Enterra Therapy system is composed of a neurostimulator or implanted neurostimulator (INS), two implantable intramuscular leads and an external programming system. The intramuscular stomach leads are implanted laparoscopically on the greater curvature of the stomach. The INS is implanted in a subcutaneous pocket typically created on the abdomen and is then connected to the leads. The INS provides the energy source that delivers the electrical pulse to the stomach muscle through the stomach leads. The generator stimulates the stomach muscle at a set of stimulation parameters determined by the physician (U.S. Food and Drug Administration [FDA], 2000b). Enterra therapy stimulates the nerves and smooth muscles of the stomach by delivering mild electrical pulses, thereby reducing nausea and vomiting symptoms associated with gastroparesis (Enterra Medical, 2025).

U.S. Food and Drug Administration (FDA):

The Enterra Therapy System is a Class III medical device regulated under the Humanitarian Device Exemption (HDE) pathway (HDE Number: H990014). This pathway is intended for devices treating conditions affecting fewer than 8,000 individuals annually in the U.S. Unlike PMA or 510(k) submissions, HDE applications are not required to demonstrate effectiveness through scientifically valid clinical investigations, but they must show that the device does not pose an unreasonable risk and that probable benefits outweigh risks compared to available alternatives.

Indication for use:

Devices in this category are indicated for the treatment of chronic, intractable (drug-refractory) nausea and vomiting secondary to gastroparesis of diabetic or idiopathic etiology, typically in adults aged 18–70 years (FDA, 2025). Effectiveness for other etiologies or age groups has not been established.

Device or Product	Identifier	Manufacturer	Decision Date
Enterra Therapy System	H990014	Medtronic, Inc.	3/31/2000
Enterra II Therapy System	H990014/S105	Enterra Medical, Inc.	9/15/2015
Enterra II MR Conditional	H990014/S227	Enterra Medical, Inc.	10/20/2023

*FDA product codes: LNQ

Note: Device or product names are provided for example purposes only. Their inclusion does not indicate endorsement or preference for any specific brand or model. Coverage decisions are not based solely on FDA approval. This list is not intended to reflect all available products or technologies.

Literature Review: The evidence in the published peer-reviewed medical literature examining the safety and effectiveness of permanent GES for the treatment of gastroparesis primarily consists of observational studies and case series and few randomized control trials (RCTs).

Ducrotte et al. (2020) conducted a multicenter, double-blind randomized controlled trial with crossover that studied the efficacy of GES in patients with refractory vomiting, with or without gastroparesis. Included patients (n=172) had chronic vomiting and/or nausea > 12 months that was related to type 1 or 2 diabetes mellitus, related to a surgical procedure (partial gastric resection surgery and/or vagotomy), or was idiopathic. Patients had normal or delayed gastric emptying with symptoms that were refractory to treatment and severe enough to affect the general condition of the patient. Patients didn't have evidence of a mechanical obstruction within the digestive tract or a neurologic disease. Patients were randomized to either the ON/OFF group (n=79) with four months of active stimulation followed by four months of sham stimulation or the OFF/ON group (n=93) with four months of sham stimulation followed by four months with active stimulation. Patients were examined at the end of each four-month period (at five and nine months after implantation). Primary endpoints measured were vomiting score, ranging from 0 (daily vomiting) to 4 (no vomiting), and the quality of life, assessed by the Gastrointestinal Quality of Life Index scoring system. Secondary endpoints were changes in other digestive symptoms, nutritional status, gastric emptying, and control of diabetes. Final analysis in the intention to treat (ITT) population was carried out in 66 patients in the ON/OFF group and in 83 patients in the OFF/ON group. During both phases of the crossover study, vomiting scores were significantly higher in the group with the device on than the control group ($p<0.001$), in diabetic and nondiabetic patients. Vomiting scores increased significantly when the device was ON in patients with delayed ($p<0.01$) or normal gastric emptying ($p=0.05$). Gastric emptying was not accelerated during the ON period compared with the OFF period. Having the GES turned on was not associated with increased quality of life. A total of 101 adverse events were reported in the study, with 45 therapy or device -related events: abdominal wall pain at the implantation site (n=28), infections at the abdominal pouch level (n=16), hematoma (n=1). In three cases, the device-related adverse events were serious enough to prompt device removal. The authors concluded that GES is effective in reducing the frequency of refractory vomiting and nausea in a subset of patients with chronic vomiting. Further studies are needed to determine predictive factors of favorable response.

McCallum et al. (2010) conducted a prospective, multicenter, double-blinded, randomized cross-over study (n=55) to evaluate the safety and efficacy of the Enterra gastric stimulation system in the treatment of intractable (drug-refractory) nausea and vomiting secondary to gastroparesis of diabetic etiology. The primary outcome measure was the reduction in weekly vomiting frequency when the device was turned on, relative to when the device was turned off during the blinded cross-over phase. Post-implantation, all patients had the stimulator turned on for six weeks and then were randomly assigned to groups that had consecutive three-month cross-over periods with the device on or off. After this period, the device was turned on in all patients with un-blinded follow-up for four months. Of the 55 subjects enrolled and implanted, 10 were not randomized. A total of 43 subjects completed the cross-over phase and 39 subjects completed 12-month visit follow up. Device-related adverse events included lead migration or dislodgements (n=3), device migrations (n=2), an implant site hematoma, and one implant site infection. The weekly vomiting frequency at 12 months decreased significantly when compared to baseline, with a median reduction of 67.8% ($p<0.001$). Gastric emptying was significantly improved at 12 months with a median retention at four hours of 20.5% compared with 46.5% at baseline ($p<0.001$). Although there were no statistical differences observed in the cross-over period, weekly vomiting frequency was reported to be somewhat better controlled during the on state than the off state. Study limitations include small sample size and loss to follow-up.

O'Grady et al. (2009) performed a meta-analysis of 13 studies evaluating GES for the treatment of medically refractory gastroparesis. Uncontrolled observational studies (n=12) and one blinded randomized control trial (RCT) (Abell, et al., 2003) were included. The findings reported from this review were that following GES, patients had statistically significant improvements in total symptom severity score (p=0.01), vomiting severity score (p<0.0001), and nausea severity score (p<0.0001). The device removal or reimplantation rate was 8.3%.

Case series, retrospective reviews and cohort studies with patient populations ranging from 9–214 support the findings that GES may significantly improve upper GI symptoms and reduce the need for nutritional support in some patients with refractory diabetic or idiopathic gastroparesis (Gourcerol, et al., 2022; Laine, et al., 2018; Shada, et al., 2018; Klinge, et al., 2017; Heckert, et al., 2016; McCallum, et al., 2011; Maranki, et al., 2008; Anand, et al., 2007; McCallum, et al., 2005).

Temporary GES: Temporary GES (tGES) or percutaneous stimulation has been investigated as a potential method for a less invasive trial prior to permanent GES insertion. With the endoscopic technique, temporary non-surgical leads are placed endoscopically on the gastric mucosa and connected to an external gastric stimulation device (Enterra; Enterra Medical, Inc). In temporary percutaneous GES (TPGES), two percutaneous unipolar leads are inserted through a plastic cannula and anchored by flexible wing-like tines to the submucosal tissue (Hasler, 2025; Abel, et al., 2019a; Abel, et al., 2019b; Atassi and Abel 2019; Abell, et al., 2015; Singh, et al., 2015). According to the manufacturer (Enterra Medical, Inc., St. Louis Park, MN) the Enterra II Neurostimulator is implanted beneath the skin in the lower abdominal region. The neurostimulator generates electrical pulses delivered by implanted leads in the antrum portion of the stomach muscle wall. The only lead that is listed as compatible to the device is the intramuscular lead which is designed for intramuscular implantation to deliver electrical current to the stomach muscle (Enterra Medical, 2025).

U.S. Food and Drug Administration (FDA): FDA approval for temporary gastric electrical stimulation was not found on the FDA site. A temporary GES can be carried out using temporary external leads that are primarily designed for external cardiac pacing. However, this represents an off-label use as temporary external leads are not FDA approved for this indication.

Literature Review: There is a paucity of studies in the published peer-reviewed medical literature evaluating temporary GES for gastroparesis or any other indication. Singh et al. (2015) published the results of a cohort study (n=551) which aimed to clarify the role of GES in gastroparesis-like syndrome (GLS), defined as gastroparesis-like symptoms with normal gastric scintigraphy. Inclusion criteria were as follows:

- gastroparesis symptoms of diabetic, surgically related or idiopathic etiology
- aged 18–70 years old
- symptoms of gastroparesis for $\geq$ one year
- refractory or intolerant to prokinetic and antiemetic drug classes
- chronic vomiting or nausea or severe dyspepsia like syndrome consistent with gastroparesis irrespective of gastric emptying test (GET) values

Patients were excluded if they were not candidates for endoscopic or surgical procedures or were pregnant. A total of 452 patients underwent gastric scintigraphy and were stratified into: delayed gastric emptying (n=273), normal gastric emptying (n=137), and rapid gastric emptying categories (n=42). Of the 551 patients in the larger cohort, 379 had tGES implantation using a temporary cardiac pacing lead (Medtronic model 6416). Outcomes measured were changes in gastric scintigraphy and total symptom score. Both components (lead and generator) were used off-label in this study. After tGES, two-hour gastric retention decreased for the delayed patients

($p < 0.01$) and increased for normal and rapid patients ($p < 0.001$). These changes were accompanied by improvements ($p < 0.001$) in vomiting, nausea, and total symptom scores in all three subgroups. Study limitations include the uncontrolled study design and the possibility of the treatment benefit being due to a placebo effect. Although study results suggest that tGES may be effective for treating GLS, well-designed RCTs are needed to support these findings.

Abell et al. (2011) published the results of a randomized, placebo-controlled, crossover trial ($n=58$) to measure the effects of endoscopically placed temporary GES (tGES) on gastroparesis symptoms. The study consisted of two consecutive, 4-day sessions (session 1 and session 2). Inclusion criteria were as follows:

- patients between the ages of 18 and 70 years, with a $\geq$ one-year history of gastroparesis symptoms from diabetic ($n=13$), postsurgical ($n=7$), or idiopathic ($n=38$) etiology
- gastroparesis symptoms refractory or intolerant to antiemetic drug classes with $\geq$ seven episodes of chronic vomiting and/or nausea per week, irrespective of gastric emptying time values

Patients with an active infection or pregnancy were excluded. Temporary GES using a temporary cardiac pacing lead (model 6414-200; Medtronic, Minneapolis, Minn) was provided to 37/58 enrolled patients (group A [$n=21$]; group B [$n=16$]). During session 1 treatment was activated for 72 continuous hours in group A, and likewise activated in group B during session 2. The primary outcome measure was a 50% improvement in baseline symptom values. Secondary outcomes were gastric emptying, electrogastrography, and quality of life measured at baseline and session close. An overall treatment effect of a slight, non-significant daily decrease in average vomiting scores ($p=0.116$) was observed by pooling stimulation effects across sessions. The single reported adverse event was dislodged electrodes for six patients in group A and seven in group B. Study limitations include the small sample size and the fact that patients were allowed to continue medication for nausea or pain (prokinetics, anti-emetics) during the trial. The small sample size and non-significant improvement in symptoms make it difficult to draw conclusions from this study.

Gastric Electrical Stimulation for Other Indications

The use of GES is currently under investigation for the treatment of obesity and type 2 diabetes mellitus (T2DM).

Obesity: GES has been proposed as a device therapy for the treatment of morbid obesity. GES for obesity is currently registered by the FDA as investigational. In Europe, however, GES is being used clinically to treat obesity. Transneuronix, Inc., (Mt. Arlington, NJ), acquired in 2005 by Medtronic Inc. (Minneapolis, MN), developed the Transcend™ Gastric Stimulation System for obesity. This implantable gastric stimulator (IGS) has not been approved by the FDA. The device includes a pulse generator, an external programmer and a gastric stimulation lead, and is implanted laparoscopically in the subcutaneous tissue. The Transcend is intended to induce satiety by delaying gastric emptying (Greenway and Zheng, 2007).

A number of unresolved issues regarding the use of GES for treatment of obesity have been identified. The mechanism of action is unclear. Proposed possibilities include: a local enteric nervous system effect, an effect mediated by the autonomic nervous system, possible central nervous system changes and neurohormonal changes. Optimal stimulation patterns are unknown, as is the importance of the number of leads and the location of electrodes. Optimal screening of patients for GES for obesity has not yet been determined. Also, the best combination of behavioral, drug, device and surgical therapies has not been determined (Abell, et al., 2006). As a result, the use of a gastric pacing device for these indications remains under investigation.

Literature Review: GES for the treatment of obesity has been evaluated in case series, randomized controlled trials (RCTs) and systematic reviews. Paulus et al. (2020) conducted a multicenter, phase 1, open prospective cohort study in the Netherlands and the USA. The study assessed the following in patients with morbid obesity: the safety of the Exilis™ gastric electrical stimulation (GES) system, the setting adjustments for chronic use and the acute gastrointestinal (GI)/feeding effects. Patients were included in this study if they were weight stable, aged 21–64 years and had a body mass index (BMI) of 40–45 kg/m² or 35–39.9 kg/m² with at least one weight-related comorbidity (e.g., nonalcoholic steatohepatitis, hypertension, dyslipidemia, obstructive sleep apnea, arthrosis). If a patient was diagnosed with diabetes mellitus, the diagnosis had to be made within the last seven years, had to be currently treated with oral agents only and had to have an HbA1c ≤ 8%. Twenty morbidly obese patients (17 female, mean BMI of 40.8 ± 0.7 kg/m²) were implanted with the Exilis™ system followed by a two-week recovery period prior to continuation of the study protocol. The study protocol included four amplitude titration visits (visits A, B, C, and D) occurring at weekly intervals. The amplitude titration visits were followed by two GI function test days performed in randomly assigned order and repeated twice (once with GES ON and once with GES OFF). Each GI function test day was preceded by a washout period (GES OFF) of seven days, and subjects were blinded to the assigned GES treatment. Testing at weeks 26 and 52 included simultaneous measurement of gastric emptying (using a breath test), gastric motility (SmartPill®), plasma concentrations of glucose and insulin, and food intake over a four-hour period in the morning following an overnight fast. Impact of Weight on Quality of Life-Lite (IWQOL-Lite) and the Multi-purpose Short Form Survey-12 (SF-12) were used to measure quality of life. Both surveys were administered at screening visit, week 0, 13, 26, and 52 postoperatively.

The authors stated that there were not any serious adverse events in all 20 subjects, with the exception of incisional hernias which had to be corrected surgically (n=2). The other adverse events were mild and were related to the IPG pocket (seroma, infection, hernia) and are most likely due to the relatively superficial placement of the device. At the 26-week and 52-week follow-up, three and four subjects (respectively) had withdrawn from the study due to not reaching the desired effect. Most of the patients that withdrew from the study had a surgical revision to Roux-en-Y gastric bypass (RYGB) or laparoscopic sleeve gastrectomy (LSG). They were not included in further analysis. At week four, 13, and 26, a significant reduction in weight loss was observed (p<0.01) but not at week 52. At this time point, the mean excess weight loss (EWL) was 14.2 ± 4.5%. There were no significant differences between GES ON and OFF in gastric emptying halftime, food intake, insulin, and glucose (all p>0.05). Author noted limitations included possible adaptation to the signal which could have resulted in loss of efficacy and the lack of a control group. Additional limitations included the small patient population and only patients from the Netherlands and the USA were enrolled and the results may not be applicable to other races or ethnic groups. The study concluded that gastric electrical stimulation with the Exilis™ system can be considered as safe. However, no significant effect on food intake, gastric emptying, or gastric motility was observed. The reduction in weight loss with Exilis™ wasn't observed in the long term. Further electrophysiological research is needed to gain more insight in optimal stimulation parameters and lead localization. No health disparities were identified by the investigators.

Cha et al. (2014) performed a systematic review (n=31 studies/1367patients) of the evidence to evaluate the effect of different types of gastric electrical stimulation (GES) on obesity. Published studies investigating the effect of GES using the Tantalus and Transcend devices, as well as vagus nerve stimulation, were included. Exclusion criteria for published studies were GES used for diseases other than obesity (e.g., gastroparesis), non-gastric stimulation, and non-clinical primary outcome. Studies were primarily non-randomized, with 4/31 randomized trials. In all studies, the generator was externalized and in most cases, they were implanted in subcutaneous layers of the anterior abdominal wall. The electrodes connected to the generator were implanted in different

locations of the stomach, depending on the type of GES. The primary outcome was weight loss, with secondary outcomes of appetite or satiety changes and biochemical marker changes. Almost all studies in each device group achieved statistically significant weight loss during the first 12 months. Only a small percentage of studies had a follow-up longer than one year and found significant weight loss maintenance. Findings were inconsistent for secondary outcomes. Gastric penetration was the most common device-related complication. In general, the level of evidence was found to be low with few studies having a large population and low loss to follow-up. Results of studies in this systematic review suggest that GES may be effective for short-term weight loss. However well-designed studies with larger patient population and long-term follow up are needed to determine safety and effectiveness of the technology for this indication.

Shikora et al. (2009) conducted a randomized, double-blinded, placebo-controlled study (n=190), the Screened Health Assessment and Pacer Evaluation (SHAPE) trial. The SHAPE study compared gastric stimulation therapy (n=96) to a standard diet and behavioral therapy regimen (n=94) in a group of obese patients. Subjects were required to be 18–65 years of age and have a BMI of 35–55 kg/m². Exclusion criteria included pregnancy, previous gastrointestinal bariatric surgery, the presence of other electrostimulation devices (e.g., pacemakers), gastrointestinal motility disorders, peptic ulcer disease, and clinically significant comorbidities such as poorly controlled diabetes. Follow-up occurred monthly for 12 months. The difference in excess weight loss (% EWL) between the control group and the treatment group was not found to be statistically significant (p=0.717) at 12 months of follow-up. These results suggest that this technology is not effective in achieving significant weight loss in severely obese individuals.

Shikora (2004) reported an update of two U.S. clinical trials for gastric stimulation in obesity. The first was an RCT in 2000 that included patients (n=103) age 18–50 who had a BMI of 40–55 kg/m² (mean 46 kg/m²). No statistical difference in the weight loss between study and control groups was found at six-month follow-up. At 29 months, the overall mean EWL increased to > 12.0%. A total of 69 patients were lost to follow-up.

The second trial (n=30), the Dual-Lead Implantable Gastric Electrical Stimulation Trial (DIGEST), was a non-randomized, open-label study of patients with a BMI 40–55 kg/m² or 35–39 kg/m² and one or more significant comorbidities. At the 12-month follow-up point, 71% of participants lost weight (54% lost > 10% of excess, and 29% lost > 20% excess). At the 16-month follow-up, mean EWL was 23%.

Several case series (n=11–91 patients) have investigated the implantation of GES for the treatment of obesity reporting varying rates of excess weight loss and improvement of comorbidities (Bohdjalian, et al., 2009; Miller, et al., 2006; Cigaina, et al., 2003). In addition to the lack of randomization, in general studies have been limited by small sample sizes and short-term follow-up.

There is insufficient evidence in the published scientific literature to support the use of gastric pacing for the treatment of morbid obesity.

Type 2 Diabetes Mellitus (T2DM): The effect of GES on HbA1c and blood glucose levels, along with changes in body weight is also being investigated. The DIAMOND® (Diabetes Improvement and MetabOlic Normalization Device), formerly known as the TANTALUS device, has been developed by MetaCure, Inc. (Kfar-Saba, Israel). The DIAMOND device consists of three pairs of bipolar electrodes. One pair is attached to the gastric fundus and the other two pairs are attached to the anterior and the posterior antrum of the stomach. The electrodes are implanted laparoscopically and connected to a pulse generator inserted into the subcutaneous tissue of the abdomen. The pulse generator uses a rechargeable battery as an external power source. The delivered electrical signal characteristics are set by a programmer within the first week after the

implantation (Lebovitz, et al., 2015). Clinical trials are now being conducted using this device for overweight and obese patients with type 2 diabetes.

Literature Review: The evidence in the published peer-reviewed medical literature examining the safety and effectiveness of GES for obese patients with T2DM consists of few case series (Lebovitz, et al., 2015; Bohdjalian, et al., 2009; Policker, et al., 2009; Sanmiguel, et al., 2009). Patient populations in these studies have ranged from 14–61, with a follow-up of primarily six–12 months. Although preliminary results suggest that GES may improve glycemic control and induce weight loss in patients with T2DM, additional evidence in the form of well-designed RCTs is needed to confirm these findings.

Professional Societies/Organizations

American College of Gastroenterology (ACG): The ACG published guidelines stated that gastric electrical stimulation may be considered for humanitarian use in patients with refractory diabetic gastroparesis or idiopathic gastroparesis (Camilleri, et al., 2022).

American Diabetes Association (ADA): In the ADA Standards of medical care in diabetes – 2025, state that although gastric electrical stimulation using a surgically implantable device has received approval from the FDA, data is very limited, and the results do not support gastric stimulation as an effective therapy in diabetic gastroparesis.

American Gastroenterological Association (AGA): In the 2025 Clinical Practice Guideline on the Management of Gastroparesis published by the AGA (Staller et al., 2025), the guideline panel issued a conditional recommendation against the routine initial use of gastric electrical stimulation (GES) in patients with gastroparesis. This intervention is reserved for select individuals whose symptoms remain refractory despite optimized medical therapy, emphasizing the importance of individualized care and shared decision-making.

National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK): The NIDDK stated that gastric electrical stimulation may be effective for some people whose nausea and vomiting do not improve with dietary changes or medications (NIDDK, 2018).

Health Equity Considerations

Health equity is the highest level of health for all people; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which people are born, grow, live, work, and age.

Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include safe housing, transportation, and neighborhoods; racism, discrimination and violence; education, job opportunities and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

The retrospective study by Friedenberg et al. (2013) studied the influence of race on symptom severity and quality of life in gastroparesis. The study included 44 (17%) nonwhites (33 African American and 11 Hispanic) and 211 (83%) whites. The study reported that nonwhite and white patients with gastroparesis differ in disease etiology and health care utilization. Nonwhite patients with gastroparesis secondary to diabetes was 55% compared with 19% of white patients ($p < 0.001$). Additionally, 49% of nonwhite patients reported ≥ 4 gastroparesis-related emergency department visits and 42% reported more ≥ 4 gastroparesis related hospitalizations, as compared with 20% and 14% of white patients, respectively. The study concluded that nonwhite patients

with gastroparesis were more likely to have diabetes as the etiology, have more severe symptoms, poorer QOL and utilized more health care resources than white patients.

Medicare Coverage Determinations

	Contractor	Determination Name/Number	Revision Effective Date
NCD		No National Coverage Determination found	
LCD		No Local Coverage Determination found	

Note: Please review the current Medicare Policy for the most up-to-date information.
(NCD = National Coverage Determination; LCD = Local Coverage Determination)

References

1. Abell TL, Bernstein RK, Cutts T, Farrugia G, Forster J, Hasler WL, et al. Treatment of gastroparesis: a multidisciplinary clinical review. *Neurogastroenterol Motil*. 2006 Apr;18(4):263-83.
2. Abell TL, Chen J, Emmanuel A, Jolley C, Sarela AI, Törnblom H. Neurostimulation of the gastrointestinal tract: review of recent developments. *Neuromodulation*. 2015 Apr;18(3):221-7; discussion 227.
3. Abell TL, Johnson WD, Kedar A, Runnels JM, Thompson J, Weeks ES, et al. A double-masked, randomized, placebo-controlled trial of temporary endoscopic mucosal gastric electrical stimulation for gastroparesis. *Gastrointest Endosc*. 2011 Sep;74(3):496-503.e3.
4. Abell TL, Kedar A, Stocker A, Beatty K, McElmurray L, Hughes M, et al. Gastroparesis syndromes: Response to electrical stimulation. *Neurogastroenterol Motil*. 2019b Mar;31(3):e13534
5. Abell T, McCallum R, Hocking M, Koch K, Abrahamsson H, LeBlanc I, et al. Gastric electrical stimulation for medically refractory gastroparesis. *Gastroenterol*. 2003 Aug;125(2).
6. Abell TL, Yamada G, McCallum RW, Van Natta ML, Tonascia J, Parkman HP, Koch KL, Sarosiek I, et al. Effectiveness of gastric electrical stimulation in gastroparesis: Results from a large prospectively collected database of national gastroparesis registries. *Neurogastroenterol Motil*. 2019a Dec;31(12):e13714.
7. American Diabetes Association. Standards of medical care in diabetes - 2025 12. Retinopathy, Neuropathy, and Foot Care: Standards of Care in Diabetes—2025. *Diabetes Care* 2025;48 (Supplement_1):S252-S265.
8. Anand C, Al-Juburi A, Familoni B, Rashed H, Cutts T, Abidi N, et al. Gastric electrical stimulation is safe and effective: a long-term study in patients with drug-refractory gastroparesis in three regional centers. *Digestion*. 2007;75(2-3):83-9.
9. Atassi H, Abell TL. Gastric Electrical Stimulator for Treatment of Gastroparesis. *Gastrointest Endosc Clin N Am*. 2019 Jan;29(1):71-83.
10. Bardaro SJ, Guerron AD, Romanelli J, Soriano I, King K, Gibbs KE, Petrick A, Lo Menzo E, Rosenthal R, Kennedy C, Gershuni V, Daly S, Leyva-Alvizo A, Tran M, Stalin V, Kothari S, Sudan R. Gastroparesis: an evidence-based review for the bariatric and foregut surgeon.

Surg Obes Relat Dis. 2023 May;19(5):403-420. doi: 10.1016/j.soard.2023.02.018. Epub 2023 Mar 4. PMID: 37080885.

11. Bohdjalian A, Prager G, Rosak C, Weiner R, Jung R, Schramm M, et al. Improvement in glycemic control in morbidly obese type 2 diabetic subjects by gastric stimulation. *Obes Surg*. 2009 Sep;19(9):1221-7.
12. Bortolotti M. Gastric electrical stimulation for gastroparesis: A goal greatly pursued, but not yet attained. *World J Gastroenterol*. 2011 Jan 21; 17(3): 273–282.
13. Camilleri M. Gastroparesis: Etiology, clinical manifestations, and diagnosis. In: UpToDate, Talley NJ (Ed). Dec 16, 2024. UpToDate, Waltham, MA. Accessed on Oct 2, 2025.
14. Camilleri M. Treatment of gastroparesis. In: UpToDate, Talley NJ (Ed). Jun 23, 2025. UpToDate, Waltham, MA. Accessed on Oct 2, 2025.
15. Camilleri M, Kuo B, Nguyen L, Vaughn VM, Petrey J, Greer K, et al. ACG Clinical Guideline: Gastroparesis. *Am J Gastroenterol*. 2022 Aug 1;117(8):1197-1220.
16. Centers for Medicare and Medicaid Services (CMS). Medicare Coverage Database. Accessed Oct 2, 2025. Available at URL address: <https://www.cms.gov/medicare-coverage-database/search.aspx>
17. Cha R, Marescaux J, Diana M. Updates on gastric electrical stimulation to treat obesity: Systematic review and future perspectives. *World J Gastrointest Endosc*. 2014 Sep 16;6(9):419-31.
18. Chu H, Lin Z, Zhong L, et al. Treatment of high-frequency gastric electrical stimulation for gastroparesis. *J Gastroenterol Hepatol*. 2012 Jun;27(6):1017-26.
19. Cigaina V, Hirschbert AL. Gastric pacing for morbid obesity: plasma levels of gastrointestinal peptides and leptin. *Obes Res*. 2003;11(12):1456-62.
20. Corvinus FM, Heinrich S, Neumann H, Hadzijasufovic E, Babic B, Lang H, et al. Minimally-invasive temporary gastric stimulation: A pilot study to predict the outcome of electronic gastric stimulation with the Enterra™ system. *Dig Liver Dis*. 2018 Oct;50(10):1030-1034.
21. Ducrotte P, Coffin B, Bonaz B, Fontaine S, Bruley Des Varannes S, Zerbib F, et al. Gastric Electrical Stimulation Reduces Refractory Vomiting in a Randomized Crossover Trial. *Gastroenterology*. 2020 Feb;158(3):506-514.e2.
22. Enterra Medical. About Enterra™ Therapy. © 2025 Enterra Medical, Inc. Accessed Oct 2, 2025. Available at URL address: <https://www.enterramedical.com/hcp/about-enterra-therapy/>
23. Friedenberg FK, Kowalczyk M, Parkman HP. The influence of race on symptom severity and quality of life in gastroparesis. *J Clin Gastroenterol*. 2013 Oct;47(9):757-61.
24. Gourcerol G, Coffin B, Bonaz B, Hanaire H, Bruley Des Varannes S, Zerbib F, et al. Impact of Gastric Electrical Stimulation on Economic Burden of Refractory Vomiting: A French Nationwide Multicentre Study. *Clin Gastroenterol Hepatol*. 2022 Aug;20(8):1857-1866.e1.

25. Greenway F, Zheng J. Electrical stimulation as treatment for obesity and diabetes. *J Diabetes Sci Technol*. 2007 Mar;1(2):251-9.
26. Grover I, Kim R, Spree DC, Lahr CJ, Kedar A, Kothari S, et al. Gastric Electrical Stimulation is an Option for Patients with Refractory Cyclic Vomiting Syndrome. *J Neurogastroenterol Motil*. 2016 Oct 30;22(4):643-649.
27. Hasanin M, Amin O, Hassan H, Kedar A, Griswold M, Abell TL. Temporary Gastric Stimulation in Patients With Gastroparesis Symptoms: Low-Resolution Mapping Multiple Versus Single Mucosal Lead Electrograms. *Gastroenterology Res*. 2019 Apr;12(2):60-66.
28. Hasler WL. Electrical stimulation for gastroparesis. In: UpToDate, Talley NJ (Ed). Aug 8, 2025. UpToDate, Waltham, MA. Accessed on Oct 2, 2025.
29. Heckert J, Sankineni A, Hughes WB, Harbison S, Parkman H. Gastric electrical stimulation for refractory gastroparesis: a prospective analysis of 151 patients at a single center. *Dig Dis Sci* 2016 Jan;61(1):168-175.
30. Horbach T, Thalheimer A, Seyfried F, Eschenbacher F, Schuhmann P, Meyer G. abiliti Closed-Loop Gastric Electrical Stimulation System for Treatment of Obesity: Clinical Results with a 27-Month Follow-Up. *Obes Surg*. 2015 Oct;25(10):1779-87.
31. Jayanthi NV, Dexter SP, Sarela AI; Leeds Gastroparesis Multi-Disciplinary Team. Gastric electrical stimulation for treatment of clinically severe gastroparesis. *J Minim Access Surg*. 2013 Oct;9(4):163-7
32. Krishnasamy S, Abell TL. Diabetic gastroparesis: principles and current trends in management. *Diabetes Ther*. 2018;9:S1-S42.
33. Klinge MW, Rask P, Mortensen LS, Lassen K, Ejksjaer N, Ehlers LH, et al. Early Assessment of Cost-effectiveness of Gastric Electrical Stimulation for Diabetic Nausea and Vomiting. *J Neurogastroenterol Motil*. 2017 Oct 30;23(4):541-549.
34. Lacy BE, Tack J, Gyawali CP. AGA Clinical Practice Update on Management of Medically Refractory Gastroparesis: Expert Review. *Clin Gastroenterol Hepatol*. 2022 Mar;20(3):491-500.
35. Laine M, Sirén J, Koskenpato J, Punkkinen J, Rantanen T, Typpö I, et al. Outcomes of high-frequency gastric electric stimulation for the treatment of severe, medically refractory gastroparesis in Finland. *Scand J Surg*. 2018 Jun;107(2):124-129.
36. Lal N, Livemore S, Dunne D, Khan I. Gastric Electrical Stimulation with the Enterra System: A Systematic Review. *Gastroenterol Res Pract*. 2015; 2015: 762972.
37. Lebovitz HE, Ludvik B, Yaniv I, Schwartz T, Zelewski M, Guttermann DD; Metacure Investigators. Treatment of Patients with Obese Type 2 Diabetes with Tantalus-DIAMOND® Gastric Electrical Stimulation: Normal Triglycerides Predict Durable Effects for at Least 3 Years. *Horm Metab Res*. 2015 Jun;47(6):456-62.
38. Levinthal DJ, et al. Systematic review and meta-analysis: Gastric electrical stimulation for gastroparesis. *Auton Neurosci*. Jan 2017;202:45-55

39. Maranki JL, Lytes V, Meilahn JE, Harbison S, FriedenberG FK, Fisher RS, et al. Predictive Factors for Clinical Improvement with Enterra Gastric Electric Stimulation Treatment for Refractory Gastroparesis. *Dig Dis Sci*. 2008 Aug;53(8):2072-8.
40. McCallum R, Lin Z, Wetzel P, Sarosiek I, Forster J. Clinical response to gastric electrical stimulation in patients with postsurgical gastroparesis. *Clin Gastroenterol Hepatology*. 2005;3:49-54.
41. McCallum RW, Snape W, Brody F, Wo J, Parkman HP, Nowak T. Gastric electrical stimulation with Enterra therapy improves symptoms from diabetic gastroparesis in a prospective study. *Clin Gastroenterol Hepatol*. 2010 Nov;8(11):947-54.
42. McCallum RW, Lin Z, Forster J, Roeser K, Hou Q, Sarosiek I. Gastric electrical stimulation improves outcomes of patients with gastroparesis for up to 10 years. *Clin Gastroenterol Hepatol*. 2011 Apr;9(4):314-319.e1.
43. Medtronic Inc. About Enterra Therapy. 2025. Accessed Oct 2, 2025. Available at URL address: <https://www.medtronic.com/us-en/patients/treatments-therapies/neurostimulator-gastroparesis.html>
44. Miller K, Hoeller E, Aigner F. The Implantable Gastric Stimulator for Obesity : An Update of the European Experience in the LOSS (Laparoscopic Obesity Stimulation Survey) Study. *Treat Endocrinol*. 2006;5(1):53-8.
45. Morales-Conde S, Alarcon Del Agua I, Busetto L, et al. Implanted closed-loop gastric electrical stimulation (CLGES) system with sensor-based feedback safely limits weight regain at 24 months. *Obes Surg*. 2018;28(6):1766-1774.
46. National Institutes of Health. National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Treatment for Gastroparesis. Jan 2018. Accessed Oct 2, 2025. Available at URL address: <https://www.niddk.nih.gov/health-information/digestive-diseases/gastroparesis/treatment>
47. O'Grady G, Egbuji JU, Du P, Cheng LK, Pullan AJ, Windsor JA. High-frequency gastric electrical stimulation for the treatment of gastroparesis: a meta-analysis. *World J Surg*. 2009 Aug;33(8):1693-701.
48. Parkman HP. Idiopathic gastroparesis. *Gastroenterol Clin North Am*. 2015 Mar;44(1):59-68.
49. Parkman HP, Yamada G, Van Natta ML, Yates K, Hasler WL, Sarosiek I, et al. Ethnic, Racial, and Sex Differences in Etiology, Symptoms, Treatment, and Symptom Outcomes of Patients With Gastroparesis. *Clin Gastroenterol Hepatol*. 2019 Jul;17(8):1489-1499.e8.
50. Pasricha PJ, Camilleri M, Hasler WL, Parkman HP. White Paper AGA: Gastroparesis: Clinical and Regulatory Insights for Clinical Trials. *Clin Gastroenterol Hepatol*. 2017 Aug;15(8):1184-1190.
51. Paulus GF, van Avesaat M, van Rijn S, Alleleyn AME, Swain JM, Abell TL. Multicenter, Phase 1, Open Prospective Trial of Gastric Electrical Stimulation for the Treatment of Obesity: First-in-Human Results with a Novel Implantable System. *Obes Surg*. 2020 May;30(5):1952-1960.

52. Policker S, Haddad W, Yaniv I. Treatment of type 2 diabetes using meal-triggered gastric electrical stimulation. *Isr Med Assoc J*. 2009 Apr;11(4):206-8.
53. Sanmiguel CP, Conklin JL, Cunneen SA, Barnett P, Phillips EH, Kipnes M, et al. Gastric electrical stimulation with the TANTALUS System in obese type 2 diabetes patients: effect on weight and glycemic control. *J Diabetes Sci Technol*. 2009 Jul 1;3(4):964-70.
54. Shada A, Nielsen A, Marowski S, Helm M, Funk LM, Kastenmeier A, et al. Wisconsin's Enterra therapy experience: a multi-institutional review of gastric electrical stimulation for medically refractory gastroparesis. *Surgery* 2018 Oct;164(4):760-765
55. Shikora SA. "What are the yanks doing?" The U.S. experience with implantable gastric stimulation (IGS) for the treatment of obesity-update on the ongoing clinical trials. *Obes Surg*. 2004;14:S40-8.
56. Shikora SA, Bergenstal R, Bessler M, Brody F, Foster G, Frank A, et al. Implantable gastric stimulation for the treatment of clinically severe obesity: results of the SHAPE trial. *Surg Obes Relat Dis*. 2009 Jan-Feb;5(1):31-7.
57. Singh S, McCrary J, Kedar A, Weeks S, Beauerle B, Weeks A, et al. Temporary Endoscopic Stimulation in Gastroparesis-like Syndrome. *J Neurogastroenterol Motil*. 2015 Oct 1;21(4):520-7.
58. Staller K, Parkman HP, Greer KB, Leiman DA, Zhou MJ, Singh S, Camilleri M, Altayar O; AGA Clinical Guidelines Committee. AGA Clinical Practice Guideline on Management of Gastroparesis. *Gastroenterology*. 2025 Oct;169(5):828-861. doi: 10.1053/j.gastro.2025.08.004. PMID: 40976635.
59. U.S. Food Drug Administration (FDA). Humanitarian Device Exemption (HDE) database. Product code(s): LNQ. Page Last Updated: Sep 29, 2025. Accessed Oct 2, 2025. Available at URL address: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfHDE/hde.cfm>
60. Zoll B, Jehangir A, Malik Z, Edwards MA, Petrov RV, Parkman HP. Gastric Electric Stimulation for Refractory Gastroparesis. *J Clin Outcomes Manag*. 2019 Jan;26(1):27-38.

Revision Details

Type of Revision	Summary of Changes	Date
Annual review	No clinical policy statement changes.	11/15/2025
Annual review	No clinical policy statement changes.	12/15/2024
Annual review	Revised the gastric electrical stimulation (GES) or gastric pacing for any other indication policy statement.	10/15/2023

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.