



Drug Coverage Policy

Effective Date 11/15/2025

Coverage Policy Number IP0075

Policy Title Armodafinil, Modafinil

Wakefulness-Promoting Agents – Armodafinil, Modafinil

- Nuvigil® (armodafinil tablets – Cephalon, generic)
- Provigil® (modafinil tablets – Cephalon, generic)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

OVERVIEW

Armodafinil and modafinil, agents with wake-promoting actions that are similar to sympathomimetic agents (e.g., amphetamine and methylphenidate), are indicated to improve wakefulness in adults with **excessive sleepiness** associated with the following conditions:^{1,2}

- **Narcolepsy.**
- **Obstructive sleep apnea/hypopnea syndrome** (approved as adjunctive therapy).
- **Shift work sleep disorder.**

Armodafinil and modafinil are Schedule IV controlled substances.^{1,2} Review of the medical literature notes many other uses of modafinil that are considered off-label or investigational. While armodafinil has not been studied off-label to the same extent as modafinil, it is expected that armodafinil will have similar clinical efficacy for these uses.

Two specialized tests, which can be performed in a sleep disorders clinic, are required to establish a diagnosis of narcolepsy.³ Polysomnogram (PSG) is an overnight recording of brain and muscle activity, breathing, and eye movements. The multiple sleep latency test assesses daytime sleepiness by measuring how quickly a person falls asleep and whether they enter rapid eye movement (REM) sleep. On the day after PSG, the patient is asked to take five short naps separated by 2 hours over the course of a day. If an individual falls asleep in < 8 minutes on average over the five naps, this indicates excessive daytime sleepiness. However, patients with narcolepsy also have an abnormally quick start to REM sleep. If REM sleep happens within 15 minutes at least two times out of the five naps and the sleep study the night before, this is likely an abnormality caused by narcolepsy.

Guidelines

Pertinent medical guidelines related to modafinil and armodafinil are summarized below.

Narcolepsy and Cataplexy

The American Academy of Sleep Medicine (AASM) practice parameters for the treatment of central disorders of hypersomnolence were updated in 2021.^{4,5}

- Modafinil, Wakix® (pitolisant tablets), Xyrem® (sodium oxybate oral solution), and Sunosi® (solriamfetol tablets) are recommended as effective treatments for daytime sleepiness due to narcolepsy and reducing disease severity in adults (Strong Recommendation for each).
- Wakix and Xyrem have also demonstrated efficacy for the treatment of cataplexy in patients with narcolepsy (Strong Recommendation for each).
- Xyrem and armodafinil have Conditional Recommendations for the treatment of narcolepsy, showing efficacy for daytime sleepiness due to narcolepsy and reducing disease severity.
- Dextroamphetamine has a Conditional Recommendation for the treatment of narcolepsy, showing efficacy for excessive daytime sleepiness and cataplexy.
- Methylphenidate has a Conditional Recommendation for the treatment of narcolepsy, showing efficacy in reducing disease severity.
- There was insufficient and inconclusive evidence to make recommendations for l-carnitine, scheduled naps, selegiline, triazolam, selective serotonin reuptake inhibitors (SSRIs), and serotonin-norepinephrine reuptake inhibitors (SNRIs).
- Modafinil and Xyrem have Conditional Recommendations for the treatment of narcolepsy in pediatric patients.
- A Strong Recommendation should be followed by clinicians under most circumstances. A Conditional Recommendation requires that the clinician use clinical knowledge and experience and strongly consider the individual patient's values and preferences to determine the best course of action.

Excessive Daytime Sleepiness Associated with Obstructive Sleep Apnea/Hypoapnea Syndrome

- According to the AASM guideline on treatment of adult obstructive sleep apnea (OSA) with positive airway pressure (PAP) [2019], PAP, compared with no therapy, is recommended to treat OSA in adults with excessive sleepiness (Strong Recommendation).⁶
- PAP therapy is recommended to be initiated using either auto-titrating positive airway pressure (APAP) at home or in-laboratory PAP titration in adults with OSA and no significant comorbidities (Strong Recommendation).
- Either continuous positive airway pressure (CPAP) or APAP is recommended for ongoing treatment of OSA in adults (Strong Recommendation).
- Use of CPAP or APAP over bilevel positive airway pressure (BPAP) is suggested in the routine treatment of OSA in adults (Conditional Recommendation).

Adjunctive/Augmentation Treatment for Major Depressive Disorder

- According to the American Psychiatric Association (APA) practice guideline for the treatment of patients with major depressive disorder (2010), modafinil (or methylphenidate) are potential treatments for sedation associated with antidepressant medications.⁷
- The APA guidelines state that modafinil has shown benefit when combined with an SSRI, related to specific effects on residual symptoms such as fatigue and hypersomnolence.
- The guidelines note that there is no clear guidance regarding the length of time modafinil should be co-administered.
- While armodafinil has not been studied for this use, it is considered to be interchangeable with modafinil for this condition.

Excessive Daytime Sleepiness Associated with Myotonic Dystrophy

- Practice parameters from the AASM, last updated in 2021, suggest that clinicians use modafinil for the treatment of hypersomnia secondary to myotonic dystrophy in adults (Conditional Recommendation).^{4,5}
- While armodafinil has not been studied for this use, it is considered to be interchangeable with modafinil for this condition.

Excessive Daytime Sleepiness Associated with Parkinson's Disease

- Practice parameters from the AASM (2021) suggest that clinicians use modafinil for the treatment of hypersomnia secondary to Parkinson's disease in adults (Conditional Recommendation).^{4,5}
- While armodafinil has not been studied for this use, it is considered to be interchangeable with modafinil for this condition.

Fatigue Associated with Multiple Sclerosis

- Practice parameters from the AASM (2021) suggest that clinicians use modafinil for the treatment of hypersomnia secondary to multiple sclerosis in adults (Conditional Recommendation).^{4,5}
- While armodafinil has not been studied for this use, it is considered to be interchangeable with modafinil for this condition.

Idiopathic Hypersomnia

Idiopathic hypersomnia, a condition similar to narcolepsy, is characterized by constant or recurrent daytime sleepiness with no other cause of sleepiness, prolonged nocturnal sleep, difficulty awakening with sleep drunkenness, and long unrefreshing naps with no history of cataplexy.⁸⁻¹⁰ The AASM practice parameters for the treatment of central disorders of hypersomnolence (2021) include recommendations for the treatment of idiopathic hypersomnia.^{4,5}

- Only modafinil has a Strong Recommendation for use.
- Clarithromycin, methylphenidate, Wakix, and Xyrem have Conditional Recommendations for the treatment of idiopathic hypersomnia in adults.

Coverage Policy

POLICY STATEMENT

Prior Authorization is required for benefit coverage of modafinil (brand and generic) and armodafinil (brand and generic). This Prior Authorization Policy also contains a Step Therapy component. When clinically appropriate, the patient is directed to try one Step 1 Product (generic modafinil or generic armodafinil) prior to brand Nuvigil or brand Provigil (Step 2). All approvals are provided for the duration noted below.

Documentation: Documentation is required where noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory tests, claims records, and/or other information.

Modafinil (brand and generic) and armodafinil (brand and generic) are considered medically necessary when ONE of the following is met:

FDA-Approved Indications

- 1. Excessive Daytime Sleepiness Associated with Narcolepsy.** Approve for 1 year if the patient meets the following (A, B, C, D, and E):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient has been evaluated using polysomnography and a multiple sleep latency test **[documentation required]**; AND
 - C)** Diagnosis of narcolepsy has been confirmed **[documentation required]**; AND
 - D)** The medication is prescribed by or in consultation with a sleep specialist physician or a neurologist; AND
 - E)** Preferred product criteria is met for the products as listed in the below tables
- 2. Excessive Daytime Sleepiness Associated with Obstructive Sleep Apnea/Hypopnea Syndrome.** Approve for 1 year if the patient meets the following (A, B, and C):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient meets one of the following (i or ii):
 - i. Armodafinil/modafinil will be used in conjunction with positive airway pressure therapy; OR
 - ii. Patient is unable to initiate or tolerate positive airway pressure therapy **[documentation required]**; AND
Note: Positive airway pressure can include continuous positive airway pressure (CPAP), auto-titrating positive airway pressure (APAP), or bilevel positive airway pressure (BPAP).
 - C)** Preferred product criteria is met for the products as listed in the below tables
- 3. Excessive Sleepiness Associated with Shift Work Sleep Disorder.** Approve for 1 year if the patient meets the following (A, B, and C):
 - A)** Patient is ≥ 18 years of age; AND
 - B)** Patient works at least five overnight shifts per month **[documentation required]**; AND
 - C)** Preferred product criteria is met for the products as listed in the below tables

Other Uses with Supportive Evidence

- 4. Adjunctive/Augmentation Treatment for Depression in Adults.** Approve for 1 year if the patient meets the following (A, B, and C):
 - A)** Patient is ≥ 18 years of age; AND

B) Patient is concurrently receiving other medication therapy for depression [**documentation required**]; AND

Note: Examples of other medications for the treatment of depression include selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs).

C) Preferred product criteria is met for the products as listed in the below tables

5. Excessive Daytime Sleepiness Associated with Myotonic Dystrophy. Approve for 1 year if the patient meets both of the following (A and B):

A) Patient is ≥ 18 years of age; AND

B) Preferred product criteria is met for the products as listed in the below tables

6. Excessive Daytime Sleepiness Associated with Parkinson's Disease. Approve for 1 year if the patient meets both of the following (A and B):

A) Patient is ≥ 18 years of age; AND

B) Preferred product criteria is met for the products as listed in the below tables

7. Fatigue Associated with Multiple Sclerosis. Approve for 1 year if the patient meets both of the following (A and B):

A) Patient is ≥ 18 years of age; AND

B) Preferred product criteria is met for the products as listed in the below tables

8. Idiopathic Hypersomnia. Approve for 1 year if the patient meets the following (A, B, and C):

A) Patient is ≥ 18 years of age; AND

B) The diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center); AND

C) Preferred product criteria is met for the products as listed in the below tables

Employer Plans:

Product	Criteria
Nuvigil (armodafinil)	The patient has tried the bioequivalent generic product, armodafinil , AND cannot take due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which would result, per the prescriber, in a significant allergy or serious adverse reaction.
Provigil (modafinil)	The patient has tried the bioequivalent generic product, modafinil , AND cannot take due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which would result, per the prescriber, in a significant allergy or serious adverse reaction.

Individual and Family Plan:

Product	Criteria
Nuvigil (armodafinil)	The patient has tried the bioequivalent generic product, armodafinil , AND cannot take due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which would result, per the prescriber, in a significant allergy or serious adverse reaction.

Product	Criteria
Provigil (modafinil)	The patient has tried the bioequivalent generic product, modafinil , AND cannot take due to a formulation difference in the inactive ingredient(s) [e.g., difference in dyes, fillers, preservatives] between the Brand and the bioequivalent generic product which would result, per the prescriber, in a significant allergy or serious adverse reaction.

Conditions Not Covered

Modafinil (brand and generic) and armodafinil (brand and generic) for any other use is considered not medically necessary, including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- 1. Attention Deficit Hyperactivity Disorder (ADHD).** The American Academy of Pediatrics clinical practice guidelines for the treatment of ADHD in children and adolescents (2011 and 2019) do not address the use of modafinil/armodafinil.^{11,12} These guidelines note that with the greater availability of approved medications for children/adolescents with ADHD, it has become increasingly unlikely that clinicians need to consider the off-label use of other medications. Many options exist for the treatment of ADHD in adults (e.g., methylphenidate, dextroamphetamine) and further large scale trials that demonstrate benefit for modafinil in adults with ADHD are needed.
- 2. Bipolar Disorder, including Bipolar Depression.** Limited data (one small study [n = 85] and case reports [n = 2]) are available that describe the use of modafinil for bipolar disorder and bipolar depression.¹³⁻¹⁵ In one study (n = 257), armodafinil was not more effective than placebo in treating bipolar depression.¹⁶ Only limited data support modafinil for this condition and more data are needed.
- 3. Cancer-Related Fatigue.** The National Comprehensive Cancer Network guidelines on cancer-related fatigue (version 2.2025 – January 14, 2025) no longer consider modafinil or armodafinil to be effective for the treatment of cancer-related fatigue and recommend against its use.¹⁷
- 4. Chronic Fatigue Syndrome.** Limited data characterize modafinil therapy in those with chronic fatigue syndrome.¹⁸ In a randomized, double-blind, crossover study in 14 patients with chronic fatigue syndrome, use of modafinil for 20 days had minimal effects on cognitive function and no significant effects on fatigue, health-related quality of life, or mood.¹⁹ More data are required to assess efficacy in this patient population.
- 5. Excessive Daytime Sleepiness Associated with Primary Insomnia.** One randomized, placebo-controlled study found that neither combination therapy with modafinil and cognitive behavioral therapy nor modafinil as monotherapy significantly decreased daytime sleepiness associated with primary insomnia.²⁰
- 6. Enhancement of Performance in Situations of Induced Sleep Deprivation.** Studies are needed to define the role/appropriateness of modafinil in these situations for the general population (as opposed to military personnel, etc.). Studies have shown that modafinil may enhance performance and sustain alertness in individuals subjected to situations that deprive

sleep (e.g., military aviation, emergency physicians).²¹⁻²⁴ Further studies are needed before its use in the general population in these types of situations can be promoted.

7. **Fibromyalgia.** Limited data are available regarding the use of modafinil in fibromyalgia with most of the data being observational.²⁵⁻²⁷ Larger-sized, randomized, placebo-controlled trials are required to better assess and validate the efficacy of modafinil in patients with fibromyalgia before it can be recommended as a therapeutic modality.
8. **Hypersomnia, Fatigue or Sleepiness Due to Other Conditions (not Idiopathic Hypersomnia, see Other Uses with Supportive Evidence).** More data are needed in specific conditions to define the role of modafinil and armodafinil.
9. **Post-Stroke Sleep-Wake Disorders or Sleep Disorders.** Sleep-wake disorders occur in approximately 20% to 40% of patients who have experienced a stroke, which includes hypersomnia and excessive daytime sleepiness. Very limited data (i.e., case reports and one small study) have explored the use of modafinil in these patients to improve alertness.^{28,29} More data are needed to determine effectiveness in this condition.
10. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

References

1. Provigil® tablets [prescribing information]. Parsippany, NJ: Teva; December 2022.
2. Nuvigil® tablets [prescribing information]. Parsippany, NJ: Teva; December 2022.
3. National Institutes of Health. Narcolepsy. National Institute of Neurological Disorders and Stroke. Last reviewed on January 10, 2025. Available at: <https://www.ninds.nih.gov/health-information/disorders/narcolepsy?search-term=narcolepsy>. Accessed on August 28, 2025.
4. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2021;17(9):1881-1893.
5. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med*. 2021;17(9):1895-1945.
6. Patil SP, Ayappa IA, Caples SM, et al. Treatment of adult obstructive sleep apnea with positive airway pressure: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2019;15(2):335-343.
7. Gelenberg A, Freeman MP, Markowitz JC, et al. Practice guideline for the treatment of patients with major depressive disorder, Third edition. American Psychiatric Association, November 2010. Available at: <https://www.ninds.nih.gov/health-information/disorders/narcolepsy?search-term=narcolepsy>. Accessed on August 28, 2025.
8. Mayer G, Benesh H, Young P, et al. Modafinil in the treatment of idiopathic hypersomnia without long sleep time—a randomized, double-blind, placebo-controlled study. *J Sleep Res*. 2015;24(1):74-81.
9. Arnulf I, Thomas R, Roy A, et al. Update on the treatment of idiopathic hypersomnia: progress, challenges, and expert opinion. *Sleep Med Rev*. 2023;69:101766. .
10. Inoue Y, Tabata T, Tsukimori N. Efficacy and safety of modafinil in patients with idiopathic hypersomnia without long sleep time: a multicenter, randomized, double-blind, placebo-controlled, parallel-group comparison study. *Sleep Med*. 2021;80:315-321.
11. American Academy of Pediatrics. ADHD: Clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics*. 2011;128(5):1007-1022.

12. Wolraich ML, Hagan JF, Allan C, et al. AAP Subcommittee on children and adolescents with attention-deficit/hyperactivity disorder. Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. *Pediatrics*. 2019;144(4):e20192528.
13. Frye MA, Frunze H, Suppes T, et al. A placebo-controlled evaluation of adjunctive modafinil in the treatment of bipolar depression. *Am J Psychiatry*. 2007;164:1242-1249.
14. Post RM, Altshuler LL, Frye MA, et al. New findings from the bipolar collaborative network: clinical implications for therapeutics. *Curr Psychiatry Rep*. 2006;8:489-497.
15. Fernandes PP, Petty F. Modafinil for remitted bipolar depression with hypersomnia. *Ann Pharmacother*. 2003;37(12):1807-1809.
16. Calabrese JR, Ketter RA, Youakim JM, et al. Adjunctive armodafinil for major depressive episodes associated with bipolar I disorder: a randomized, multicenter, double-blind, placebo-controlled, proof-of-concept study. *J Clin Psychiatry*. 2010;71:1363-1370.
17. The NCCN Cancer-Related Fatigue Clinical Practice Guidelines in Oncology (Version 2.2025 – January 14, 2025). © 2025 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 28, 2025.
18. Turkington D, Hedwat D, Rider I, Young AH. Recovery from chronic fatigue syndrome with modafinil. *Hum Psychopharmacol*. 2004;19(1):63-64.
19. Randall DC, Cafferty FH, Shneerson JM, et al. Chronic treatment with modafinil may not be beneficial in patients with chronic fatigue syndrome. *J Psychopharmacol*. 2005;19(6):647-660.
20. Perlis ML, Smith MT, Orff H, et al. The effects of modafinil and cognitive behavior therapy on sleep continuity in patients with primary insomnia. *Sleep*. 2004;27(4):715-725.
21. Gill M, Haerich P, Westcott K, et al. Cognitive performance following modafinil versus placebo in sleep-deprived emergency physicians: a double-blind, randomized crossover study. *Acad Emerg Med*. 2006;13:158-165.
22. Caldwell JA, Caldwell JL, Smythe NK, Hall KK. A double-blind, placebo-controlled investigation of the efficacy of modafinil for sustaining the alertness and performance of aviators: a helicopter simulator study. *Psychopharmacol*. 2000;150:272-282.
23. Bonnet MH, Balkin TJ, Dinges DF, et al. The use of stimulants to modify performance during sleep loss: a review by the sleep deprivation and stimulant task force of the American Academy of Sleep Medicine. *Sleep*. 2005;28(9):1163-1187.
24. Caldwell JA, Caldwell JL. Fatigue in military aviation: an overview of US military-approved pharmacological countermeasures. *Aviat Space Environ Med*. 2005;76(7):C39-C51.
25. Schaller JL, Behar D. Modafinil in fibromyalgia treatment. *J Neuropsychiatry Clin Neurosci*. 2001;13(4):530-531.
26. Schwartz TL, Rayancha S, Rashid A, et al. Modafinil treatment for fatigue associated with fibromyalgia. *J Clin Rheumatol*. 2007;13(1):52.
27. Pachas WN. Modafinil for the treatment of fatigue of fibromyalgia. *J Clin Rheumatol*. 2003;9(4):282-285.
28. Bassetti CL. Sleep and stroke. *Semin Neurol*. 2005;25(1):19-32.
29. Bivard A, Lillicrap T, Krishnamurthy V, et al. MIDAS (modafinil in debilitating fatigue after stroke): a randomized, double-blind, placebo-controlled, cross-over trial. *Stroke*. 2017;48(5):1293-1298.

Revision Details

Type of Revision	Summary of Changes	Date
Annual Revision	Updated title from 'Armodafinil/Modafinil' to 'Wakefulness-Promoting Agents – Armodafinil, Modafinil'	10/15/2024

	Excessive Daytime Sleepiness Associated with Narcolepsy. Updated 'Treatment of Excessive Daytime Sleepiness Associated with Narcolepsy (Type 1 or 2)' TO 'Excessive Daytime Sleepiness Associated with Narcolepsy.' Removed 'Daily periods of irrepressible need to sleep or lapses into sleep during waking hours, occurring for at least three months' Updated 'Documentation of ONE of the following: (i) Diagnosis of narcolepsy type 1 and ONE of the following: (a) Mean Sleep Latency Test (MSLT) performed according to standard techniques, showing a mean sleep latency of less than or equal to 8 minutes <u>and</u> two or more sleep-onset rapid eye movement periods (SOREMPs) following a nocturnal polysomnogram (PSG) that rules out other causes of excessive daytime sleepiness, (b) A SOREMP (within 15 minutes of sleep onset) on a nocturnal PSG; (ii) Diagnosis of narcolepsy type 2 <u>and</u> Mean Sleep Latency Test (MSLT) performed according to standard techniques, showing a mean sleep latency of less than or equal to 8 minutes <u>and</u> two or more sleep-onset rapid eye movement periods (SOREMPs) following a nocturnal polysomnogram (PSG) that rules out other causes of excessive daytime sleepiness. A SOREMP (within 15 minutes of sleep onset) on a nocturnal PSG may replace one of the SOREMPs on the MSLT' TO 'Patient has been evaluated using polysomnography and a multiple sleep latency test' Removed 'The hypersomnolence and/or MSLT findings are not better explained by other causes such as insufficient sleep, delayed sleep phase disorder, or the effect of medication or substances or their withdrawal' Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Added 'Diagnosis of narcolepsy has been confirmed, according to the prescriber' Excessive Daytime Sleepiness Associated with Obstructive Sleep Apnea/Hypoapnea Syndrome. Removed 'Daily periods of irrepressible need to sleep or lapses into sleep during waking hours, occurring for at least three months' Removed 'Documentation of diagnosis of Obstructive Sleep Apnea (OSA)/Hypoapnea Syndrome (OSAHS) is confirmed by sleep study' Removed 'The hypersomnolence and/or sleep study findings are not better explained by other	
--	--	--

	causes such as insufficient sleep, delayed sleep phase disorder, or the effect of medication or substances or their withdrawal' Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Removed 'Documentation of inadequate response to at least 1 month of non-pharmacologic treatment for OSA (for example, continuous positive airway pressure [CPAP])' Removed 'Armodafinil (Nuvigil) or modafinil (Provigil) will be used in combination with non-pharmacologic treatment for OSA/OSAHS, unless contraindicated or intolerant' Added 'Patient meets one of the following (i or ii): (i)Armodafinil/modafinil will be used in conjunction with continuous positive airway pressure therapy; OR (ii) Patient is unable to initiate or tolerate continuous positive airway pressure therapy;" Excessive Sleepiness Associated with Shift Work Sleep Disorder. Removed 'Documentation of insomnia and/or excessive sleepiness, accompanied by a reduction of total sleep time, which is associated with a recurring work schedule that overlaps the usual time for sleep' Removed 'Documentation of sleep log, completed on work and free days, demonstrating a disturbed sleep and wake pattern' Removed 'Documentation that the sleep and/or wake disturbance cannot be better explained by another cause (for example, concurrent sleep disorder, medical or neurological disorder, mental disorder, medication use, poor sleep hygiene, substance use disorder)' Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Adjunctive/Augmentation Treatment for Depression in Adults. Removed 'Medication is prescribed by, or in consultation with, a neurologist or psychiatrist" Excessive Daytime Sleepiness Associated with Myotonic Dystrophy. Removed 'Daily periods of irrepressible need to sleep or daytime lapses into sleep occurring for at least three months' Removed 'Medication is prescribed by, or in consultation with, a neurologist, pulmonologist, or sleep specialist'	
--	--	--

	Excessive Daytime Sleepiness Associated with Parkinson's Disease. Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Fatigue Associated with Multiple Sclerosis. Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Idiopathic Hypersomnia. Removed pulmonologist from 'Medication is prescribed by, or in consultation with' bullet Removed 'Daily periods of irrepressible need to sleep or lapses into sleep during waking hours, occurring for at least three months' Removed 'Documentation of Multiple Sleep Latency Test (MSLT) performed according to standard techniques demonstrating an average sleep latency of less than or equal to 8 minutes with a total of less than 2 sleep onset rapid eye movement periods (SOREMPs)' Removed 'Documented absence of cataplexy' Removed 'The hypersomnolence and/or MSLT findings are not better explained by other causes such as insufficient sleep, delayed sleep phase disorder, or the effect of medication or substances or their withdrawal'	
Selected Revision	Added "Documentation: Documentation is required where noted in the criteria. Documentation may include, but not limited to, chart notes, laboratory tests, claims records, and/or other information." Excessive Daytime Sleepiness Associated with Narcolepsy. Updated criteria from "Patient has been evaluated using polysomnography and a multiple sleep latency test" to "Documentation that the patient has been evaluated using polysomnography and a multiple sleep latency test." Updated criteria from "Diagnosis of narcolepsy has been confirmed, according to the prescriber" to "Documented diagnosis of narcolepsy has been confirmed." Excessive Daytime Sleepiness Associated with Obstructive Sleep Apnea/Hypoapnea Syndrome. Updated criteria from "Patient is unable to initiate or tolerate continuous positive airway pressure therapy" to "Documentation that the patient is unable to initiate or tolerate continuous positive airway pressure therapy."	3/1/2025

	Excessive Sleepiness Associated with Shift Work Sleep Disorder. Updated criteria from "Patient works at least five overnight shifts per month" to "Documentation that the patient works at least five overnight shifts per month." Adjunctive/Augmentation Treatment for Depression in Adults. Updated criteria from "Patient is concurrently receiving other medication therapy for depression" to "Documentation that the patient is concurrently receiving other medication therapy for depression." Idiopathic Hypersomnia. Updated criteria from "The diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)" to "Documented diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)."	
Annual Revision	Excessive Daytime Sleepiness Associated with Obstructive Sleep Apnea. Criterion requiring use in conjunction with continuous positive airway pressure or patient is unable to initiate or tolerate continuous positive airway pressure was changed to use in conjunction with positive airway pressure or patient is unable to initiate or tolerate positive airway pressure with a Note that positive airway pressure can include continuous positive airway pressure (CPAP), auto-titrating positive airway pressure (APAP), or bilevel positive airway pressure (BPAP). Updated documentation verbiage to "[documentation required]." Idiopathic Hypersomnia. Removed documentation requirement from "Documented diagnosis is confirmed by a sleep specialist physician or at an institution that specializes in sleep disorders (i.e., sleep center)."	11/15/2025

The policy effective date is in force until updated or retired.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.