

Opioid Therapy – Individual and Family Plans

Table of Contents

Overview	1
Medical Necessity Criteria	1
Reauthorization Criteria	9
Authorization Duration	9
Background	9
References	13
Revision History	13

Related Coverage Resources

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Each coverage request should be reviewed on its own merits. Medical directors are expected to exercise clinical judgment where appropriate and have discretion in making individual coverage determinations. Where coverage for care or services does not depend on specific circumstances, reimbursement will only be provided if a requested service(s) is submitted in accordance with the relevant criteria outlined in the applicable Coverage Policy, including covered diagnosis and/or procedure code(s). Reimbursement is not allowed for services when billed for conditions or diagnoses that are not covered under this Coverage Policy (see "Coding Information" below). When billing, providers must use the most appropriate codes as of the effective date of the submission. Claims submitted for services that are not accompanied by covered code(s) under the applicable Coverage Policy will be denied as not covered. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This policy supports medical necessity review for ALL of the following:

- Immediate-release opioid analgesics (refer to [Appendix 1](#) for products)
- Extended-release opioid analgesics (refer to [Appendix 2](#) for products)
- Injectable opioid analgesics (refer to [Appendix 3](#) for products)

Methadone for the treatment of Opioid Use Disorder (OUD) is covered without prior authorization as part of an opioid treatment program

Medical Necessity Criteria

Opioid analgesics are considered medically necessary when the following are met:

- I. **Immediate-release opioid analgesics in an opioid naïve individual are considered medically necessary when the individual meets ONE of the following:**
 1. Documented diagnosis of active cancer treatment (defined as receiving antineoplastic or antitumor therapy) requiring treatment for cancer-related pain, end-of-life care (including hospice or palliative care), **OR** sickle cell disease
 1. Preferred product criteria is met for the product(s) as listed in the below table
 2. **For the management of acute pain (for example, pain lasting less than 90 days).** Individual meets **ALL** of the following:
 - A. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - B. **ONE** of the following:
 - i. For the management of acute dental pain: Attestation that an initial treatment regimen exceeding 3 days is medically necessary (for example, the individual is not a candidate for less than 3 days of therapy)
 - ii. For the management of acute non-dental pain: Attestation that an initial treatment regimen exceeding 7 days is medically necessary (for example, the individual is not a candidate for less than 7 days of therapy)
 - C. Preferred product criteria is met for the product(s) as listed in the below table
 3. **For the management chronic pain (for example, pain lasting more than 90 days).** Individual meets **ALL** of the following:
 - A. Failure, contraindication, or intolerance to nonopioid pharmacologic therapies intended to treat pain
 - B. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - C. Attestation that opioids will be prescribed in accordance with current clinical practice guidelines **AND** that an assessment of risks, harms, and goals consistent with an opioid agreement (or comprehensive treatment plan) has been undertaken
 - D. Preferred product criteria is met for the product(s) as listed in the below table

Immediate-Release Opioid Analgesic Products and Criteria:

Product	Criteria
Acetaminophen/caffeine/dihydrocodeine	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of failure, contraindication, or intolerance to BOTH of the following short acting narcotics: A. hydrocodone / acetaminophen B. oxycodone / acetaminophen (generic for Percocet)
Apadaz™ (benzhydrocodone / acetaminophen) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of failure, contraindication, or intolerance to BOTH of the following short acting narcotics: A. hydrocodone / acetaminophen B. oxycodone / acetaminophen (generic for Percocet)
Dilaudid® (hydromorphone) oral solution	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of hydromorphone oral solution (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive

Product	Criteria
	ingredient(s) which would result in a significant allergy or serious adverse reaction B. Failure, contraindication, or intolerance to FOUR of the following short acting narcotics: i. hydrocodone / acetaminophen ii. morphine iii. oxycodone iv. oxycodone / acetaminophen (generic for Percocet) v. oxymorphone
Dilaudid® (hydromorphone) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of hydromorphone tablet (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction B. Failure, contraindication, or intolerance to FOUR of the following short acting narcotics: i. hydrocodone / acetaminophen ii. morphine iii. oxycodone iv. oxycodone / acetaminophen (generic for Percocet) v. oxymorphone
Hycodan® (hydrocodone / homatropine) oral solution	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documented trial of hydrocodone/homatropine oral solution (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction
Hycodan® (hydrocodone / homatropine) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documented trial of hydrocodone/homatropine tablet (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction
Nalocet® (oxycodone / acetaminophen) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Intolerance to oxycodone / acetaminophen (generic for Percocet) B. Failure, contraindication, or intolerance to hydrocodone / acetaminophen
Nucynta® (tapentadol) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of Failure, contraindication, or intolerance to ALL of the following:

Product	Criteria
	A. hydrocodone / acetaminophen B. morphine C. oxycodone D. oxycodone / acetaminophen (generic for Percocet) E. oxymorphone
Oxaydo® (oxycodone tablet)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Intolerance to oxycodone B. Failure, contraindication, or intolerance to ALL of the following short acting narcotics: i. hydrocodone / acetaminophen ii. hydromorphone iii. morphine iv. oxymorphone
Oxycodone tablets (generic for RoxyBond, manufactured by Ohemo Life Sciences)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of ONE of the following: A. An abuse-deterrent short-acting opioid is required B. The patient has tried the oxycodone tablets (generic for Roxicodone) AND cannot take due to a formulation difference in the inactive ingredient(s) [for example, difference in dyes, fillers, preservatives] between oxycodone tablets (generic for RoxyBond) and oxycodone tablets (generic for Roxicodone), which, per the prescriber, would result in a significant allergy or serious adverse reaction.
Oxycodone / Acetaminophen 2.5 mg/ 300 mg, 5 mg/ 300 mg, 7.5 mg/ 300 mg, 10 mg/ 300 mg tablets and capsules	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Intolerance to generic oxycodone / acetaminophen tablets (generic for Percocet) B. Failure, contraindication or intolerance to hydrocodone / acetaminophen
Oxycodone / Acetaminophen 5 mg / 325 mg oral solution	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of ONE of the following: A. Intolerance to generic oxycodone / acetaminophen tablets (generic for Percocet) B. Inability to swallow tablets
Percocet® (oxycodone / acetaminophen) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of oxycodone/acetaminophen tablet (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction

Product	Criteria
	B. Failure, contraindication, or intolerance to hydrocodone / acetaminophen
Prolate® (oxycodone/ acetaminophen) oral solution	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of ONE of the following: A. Intolerance to generic oxycodone / acetaminophen tablets (generic for Percocet) B. Inability to swallow tablets
Qdolo® (tramadol) oral solution	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. 18 years of age or older B. Inability to swallow tramadol 50 mg immediate-release tablets
Tramadol oral solution (generic for Qdolo)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. 18 years of age or older B. Inability to swallow tramadol 50 mg immediate-release tablets
Roxicodone® (oxycodone)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of oxycodone tablet (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction B. Failure, contraindication, or intolerance to FOUR of the following short acting narcotics: i. hydrocodone / acetaminophen ii. hydromorphone iii. morphine iv. oxycodone / acetaminophen (generic for Percocet) v. oxymorphone
RoxyBond™ (oxycodone)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of ONE of the following: A. An abuse-deterrent short-acting opioid is required B. The patient has tried the oxycodone tablets (generic for Roxicodone) AND cannot take due to a formulation difference in the inactive ingredient(s) [for example, difference in dyes, fillers, preservatives] between RoxyBond and oxycodone tablets (generic for Roxicodone), which, per the prescriber, would result in a significant allergy or serious adverse reaction.

Product	Criteria
Seglantis® (celecoxib and tramadol)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. 18 years of age or older B. Inability to use tramadol tablets and celecoxib capsules concurrently
Tramadol 25 mg tablet	In addition to the criteria detailed above, the following: 1. Inability to obtain prescribed dose with tramadol 50 mg immediate-release tablets
Tramadol 75 mg tablet	In addition to the criteria detailed above, the following: 1. The patient's prescribed dose cannot be obtained with tramadol 50 mg. <u>Note:</u> The patient is NOT required to split the 50 mg tablets in half.
Tramadol 100 mg tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of intolerance or inability to use tramadol 50 mg immediate-release tablets
Trezix™ (dihydrocodeine / acetaminophen / caffeine) tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Intolerance to generic dihydrocodeine / acetaminophen / caffeine capsules B. Failure, contraindication, or intolerance to BOTH of the following short acting narcotics: i. hydrocodone / acetaminophen ii. oxycodone / acetaminophen (generic for Percocet)

II. **Extended-release opioid analgesics are considered medically necessary when the individual meets ONE of the following:**

1. Documented diagnosis of active cancer treatment (defined as receiving antineoplastic or antitumor therapy) requiring treatment for cancer-related pain, end-of-life care (including hospice or palliative care), **OR** sickle cell disease
Preferred product criteria is met for the product(s) as listed in the below table
2. Diagnosis of pain severe enough to require daily, around-the-clock, long-term opioid treatment. Individual meets **ALL** of the following:
 - A. Failure, contraindication, or intolerance to a minimum one week trial of immediate-release opioids
 - B. Failure, contraindication, or intolerance to nonopioid pharmacologic therapies intended to treat pain
 - C. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - D. Documentation that opioids will be prescribed in accordance with current clinical practice guidelines **AND** that an assessment of risks, harms, and goals consistent with an opioid agreement (or comprehensive treatment plan) has been undertaken
 - E. Preferred product criteria is met for the product(s) as listed in the below table

Extended-Release Opioid Analgesic Products and Criteria:

Product	Criteria
Conzip™ (tramadol extended-release capsule)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of failure, contraindication, or intolerance to tramadol 50 mg tablets
Tramadol extended-release capsule (generic for Conzip)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of failure, contraindication, or intolerance to tramadol 50 mg tablets
Hysingla™ ER (hydrocodone bitartrate)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of hydrocodone bitartrate ER (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction B. Failure, contraindication, or intolerance to Xtampza ER
MS Contin® (morphine sulfate) extended-release tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of BOTH of the following: A. Trial of morphine sulfate ER (the bioequivalent generic product) AND cannot take due to a formulation difference in the inactive ingredient(s) which would result in a significant allergy or serious adverse reaction B. Failure, contraindication, or intolerance to ALL of the following: i. hydrocodone bitartrate ER ii. hydromorphone ER iii. oxymorphone ER iv. Xtampza ER
Nucynta® ER (tapentadol)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. Documentation of failure, contraindication, or intolerance to BOTH of the following: A. hydrocodone bitartrate ER B. Xtampza ER
Oxycontin® (oxycodone HCl)	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment 2. 11 years of age to 17 years of age 3. Documentation of failure, contraindication, or intolerance to BOTH of the following: A. hydrocodone bitartrate ER B. Xtampza ER
Oxycodone extended-release tablet	In addition to the criteria detailed above, ONE of the following: 1. Established therapy for an individual in hospice or end of life care or active cancer treatment

Product	Criteria
	2. 11 years of age to 17 years of age 3. Documentation of failure, contraindication, or intolerance to BOTH of the following: A. hydrocodone bitartrate ER B. Xtampza ER

III. Methadone and Fentanyl Transdermal Patches are considered medically necessary when the individual meets ONE of the following:

1. Documented diagnosis of active cancer treatment (defined as receiving antineoplastic or antitumor therapy) requiring treatment for cancer-related pain, end-of-life care (including hospice or palliative care), **OR** sickle cell disease
2. **ALL** of the following:
 - A. Diagnosis of pain severe enough to require daily, around-the-clock, long-term opioid treatment
 - B. Failure, contraindication, or intolerance to a minimum one week trial of immediate-release opioids
 - C. Opioid tolerant [required daily dosage for pain management exceeds 60 morphine milligram equivalents (MME)]
 - D. Alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain
 - E. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - F. Documentation that opioids will be prescribed in accordance with current clinical practice guidelines **AND** that an assessment of risks, harms, and goals consistent with an opioid agreement (or comprehensive treatment plan) has been undertaken
 - G. Medication prescribed by, or in consultation with, a board certified pain management specialist

IV. Daily dose of all opioid analgesics exceed 120 morphine milligram equivalents (120 MME), opioid analgesics are considered medically necessary when the individual meets ONE of the following:

1. Documented diagnosis of active cancer treatment (defined as receiving antineoplastic or antitumor therapy) requiring treatment for cancer-related pain, end-of-life care (including hospice or palliative care), **OR** sickle cell disease
2. **ALL** of the following:
 - A. Failure, contraindication, or intolerance to nonopioid pharmacologic therapies intended to treat pain
 - B. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - C. Documentation that opioids will be prescribed in accordance with current clinical practice guidelines **AND** that an assessment of risks, harms, and goals consistent with an opioid agreement (or comprehensive treatment plan) has been undertaken
 - D. Consideration of additional precautions intended to reduce the risk of serious harm associated with high dose opioids (for example, education and provision of naloxone)
 - E. Quarterly reassessment of opioid therapy benefits and risks specific to diagnosis and treatment goals
 - F. Medication prescribed by, or in consultation with, a board certified pain management specialist

- V. **Daily dose of all opioid analgesics exceed 200 morphine milligram equivalents (200 MME), opioid analgesics are considered medically necessary when the individual meets ONE of the following:**
1. Documented diagnosis of active cancer treatment (defined as receiving antineoplastic or antitumor therapy) requiring treatment for cancer-related pain, end-of-life care (including hospice or palliative care), **OR** sickle cell disease
 2. **ALL** of the following:
 - A. Failure, contraindication, or intolerance to nonopioid pharmacologic therapies intended to treat pain
 - B. Attestation that the history of controlled substance prescriptions has been checked using the state prescription drug monitoring program (PDMP)
 - C. Documentation that opioids will be prescribed in accordance with current clinical practice guidelines **AND** that an assessment of risks, harms, and goals consistent with an opioid agreement (or comprehensive treatment plan) has been undertaken
 - D. Quarterly reassessment of opioid therapy benefits and risks specific to diagnosis and treatment goals
 - E. Consideration of additional precautions intended to reduce the risk of serious harm associated with high dose opioids (for example, education and provision of naloxone)
 - F. The provider has performed an individualized behavioral health screening to assess the risks and benefits of the opioid dose (for example, PHQ-9, GAD-7, PC-PTSD)
 - G. The provider has screened for substance abuse risk to assess the risks and benefits of the opioid dose (for example, DIRE, ORT, PDUQ, PMQ)
 - H. Medication prescribed by, or in consultation with, a board certified pain management specialist

Resources for calculating morphine milligram equivalents can be found in [Appendix 4](#)

When coverage is available and medically necessary, the dosage, frequency, duration of therapy, and site of care should be reasonable, clinically appropriate, and supported by evidence-based literature and adjusted based upon severity, alternative available treatments, and previous response to therapy.

Receipt of sample product does not satisfy any criteria requirements for coverage.

Reauthorization Criteria

Opioid analgesics are considered medically necessary for continued use when the above medical necessity criteria are met **AND** there is documentation of beneficial response.

Authorization Duration

Initial approval duration is up to 12 months
Reauthorization approval duration is up to 12 months

Background

OVERVIEW

All of the long-acting (LA) opioids are indicated for the **management of severe and persistent pain that requires an extended treatment period with a daily opioid analgesic** and for which alternative treatment options are inadequate.¹⁻⁹ OxyContin is the only product specifically indicated in pediatric patients 11 years to 18 years of age.³ Nucynta ER is the only product also indicated for the management of neuropathic pain associated with diabetic peripheral neuropathy in adults.¹

The currently available LA opioids are buprenorphine, fentanyl, hydrocodone, hydromorphone, morphine sulfate, oxycodone, oxymorphone, tapentadol, and tramadol.¹⁻⁹

Guidelines

In 2022, the **Centers for Disease Control and Prevention (CDC)** published an updated guideline for prescribing opioids for pain.¹⁰ Nonopioid therapies are at least as effective as opioids for many common types of acute pain, and nonopioid therapies are preferred for subacute and chronic pain. Clinicians should maximize the use of nonpharmacologic and nonopioid pharmacologic therapies as appropriate for the specific condition and patient and only consider initiating opioid therapy if expected benefits for pain and function are anticipated to outweigh risks to the patient. Multiple noninvasive nonpharmacologic interventions (e.g., aerobic, aquatic, or resistance exercises, weight loss, psychological therapy, spinal manipulation, low-level laser therapy, massage, mindfulness-based stress reduction, yoga, tai chi, qigong, acupuncture, cognitive behavioral therapy, and spinal manipulation) are associated with improvements in pain, function, or both, that are sustained after treatment and are not associated with serious harms. Non-opioid drugs (e.g., tricyclic antidepressants, serotonin and norepinephrine reuptake inhibitor [SNRI] antidepressants, duloxetine, selected antiseizure medications [e.g., pregabalin, gabapentin, oxcarbazepine], capsaicin and lidocaine patches, and nonsteroidal anti-inflammatory drugs [NSAIDs]) are associated with small to moderate improvements in chronic pain and function for certain chronic pain conditions.

Before initiating opioid therapy for patients with pain, clinicians should discuss with patients the realistic benefits and known risks of opioid therapy.¹⁰ Before starting ongoing opioid therapy for patients with subacute or chronic pain, clinicians should work with patients to establish treatment goals for pain and function and consider how opioid therapy will be discontinued if benefits do not outweigh risks. When opioids are initiated, clinicians should prescribe the lowest effective dosage of immediate-release opioids for no longer than needed for the expected duration of pain severe enough to require opioids. During ongoing opioid therapy, clinicians should collaborate with patients to evaluate and carefully weigh the benefits and risks of continuing opioid therapy and exercise care when increasing, continuing, or reducing opioid dosage. The guideline recommends that clinicians should not initiate opioid treatment with LA opioids for patients who are opioid-naïve and should not prescribe LA opioids for intermittent use. LA opioids should be reserved for severe, continuous pain. Before starting and periodically during continuation of opioid therapy, clinicians should evaluate the risk for opioid-related harms and should work with patients to incorporate relevant strategies to mitigate risk, including offering naloxone and reviewing potential interactions with any other prescribed medications or substances used. When prescribing initial opioid therapy and periodically during opioid therapy, clinicians should review the patient's history of controlled substance prescriptions using state prescription drug monitoring program (PDMP) data to determine whether the patient is receiving opioid dosages or combinations that put the patient at high risk for overdose. When prescribing opioids for subacute or chronic pain, clinicians should consider the benefits and risks of toxicology testing to assess for prescribed medications as well as other prescribed and nonprescribed controlled substances.

The 2020 **American Society of Hematology** guideline for the management of acute and chronic pain in patients with sickle cell disease states that pain causes significant morbidity for those living with sickle cell disease and manifests as acute intermittent pain, chronic daily pain, and acute-on-chronic pain.¹¹ For adults and children with chronic pain who are receiving chronic opioid therapy, are functioning well, and have perceived benefit, the guideline suggests shared decision making for continuation of chronic opioid therapy. For adults and children with chronic pain who are receiving chronic opioid therapy, are functioning poorly, or are at high risk for aberrant opioid use or toxicity, the guideline suggests against continuation of chronic opioid therapy.

Appendix 1

Immediate-release opioid analgesics include the following:

Acetaminophen/codeine tablet
Benzhydrocodone/acetaminophen (Apadaz™)
Butorphanol tartrate spray
Codeine sulfate tablets and solution
Codeine/carisoprodol/aspirin
Dihydrocodeine/acetaminophen/caffeine (Dvorah, Trezix™)
Dihydrocodeine/aspirin/caffeine
Hydrocodone/acetaminophen (Lorcet®, Lorcet® HD, Lorcet® Plus, Lortab®, Stagesic-10, Zydone®)

Hydrocodone/ibuprofen
Hydromorphone tablets, solution, suppositories (Dilaudid®)
Levorphanol tablets
Meperidine tablets and solution
Morphine sulfate tablets, solution, suppositories
Nucynta® (tapentadol)
Opium Tincture
Opium/Belladonna alkaloids (Belladonna-Opium) suppository
Oxycodone hydrochloride capsules, concentrate, solution, tablets (Oxaydo®)
Oxycodone/acetaminophen (Endocet, Nalocet, Percocet®, Primlev™; Prolate)
Oxycodone/aspirin tablets
Oxycodone/ibuprofen tablets
Oxymorphone tablets
Pentazocine/naloxone
Roxicodone® (oxycodone) tablets
RoxyBond™ (oxycodone) tablets
Tramadol (Qdolo and its Authorized Generic) solution, (Ultram®) tablets
Tramadol/acetaminophen (Ultracet®) tablets

Appendix 2

Extended-release opioid analgesics include the following:

Conzip™ (tramadol ER capsules)
Fentanyl patches
Hydrocodone bitartrate ER (generic for Hysingla ER)
Hydrocodone bitartrate ER (generic for Zohydro ER)
Hydromorphone hydrochloride ER tablets
Hysingla® ER (hydrocodone bitartrate)
Levorphanol tablets
Morphabond ER™ (morphine sulfate)
MS Contin® (morphine sulfate)
Nucynta® ER (tapentadol)
Oxycodone hydrochloride ER tablets
Oxycontin® (oxycodone HCl)
Oxymorphone hydrochloride ER tablets
Tramadol hydrochloride ER capsules/tablets
Xtampza® ER (oxycodone myristate)

Appendix 3

Alfentanil ampule
Hydromorphone ampule
Hydromorphone carpuject
Hydromorphone Isecure syringe
Hydromorphone syringe (Dilaudid)
Hydromorphone vial
Hydromorphone / Bupivacane
Hydromorphone / Ropivacaine / Sodium chloride
Hydromorphone / Sodium chloride
Hydromorphone / Water
Meperidine ampule (Demerol)
Meperidine carpuject (Demerol)
Meperidine vial (Demerol)
Meperidine / Sodium chloride
Methadone vial
Morphine ampule (Duramorph, Infumorph)

Morphine auto-injector
Morphine carpject
Morphine Isecure syringe
Morphine syringe
Morphine vial (Mitgo)
Morphine / Sodium chloride
Morphine / D5W
Remifentanil vial (Ultiva)
Sufentanil ampule
Sufentanil vial

Appendix 4

Morphine Milligram Equivalent (MME) Dose Calculation

A commonly used method to compare opioid analgesic doses is to calculate its dose in terms of morphine milligram equivalents. To calculate the morphine milligram equivalent (MME) dose one starts by adding the total daily milligram amount of each opioid an individual takes. Next, each opioid total dose should be converted to a morphine milligram equivalent by multiplying the total dose by the conversion factor associated with the identity of the patients' opioid. The conversion factors are shown in the table below. Once all opioid doses have been converted to a morphine milligram equivalent dose, they should be added together to calculate a total MME dose per day. It is important to note that this calculation should only be used as an approximation for comparative purposes. If using a similar conversion to switch an individual to a different opioid, a dose reduction must be made to account for incomplete tolerance and prevent an overdose.

Opioid	Conversion Factor
Codeine	0.15
Fentanyl (transdermal) in mcg / hr	2.4
Hydrocodone	1
Hydromorphone	4
Methadone	
1 – 20 mg / day	4
21 – 40 mg / day	8
41 – 60 mg / day	10
≥ 61 – 80 mg / day	12
Morphine	1
Oxycodone Hydrochloride	1.5
Oxycodone Myristate (Xtampza ER)	1.67
Oxymorphone	3
Tapentadol	0.4

Source: Adapted from CDC Guideline for Prescribing Opioids for Chronic Pain

Example: An individual is using 20 mg of extended-release oxycodone twice daily and 5 mg / 300 mg of immediate-release hydrocodone / acetaminophen three times daily.

- 1) Calculate the total daily dose of each opioid
 - a. 20 mg oxycodone twice daily = 40 mg oxycodone
 - b. 5 mg hydrocodone three times daily = 15 mg hydrocodone
- 2) Convert to a morphine milligram equivalent dose by multiplying the total daily dose of each opioid by the conversion factor
 - a. 40 mg oxycodone x 1.5 = 60 morphine milligram equivalents
 - b. 15 mg hydrocodone x 1 = 15 morphine milligram equivalents
- 3) Add the morphine milligram equivalent doses to calculate the total morphine milligram equivalent dose per day
 - a. 60 + 15 = 75 morphine milligram equivalents per day

References

1. Nucynta® ER extended-release oral tablets [prescribing information]. Stoughton, MA: Collegium; March 2021.
2. Embeda® extended-release capsules [prescribing information]. New York, NY: Pfizer; October 2019.
3. Avinza® capsules [prescribing information]. New York, NY: Pfizer; May 2014.
4. MS Contin® tablets [prescribing information]. Stamford, CT: Purdue; March 2021.
5. OxyContin® tablets [prescribing information]. Stamford, CT: Purdue; October 2021.
6. Oxymorphone ER tablets [prescribing information]. Bridgewater, NJ: Amneal; June 2022.
7. Exalgo® extended-release tablets [prescribing information]. Webster Groves, MO: SpecGx; October 2019.
8. Zohydro® ER extended-release capsules [prescribing information]. Morristown, NJ: Currax; March 2021.
9. Hysingla™ ER extended-release tablets [prescribing information]. Stamford, CT: Purdue; March 2021.
10. Xtampza ER® extended-release capsules [prescribing information]. Cincinnati, OH: Patheon; March 2021.
11. Arymo® ER extended-release tablets [prescribing information]. Wayne, PA: Egalet; October 2019.
12. Conzip® extended-release capsules [prescribing information]. Bridgewater, NJ: Vertical; September 2021.
13. Belbuca® buccal film [prescribing information]. Raleigh, NC: BioDelivery Sciences; June 2022.
14. Duragesic® transdermal system [prescribing information]. Titusville, NJ: Janssen; March 2021.
15. Dolophine® [prescribing information]. Eatontown, NJ: West-Ward; June 2021.
16. Dowell D, Ragan KR, Jones CM, et al. CDC Clinical Practice Guideline for Prescribing Opioids for Pain - United States, 2022. *MMWR Recomm Rep.* 2022; 71(3):1-95.
17. Brandow AM, Carroll CP, Creary S, et al. American Society of Hematology 2020 guidelines for sickle cell disease: management of acute and chronic pain. *Blood Adv.* 2020; 4(12):2656-2701.

Revision History

Type of Revision	Summary of Changes	Date
Selected Revision	Preferred Product Table: Added preferred product criteria for Tramadol 75mg tablet.	04/01/2025
Selected Revision	Appendix 2 Removed Kadian from the appendix.	06/01/2025
Selected Revision	Updated policy with clarification regarding Methadone for the treatment of Opioid Use Disorder (OUD)	9/15/2025

The policy effective date is in force until updated or retired.

"Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group. © 2025 The Cigna Group.